

Effect of different doses of nitrogen and potassium on growth and yield of potato (*Solanum tuberosum* L.) under New Alluvial Zone of West Bengal

Abstract

An experiment was conducted during *rabi* season of 2012-13 and 2013-14 at Instructional Farm, Jaguli, Nadia, West Bengal to find out the different doses of nitrogen and potassium on growth and yield of potato. The experiment was designed in RBD with 10 treatments replicated thrice in potato cultivated variety *Kufri Jyoti*. Different doses of nitrogen and potassium were considered as the treatments. For all the treatments P_2O_5 dose was 150 kg ha^{-1} . The size of the experimental plots were 12 square meter and seed tubers were planted with 50 cm X 20 cm spacing. In this experiment, it was observed that the growth attributes like plant height, leaf area index, dry matter accumulation of tubers at 80 DAP, crop growth rate at 60-80 DAP were highest with the application of $250\text{ kg ha}^{-1}\text{ N}$, $200\text{ kg K}_2\text{O}$ (T_9) and statistically at par with $300\text{ kg ha}^{-1}\text{ N}$, $150\text{ kg ha}^{-1}\text{ K}_2\text{O}$ (T_{10}). Again among the yield parameters T_9 recorded the highest tuber number per square meter and tuber yield which was closely followed by T_{10} . Highest B:C ratio was also observed in T_9 . This result proves that T_9 can be recommended to get better growth and economic yield of potato than T_{10} (farmer practice dose) in the new alluvial soil of West Bengal.

Key words: potato, different doses of nitrogen and potassium, tuber growth and yield

INTRODUCTION

Worldwide one of the most important food crops is potato (*Solanum tuberosum* L.). After rice and wheat in terms of human consumption its rank is third. Its tubers are a good source of carbohydrates, proteins, vitamins, and minerals for human nutrition (Blagoeva *et al.*, 2004). Potato can provide necessary nutrients for the people of the low income group through meeting vegetable demand (Islam *et al.*, 2009; Hossain and Miah, 2012). It can produce more dry matter per unit area and per unit time compared to cereals. The high rate of dry matter production results in large amount of nutrient removal per unit time and most of the soils are unable to meet the demand. Thus it is essential to apply nutrient from external sources such as fertilizers. The production of potato depends on many factors, among them judicious application of nitrogen (N) and potassium (K) play a vital role. For producing 25 to 30 tonne ha^{-1} tuber a mature potato crop removes 120 to 140 kg N ha^{-1} (Patel and Patel, 2001). Tuber dry matter yield and the nitrogen content in potato plants are increased through nitrogen fertilization (Sharifi *et al.*, 2007; Neshev *et al.*, 2014). This crop feeds heavily on soil potassium and the tubers remove 1 to 5 times the amount of nitrogen and 4 to 5 times the amount of phosphate. Because of its high K requirement potato also acts as an indicator crop for K (Fageria *et al.*, 1997). Potassium deficient plants are short with pale-green leaves and later in vegetation the leaves ends and tops become necrotic (Kerin and Berova, 2008; Kumar and Sharma, 2013). Considering the significance of N and K on productivity of potato an

experiment was conducted to study the effect of different levels of nitrogen and potassium on yield of potato grown on new alluvial soils.

MATERIALS AND METHODS

A field experiment was conducted in Gangetic alluvium sandy loam neutral soil (pH 6.9) at the Instructional Farm, Jaguli of Bidhan Chandra Krishi Viswavidyalaya, Kalyani, Nadia (22.93°N, 88.53°E and 9.75 m altitude) during *rabi* season of 2012-13 and 2013-14. The experiment was designed with three replication in randomized block design (RBD), considering ten treatments [T₁: 150 kg ha⁻¹ N, 100 kg ha⁻¹ K₂O, T₂: 150 kg ha⁻¹ N, 150 kg ha⁻¹ K₂O, T₃: 150 kg ha⁻¹ N, 200 kg ha⁻¹ K₂O, T₄: 200 kg ha⁻¹ N, 100 kg ha⁻¹ K₂O, T₅: 200 kg ha⁻¹ N, 150 kg ha⁻¹ K₂O, T₆: 200 kg ha⁻¹ N, 200 kg ha⁻¹ K₂O, T₇: 250 kg ha⁻¹ N, 100 kg ha⁻¹ K₂O, T₈: 250 kg ha⁻¹ N, 150 kg ha⁻¹ K₂O, T₉: 250 kg ha⁻¹ N, 200 kg ha⁻¹ K₂O, T₁₀: 300 kg ha⁻¹ N, 150 kg ha⁻¹ K₂O] in 4.0 x 3.0 m size plots. For all the treatments P₂O₅ dose was 150 kg ha⁻¹. Seed tubers of cultivated variety *Kufri Jyoti* were planted with 50 cm X 20 cm spacing in the third week of November and harvested in fourth week of February. All other standard agronomic practices including plant protection measures recommended for potato tuber production were followed. Observations on plant height, leaf area index (LAI), dry matter accumulation of tuber, tuber number per square meter, tuber yield were recorded and analyzed using the analysis of variance technique. Crop growth rate (g m⁻² day⁻¹) was derived by adopting the procedure recommended by Watson (1958) with certain modification.

$$\text{Crop growth rate (CGR)} = \frac{W_2 - W_1}{t_2 - t_1}$$

Where, W₂ and W₁ were the dry weight of sample plants on two different time t₂ and t₁ respectively.

Economic analysis was performed considering local market rates for inputs and the produce.

RESULTS AND DISCUSSION

Plant height

At 80 DAP (days after planting) among all the treatments 250 kg ha⁻¹ N and 200 kg ha⁻¹ K₂O (T₉) recorded the highest plant height i.e. 71.10 cm where as lowest plant height (44.97 cm) was found in 150 kg ha⁻¹ N and 100 kg ha⁻¹ K₂O (T₁). These observations corroborate the findings obtained by Kumar *et al.*, 2008 and Zelelew *et al.*, 2016. According to Kumar *et al.* (2008) there was increased plant height in potato due increased N dose upto 180 kg ha⁻¹. Zelelew *et al.*, 2016 demonstrated that application of 150 kg ha⁻¹ K₂O recorded highest plant in potato.

Leaf area index

250 kg ha⁻¹ N and 200 kg ha⁻¹ K₂O (T₉) resulted the highest i.e. 1.652 leaf area index (LAI) at 80 DAP which established the fact that nitrogen and potassium had profound influence on the growth of potato. However these observations are in consistent with the

findings obtained by Watson *et al.* (1963) and Veeranna and Khalak (1997). 200 kg ha⁻¹ N and 100 kg ha⁻¹ K₂O (T₄) recorded the lowest (1.173) LAI at 80 DAP.

Crop growth rate

The highest crop growth rate (15.56 g m⁻² day⁻¹) was observed at 60-80 DAP by the application of 250 kg ha⁻¹ N and 200 kg ha⁻¹ K₂O (T₉) which is statistically at par with 300 kg ha⁻¹ N and 150 kg ha⁻¹ K₂O (T₁₀) (15.35 g m⁻² day⁻¹). The lowest crop growth rate i.e. 8.25 g m⁻² day⁻¹ was found in by the application of 150 kg ha⁻¹ N and 100 kg ha⁻¹ K₂O (T₁). This result had the similar trend with the observations of Yadav *et al.* (1999).

Dry matter accumulation of tuber

At 80 DAP T₉ (application of 250 kg ha⁻¹ N and 200 kg ha⁻¹ K₂O) recorded maximum (622.57 g m⁻²) tuber dry matter accumulation. The minimum dry matter accumulation of tuber was observed in T₁ (application of 150 kg ha⁻¹ N, 100 kg ha⁻¹ K₂O) at 80 DAP in potato. Comparable dry matter accumulation of tuber was observed by Zhao *et al.* in 2005.

Tuber number per square meter

The maximum number of tuber per square meter was found in T₉ (application of 250 kg ha⁻¹ N and 200 kg ha⁻¹ K₂O) i.e. 69.70 m⁻² which was statistically at par with T₈ (application of 250 kg ha⁻¹ N and 150 kg ha⁻¹ K₂O) and T₁₀ (application of 250 kg ha⁻¹ N and 150 kg ha⁻¹ K₂O). This result reveals that higher levels of N and K increase the number of tuber per square meter.

Tuber yield

The tuber yield differed significantly due to different doses of nitrogen and potassium on the crop. T₉ (application of 250 kg ha⁻¹ N and 200 kg ha⁻¹ K₂O) produced highest yield (27.03 t ha⁻¹). The next best treatment was T₁₀ (application of 250 kg ha⁻¹ N and 150 kg ha⁻¹ K₂O) which recorded 25.35 t ha⁻¹ tuber yield. The increase in tuber yield due to application of N and K is also documented by Sharma and Arora in 1988.

B:C ratio

Maximum B:C ratio was obtained (1.74) in T₉ (application of 250 kg ha⁻¹ N and 200 kg ha⁻¹ K₂O) followed by T₁₀ (application of 250 kg ha⁻¹ N and 150 kg ha⁻¹ K₂O) i.e. 1.66 and T₈ (application of 250 kg ha⁻¹ N and 150 kg ha⁻¹ K₂O) i.e. 1.62. The minimum B:C ratio was found in T₁ (application of 150 kg ha⁻¹ N and 100 kg ha⁻¹ K₂O) i.e. 0.99 due to poor growth and productivity of crop.

CONCLUSION

From the experiment it can be concluded that application of 250 kg ha⁻¹ N and 200 kg ha⁻¹ K₂O significantly increased the plant height, leaf area index, crop growth rate, dry matter accumulation of tuber and tuber number per square meter which eventually increased the tuber yield and B:C ratio in *Kufri Jyoti* variety of potato. Therefore, it can be recommended that farmers of new alluvial zone of West Bengal may apply 250 kg ha⁻¹ N and 200 kg ha⁻¹ K₂O along with 150 kg K₂O ha⁻¹ to get optimum tuber yield and maximum B:C ratio.

REFERENCE

Blagoeva, V., Iliev, E., Nikolova, M. 2004. Potatoes – cultivation, diseases and pests, and storage. *Enjovche Sofia*. p.105.

- Zeleeuw, D. Z., Lal, S., Kidane, T. T. and Ghebreslassie, B. M. 2016. Effect of Potassium Levels on Growth and Productivity of Potato Varieties. *American Journal of Plant Sciences*.7:1629-1638.
- Fageria, N. K., Baligar, V.C. and Jones, C.A. 1997. Growth and Mineral Nutrition of Field Crops. 2nd Edition. *Marcel Dekker Inc.*, Rome.
- Hossain, M.A., Miah, M. A. M .2012. Post harvest losses and technical efficiency of potato storage systems in Bangladesh. Available at: http://www.nfpcsp.org/agridrupal/sites/default/files/Ayub-final_Report_CF2.pdf (Accessed 15 May 2012)
- Islam, M.Z., Zamam, M.M., Hossain, M.M., Hossain, A. 2009. Integrated nutrient management with liming for potato production in North-West region of Bangladesh. Annual Report 2008-2009, Tuber Crops Research Centre, Bangladesh Agricultural Res Inst, Gazipur, Bangladesh.
- Kerin, V., Berova, M. 2008. Leaf nutrition of plants. *Videnov and son.* p. 124.
- Kumar, P., Sharma, M. 2013. Nutrient Deficiencies of Field Crops: Guide to Diagnosis and Management Kindle Edition. *CABI*. p.400.
- Kumar, A., Tripathi, H.R., Yadav, R.A. and Yadav, D.S. 2008. Diversification of rice (*Oryza Sativa*)- wheat (*Triticum aestivum*) cropping systems for sustainable production in eastern Uttar Pradesh. *Indian Journal of Agronomy*. **53**(1):18-21.
- Neshev, N., Manolov, I., Chalova, V., Yordanova, N. 2014. Effect of nitrogen fertilization on yield and quality parameters of potatoes.. *Journal of mountain agriculture on the Balkans*. **17**(3): 615-627.
- Patel, J.C. and Patel, B.K. 2001. Response of potato to nitrogen under drip and furrow methods of irrigation. *Indian Potato Assoc.* **28**(2-4): 293-295.
- Sharifi, M., Zebarth, B., Hajabbasi, M., Kalbasi, M. 2007. Dry matter and nitrogen accumulation and root morphological characteristics of two clonal selections of 'russet norkotah' potato as affected by nitrogen fertilization. *Journal of plant nutrition*. p. 2243-2253.
- Veeranna, H.K. and khalak, A.1997.Effect of spacing and fertilizer levels on growth parameter and dry matter production of potato crop raised from TPS seedlings. *Crop Res.* **14**(1):119-126.
- Watson, D. J. (1958) .The dependence of assimilation rate on leaf area index, *Annals of Botany*.**22**:22-54.
- Watson, D.J., Thorne, G.N. and French, S.A.W. 1963. Analysis of growth and yield of winter and spring varieties of potato. *Annals of Botany*.**27**:1-22.
- Yadav, K.S., Singh, N., Nehra, B.K., Narula, N. and Lakshminarayana, K. 1999. Azotobactor bio-fertilizer – apotential nitrogen supplement in potato (*Solanum tuberosum* L.) cv. Kufri Sutley. In: *Abstracts Global conference on Potato*, IARI, New Delhi, 6-11 Dec.,1999, pp.123.

Zhao, D., Reddy, K.R., Kakani, V.J. and Reddy, V.R. 2005. Nitrogen deficiency effects on plant growth, leaf photosynthesis and hyper spectral reflectance properties of Sorghum. *European Journal of Agronomy*. **22**:391-403.

UNDER PEER REVIEW

Table 1: Effect of different doses of nitrogen and potassium on plant height, leaf area index, crop growth rate, dry matter accumulation, tuber number per square meter and tuber yield of potato (*Two years pooled data*)

| Treatment | Details | Plant height at 80 DAP (cm) | Leaf area index at DAP | Crop growth rate (g m⁻² day⁻¹) at 60-80 DAP | Dry matter accumulation of tuber at 80 DAP (g m⁻²) | Tuber number per square meter | Tuber yield (t ha⁻¹) | B:C ratio |
|-----------------------|---|------------------------------------|-------------------------------|--|--|--------------------------------------|--|------------------|
| T₁ | 150 kg ha ⁻¹ N, 100 kg ha ⁻¹ K ₂ O | 44.97 | 1.271 | 8.25 | 209.07 | 36.34 | 14.52 | 0.99 |
| T₂ | 150 kg ha ⁻¹ N, 150 kg ha ⁻¹ K ₂ O | 46.84 | 1.433 | 11.53 | 298.10 | 43.90 | 15.70 | 1.05 |
| T₃ | 150 kg ha ⁻¹ N, 200 kg ha ⁻¹ K ₂ O | 50.07 | 1.283 | 13.71 | 351.44 | 57.20 | 18.82 | 1.24 |
| T₄ | 200 kg ha ⁻¹ N, 100 kg ha ⁻¹ K ₂ O | 46.98 | 1.173 | 11.98 | 345.21 | 54.56 | 16.70 | 1.12 |
| T₅ | 200 kg ha ⁻¹ N, 150 kg ha ⁻¹ K ₂ O | 52.08 | 1.383 | 14.82 | 512.12 | 58.91 | 21.99 | 1.45 |
| T₆ | 200 kg ha ⁻¹ N, 200 kg ha ⁻¹ K ₂ O | 59.99 | 1.482 | 14.02 | 580.31 | 62.28 | 23.21 | 1.51 |
| T₇ | 250 kg ha ⁻¹ N, 100 kg ha ⁻¹ K ₂ O | 58.11 | 1.421 | 14.22 | 598.41 | 64.40 | 22.94 | 1.53 |
| T₈ | 250 kg ha ⁻¹ N, 150 kg ha ⁻¹ K ₂ O | 65.05 | 1.382 | 14.97 | 600.40 | 68.03 | 24.96 | 1.62 |
| T₉ | 250 kg ha ⁻¹ N, 200 kg ha ⁻¹ K ₂ O | 71.10 | 1.652 | 15.56 | 622.57 | 69.70 | 27.03 | 1.74 |
| T₁₀ | 300 kg ha ⁻¹ N, 150 kg ha ⁻¹ K ₂ O | 70.09 | 1.531 | 15.35 | 611.45 | 67.87 | 25.35 | 1.66 |
| S. Em(±) | | 1.11 | 0.05 | 0.27 | 3.42 | 1.93 | 0.54 | |
| CD(at 5%) | | 3.29 | 0.14 | 0.8 | 10.15 | 5.73 | 1.61 | |