

# **Analysis of the *Atriplex* Subjected to *Claroideoglossum etunicatum* and to the Desalinator Reject**

## **ABSTRACT**

The objective of this work was to analyze the minerals extracted from the soil and absorbed by *Atriplex nummularia* Lind. submitted to *Claroideoglossum etunicatum* and to the desalinator reject. The experiment was conducted in a greenhouse at the Agronomic Institute of Pernambuco - IPA, Recife, Pernambuco, Brazil. The experimental design was of randomized blocks with the treatments constituted in a factorial scheme, in five levels of salinity: AC: 2.87 mS / cm; T1: 11.54 mS / cm; T2: 12.04 mS / cm; T3: 13,13 and T4: 14,16 mS / cm, associated with the presence and absence of AMF, presence and absence of nutrient solution and autoclaved and non-autoclaved soil. 8.0ml of Hoagland & Arnon complete nutrient solution was added every fortnight. After five months, the contents of the elements absorbed by the plant and present in the soil were evaluated. It was observed that in non-autoclaved soil *Atriplex* absorbed higher nutrient content. Furthermore, the best treatment was the T4 of EC of 14.16 mS/cm + AMF + Hoagland & Arnon solution. Therefore, the high sodium content absorbed (22%) by *Atriplex* evidences the potential of its use in phytoextraction programs in soils affected by salts.

**Keywords:** Mineral nutrition, saline soils, salinity tolerant plants, salt grass.

## **1. INTRODUCTION**

Excess salts and sodium are one of the main factors responsible for soil degradation, causing negative impacts on agricultural production and the sustainability of ecosystems, especially in arid and semi-arid regions [1, 2].

Moreover, as mechanisms for providing nutrients for seedling production there is the use of microorganisms. The mutualistic symbiosis between certain soil fungi and plant roots is

called mycorrhiza. Among the various types of mycorrhiza, the arbuscular mycorrhiza that occur in most forest species stand out, and it is notable for its nutritional benefits, greater resistance to abiotic stress factors and greater tolerance to salinity [3, 4].

The knowledge of the mineral contents in *Atriplex nummularia* can provide subsidies for a management program in saline areas, whose objectives are the extraction of salts and/or the use of this plant as forage [5, 6].

Thus, the objective of this work was to analyze the minerals present in the soil and absorbed by *Atriplex nummularia* Lind. submitted to *Claroideoglossum etunicatum* and to the desalinator reject.

## 2. MATERIAL AND METHODS

The experiment was conducted in a greenhouse at the headquarters of Agronomic Institute of Pernambuco (IPA), Recife, Pernambuco, Brazil.

The soil used was obtained from the IPA Experimental Station, in the city of São Bento do Una, Pernambuco, air dried, dewormed, homogenized and sieved in a 2 mm mesh. Then part of the soil was weighed (8 kg) and used naturally, and the other part, after weighing, was autoclaved at 120 °C for 1 hour. Finally, the soils were transferred to 80 polyethylene vessels. In addition, a soil sample was collected and analyzed in the IPA Soil Fertility Laboratory, according to Tables 1 and 2.

**Table 1. Chemical characteristics of the soil used in the experiment, with precision of 0.1mg.**

P mg/dm <sup>3</sup>	pH (H <sub>2</sub> O)	cmolc/dm <sup>3</sup>					
		Ca	Mg	Na	K	Al	H
350	7,8	16	3,9	3,0	0,7	0,0	0,0

**Table 2. Physical characteristics of the soil used in the experiment, with precision of 0.1g.**

Dap	Dr	Coarse sand	Sand	Silte	Clay	Texture
(g/cm <sup>3</sup> )		4	21	56	19	Franco-silty
1,3	2,6					

The reject used came from the desalinador implanted in the city of Riacho das Almas, Pernambuco with the following characteristics: Electrical conductivity = 11.54 mS / cm at 25 °C, Ca + 2 = 403 mg / L, Mg + 2 = 393.09 mg / L, Na + = 200 mg / L and K + = 40 mg / L, RAS = 23.67, pH = 7.9, Classification for irrigation = C4S4 (Very high salinity water and high sodium concentration; under these conditions it is not suitable for irrigation).

The water (absolute control) used in the experiment contains the following characteristics: Electrical conductivity = 2.87 mS / cm at 25 °C, Ca + 2 = 10.21 mg / L, Mg + 2 = 9.05 mg / L, NA+ = 10.4 mg / L and K+ = 11.6 mg / L, RAS = 0.57, pH = 6.4, Classification for irrigation = C2S1 (Average salinity water, for irrigation whenever there is a moderate degree of irrigation).

*Atriplex nummularia* seedlings with 120 days of age were used, multiplied by means of vegetative propagation by cutting for 30 days. After rooting, the best seedlings were selected to be transplanted into the vessel.

The arbuscular mycorrhizal fungus, *Claroideoglomus etunicatum*, was obtained from the AMF Inoculum Bank of the IPA Soil Microbiology Laboratory, where they are kept under refrigeration at  $\pm 4^{\circ}\text{C}$ . During transplanting, each treatment received 50 g of soil-inoculum with 50 spores.

The nutrient solution used was that of [7], in which 8 ml per vessel was applied every 15 days for 5 months.

The experimental design was a randomized block design consisting of a factorial scheme of 5 irrigation levels (AC: water – EC = 2.87 mS/cm; T1: reject – EC= 11.54 mS/cm; T2: reject plus 7g NaCl – EC= 12.04 mS/cm; T3: reject plus 14g NaCl – EC=13.13 and T4: reject plus 21g NaCl – EC= 14.16 mS/cm), associated to 2 levels of AMF (presence and absence), 2 levels of nutrient solution (presence and absence), and 2 soil levels (autoclaved and non-autoclaved). A 5x2x2x2 factorial with 2 replicates was used, totaling 80 experimental units.

After 5 months, the growth of the *A. nummularia* was monitored by measuring the height, then the aerial part and the root of the *Atriplex nummularia* were collected, separating them at

the height of the plant colon and washed with distilled water. After this, all the material was packed in paper bags and dried in an air circulation oven at 60°C for 72 hours.

Then, the material was ground in a Wiley-type mill equipped with a 42 mm aperture sieve to determine the contents of the absorbed elements (P, K, Ca, Mg) by means of nitroperchloric digestion [8], the content of the elements absorbed (P, K, Ca, Mg, Na) and the total nitrogen were determined by the microkjeldhal method [8].


A soil sample was also collected for complete chemical analysis [9] and phosphorus determinations with Melich-1 solution and reading on the atomic absorption spectrophotometer.

The obtained data were analyzed statistically comparing the means between the treatments through the analysis of variance - Anova and the test of Tukey to 5% of probability using the software Statistica version 10.

### 3. RESULTS AND DISCUSSION

The results obtained in the laboratory determinations regarding the absorbed nutrients N, Ca, Na, Mg and K are presented in Figure 1.

**Figure 1. Elements absorbed by *Atriplex nummularia* depending on the treatments and height.**



**Legend:** Treatments (AC=1; AC+AMF=2; AC+SHA=3; AC+AMF+SHA=4; T1=5; T1+AMF=6; T1+SHA=7; T1+AMF=8; T2=9; T2+AMF=10; T2+SHA=11; T2+AMF=12; T3=13; AC+AMF=14; T3+SHA=15; T3+AMF=16; T4=17; T4+AMF=18; T4+SHA=19; T4+FMA=20; \*AC= 21; \*AC+AMF=22; \*AC+SHA=23; \*AC+AMF+SHA= 24 \*T1=25; \*T1+AMF=26; \*T1+SHA=27; \*T1+AMF=28; \*T2=29; \*T2+AMF=30; \*T2+SHA=31; \*T2+AMF=32; \*T3=33; \*AC+AMF=34; \*T3+SHA=35; \*T3+AMF=36; \*T4=37; \*T4+AMF=38; \*T4+SHA=39; \*T4+AMF=40).

AC= water; T1= reject; T2= reject + 7gNaCl; T3= reject + 14g NaCl; T4= reject + 14gNaCl.

AMF= Arbuscular Mycorrhizal Fungus (*Claroideoglomus etunicatum*).

SHA= Nutrient solution of Hoagland and Arnon (1950).

In Figure 1 it can be observed, for the calcium element, that the best absorption was for the treatment T4 (reject + 21g NaCl) + AMF + nutrient solution with 5.6%. In relation to the other treatments, the treatment with fungus plus nutrient solution stood out in the absorption of this element, being T2 (reject + 7g NaCl) = 4.4% and T3 (reject + 14g NaCl) = 4.7%.

The contents of the elements absorbed by Atriplex with mycorrhizal inoculation, according to [10], in research with peach palm with mycorrhizal fungus followed the decreasing order  $P > K > Mg > Ca$ .

For the nitrogen, it was observed that the best treatment was the T1 (reject) with nutrient solution, corresponding to 7%. Among the other variables, it was observed that the treatment with nutrient solution stood out in the absorption of this element, presenting  $T4 = 5.8\%$  (Figure 1).

Table 3 shows that the T4 treatment (reject plus 21g NaCl) + nutrient solution + AMF was the most significant in relation to the amount of sodium absorbed, with 22%. In addition, among all treatments, the AMF + nutrient solution association was the most expressive for absorbed sodium, with AC (water) = 19.08%, T2 (reject plus 7g NaCl) = 22%, T3 (reject plus 14g NaCl) = 21% and T4 = 22%; only T1 (reject) was more significant for treatment with nutrient solution (19%).

The best treatment for potassium absorption (Table 3) consisted of T3 + AMF treatment with 7.0%. For magnesium (Table 3) it was observed that the best absorption corresponded to the T4 treatment with 17%. Among the other factors, the best absorption for this element consisted of treatments in the absence of fungus and nutrient solution: T2 = 18%, T3 = 19% and T4 = 20%.

In relation to the height of the aerial part of the plant, it was observed that the T4 + AMF + solution of Hoagland and Arnon indicated the highest growth with 110 cm.

**Table 3. Result of the Tukey test at 5% in relation to the elements absorbed by Atriplex in non-autoclaved soil.**

Absorbed nutrients	
Treatments	Potassium (%)
T3 + AMF	7,00 a
T4 + AMF	6,98 ab
T3 + AMF + SH	6,92 ab

T4 + SH	6,66 abc
T3	6,50 abcd
T4	6,32 abcdf
T4 + AMF + SN	6,00 abcdfc
<b>Treatments</b>	<b>Magnesium (%)</b>
T4	20,4 a
T4 + AMF + SH	19,9 ab
T1 + AMF	19,8 ab
AC + AMF	19,5 abc
AC + SHA	19,3 abcd
<b>Treatments</b>	<b>Sodium (%)</b>
T4 + AMF + SHA	22,4 a
T2 + AMF + SHA	22,0 ab
T4 + AMF	21,6 abc
T4 + SH	21,0 abcd
T3 + AMF + SHA	20,9 abcd
<b>Treatments</b>	<b>Height (cm)</b>
T4 + AMF + SHA	110 a
T1 + AMF	106 ab
AC + AMF*	102 abc
AC + SHA	98 abcd

(\*) Autoclaved soil; AC = water; T1 = reject; T2 = reject + 7g NaCl; T3 = reject + 14g NaCl; T4 = reject + 20g NaCl

Different letters indicate significant difference at the 5% probability level.

AC = water; T1 = reject; T2 = reject + 7g NaCl; T3 = reject + 14g NaCl; T4 = reject + 20g NaCl.

AMF = Arbuscular Mycorrhizal Fungus (*Claroideoglomus etunicatum*).

SHA = Solution of Hoagland and Arnon (1950).







[11] indicate that the mutual association between arbuscular mycorrhizal fungi can provide the host plant with greater resistance to the effects of saline stress and to water deficit. Thus, as *Atriplex nummularia* is a salinity-tolerant plant, the association of AMF may contribute to resistance to the deleterious effects of salinity. They showed in their results the presence of sodium in greater quantity in the aerial part than in relation to the root of all the treatments used, which reinforces the hyperaccumulation of sodium in the aerial part of *Atriplex nummularia*.

In Figure 2, the analytical results obtained for non-autoclaved soil, the nutrient content increases as the electrical conductivity of the treatments increases. In this soil, it was observed that the chemical attributes presented higher levels for the phosphorus with 377 mg / dm<sup>3</sup>, followed by sodium with 54.20 cmol.kg<sup>-1</sup>, calcium with 50.97 cmol.kg<sup>-1</sup>, and magnesium 18.84 cmol.kg<sup>-1</sup> and, finally, potassium with 0.75 cmol.kg<sup>-1</sup>. All these results were for the treatment T4 (reject + 21g NaCl) with nutrient solution and AMF, except for the potassium whose highest content was in the treatment T2 (reject + 7gNaCl) with nutrient solution and AMF.

The soil elements were determined before the experiment with (Ca= 16 cmolc/dm<sup>3</sup>; Mg= 3.9 cmolc/dm<sup>3</sup>; Na= 3.0 cmolc/dm<sup>3</sup> and K= 0.7 cmolc/dm<sup>3</sup>), when compared with the final soil collection, presented increasing contents, except for potassium with values reaching 0.5 cmolc / dm<sup>3</sup>.

**Figure 2. Nutrients P, Ca, Mg, Na and K and pH of non-autoclaved and autoclaved soil.**





**Legend:** Treatments (AC=1; TA+AMF=2; AC+SHA=3; AC+AMF+SHA=4; T1=5; T1+AMF=6; T1+SHA=7; T1+AMF=8; T2=9; T2+AMF=10; T2+SHA=11; T2+AMF=12; T3=13; AC+AMF=14; T3+SHA=15; T3+AMF=16; T4=17; T4+AMF=18; T4+SHA=19; T4+AMF=20; \*AC= 21; \*AC+AMF=22; \*AC+SHA=23; \*AC+AMF+SHA= 24 \*T1=25; \*T1+AMF=26; \*T1+SHA=27; \*T1+AMF=28; \*T2=29; \*T2+AMF=30; \*T2+SHA=31; \*T2+AMF=32; \*T3=33; \*AC+AMF=34; \*T3+SHA=35; \*T3+AMF=36; \*T4=37; \*T4+AMF=38; \*T4+SHA=39; \*T4+AMF=40).

AC= water; T1= reject; T2= reject + 7gNaCl; T3= reject + 14g NaCl; T4= reject + 14gNaCl

AMF = Arbuscular Mycorrhizal Fungus (*Claroideoglomus etunicatum*).

SHA = Solution of Hoagland and Arnon (1950).

[12] demonstrated in the experiment on the growth of *Atriplex nummularia* submitted to EC concentrations of 0 to 40 dSm<sup>-1</sup>, that EC: 0 (control) showed the best growth of the plant. There was a slight decrease in soil pH of most treatments, which can be attributed to the ability of *Atriplex nummularia* to absorb sodium. [13] observed reductions in pH values with cultivar of *Atriplex nummularia* with EC: 63.45 µS cm<sup>-1</sup>.

#### 4. CONCLUSION

The high absorption of Na and Mg by *Atriplex nummularia* demonstrates the potential of using this plant in phytoextraction programs in salt affected soils.

*Atriplex nummularia* was not negatively influenced by irrigation with the reject and NaCl treatments. In addition, *Atriplex* behaved as a hyperaccumulating sodium plant, absorbing large amounts of the element with 22%.

The treatments with *Claroideoglomus etunicatum* plus nutrient solution and solely with AMF were the most significant for plant height.

In non-autoclaved soil, *Atriplex nummularia* absorbed more nutrients than the autoclaved soil. The best treatment consisted of T4 (reject + 21g NaCl) + AMF (*Claroideoglomus etunicatum*) + SHA (Hoagland and Arnon nutrient solution).

#### REFERENCES

1. Araújo, MM, Sousa, GM, Rodrigues, LRCC. Techniques used in the recovery of soils affected by salts. *Agroecology in the Semi-Arid*, 2017; 1 (2).
2. Azevedo, LC. Salinity of soil in protected environment. *Revista Exact and Earth Sciences and Agrarian Sciences*, 2018; 13 (1): 52-69.
3. Martins, RMS, Melloni, R, Melloni, EG. Mycelial growth of arbuscular mycorrhizal fungi and formation of mycorrhiza in soil contaminated by cadmium. *Scientia Agraria*. 2017.

4. Silva, EP, Ferreira, PAA, Furtini Neto, AE, Soares, CRF. Arbuscular mycorrhizae and phosphate in the development of Australian cedar seedlings. *Forest Science*, 2017; 27 (4): 1269-1281.
5. Rodrigues, CTA, Silva, TGF, Carvalho, HFS, Moura, MSB. Definitions of production environments for the Atriplex culture in the state of Pernambuco. *Journal of Environmental Analysis and Progress*, 2017; 2 (3): 302-311.
6. Global mechanisms of plant response to salt stress: physiological and molecular levels and implications in biotechnology 6. Tang, X, Mu, X, Shao, H, Wang, H, Brestic, M. *Critical Reviews in Biotechnology*, 2015; 35: 425-437.
7. Hoagland, DR, Arnon, DI. The waterculture method for growing plants without soil - CA: Agricultural Experiment Station, University of California. Berkeley. 1950.
- 8 Embrapa. National Center for Soil Research (Rio de Janeiro, RJ). Brazilian system of soil classification. Brasília: Embrapa-SPI / Embrapa-CNPq, 1999: 412.
9. Embrapa. National Center for Soil Research (Rio de Janeiro, RJ). Manual of methods of soil analysis. 2.ed. Rio de Janeiro, 1997: 212.
10. Brito, VN, Tellechea, FRF, Heitor, LC, Freitas, MSM, Martins, MA. Arbuscular mycorrhizal fungi and phosphate fertilization in the presence of paricá seedlings. *Forest Science*, 2017; 27 (2): 485-497.
11. Berude, MC, Almeida, OS, Riva, MM, Cabanez, PA, Amaral, AA. Mycorrhiza and its agroecological importance. *Encyclopedia Biosphere*, 2015; 11 (22): 133-146.
12. Melo, HF. Growth, biomass production and ion accumulation in *Atriplex nummularia* Lindl. Cultivated under abiotic stress. *Brazilian Journal of Agricultural Engineering*, 2016; 2 (2).
13. Cunha JC, Freire, MBGS, Silva, JDGS. Phytoextraction potential of *Atriplex nummularia* Plants under Nitrogen and Phosphate Fertilization. 2017.

UNDER PEER REVIEW