

Effect of pine needle mulch and irrigation frequency on the yield of *Origanum syriacum* under open field condition.

Abstract

Aims: Due to the intensive harvest from natural habitat and increasing competition as a source of food, *O.syriacum* was introduced into cultivation and soon became one of the promising crops for many farmers in rural areas. The aim of the study is to investigate the effect of pine needle mulch(PN) and irrigation frequency(IF) at 5, 10 and 15 days on the yield under open field conditions.

Study design: The study was organized in randomized complete block design(RCBD) with three replicates.

Place and duration: The study was conducted at the Lebanese agricultural research institute (LARI) in Lebaa station (33°32.681' North, 35°27.088' East, 354 m a.s.l.), south Lebanon, province of Jizzin, under open field condition for two seasons during 2016.

Results: The results indicated a significant effect of mulch in both spring and fall seasons $F(1, 172) = 59.34, p < .0001, \eta^2 = .24$, $F(1, 172) = 77.71, p < .0001, \eta^2 = .30$ respectively. The two-way anova showed no significant effect of irrigation frequency on the average and total plant fresh weight (PFW) across two seasons $F(2, 349) = .80, p = .4522, \eta^2 = .0038$, $F(2, 159) = 1.07, p = .3460, \eta^2 = .007$ respectively. Very little variance of the plant fresh weight (0.2% – spring and 3.1% – fall) was attributed to the interaction of mulch and irrigation frequency. In a comparison between percent weight contribution(PWC) of a different plant part, leaves formed the highest proportion(48.6%) compared to the inflorescence and stem. The fresh weight of weeds(g/m^2) were significantly less in plots mulched with pine needles compared to the control ($F(1,66)=24.43, p<.0001, \eta^2=.75$).

Conclusion: Applying pine needle mulch enhanced most of the yield parameters. Under calcareous clayey soil, *Origanum syriacum* could be irrigated at 10 or 15 days frequencies without remarkable decrease in the yield.

Keywords: *Intensive harvest, fresh weight, Lebaa, percent weight contribution.*

1. INTRODUCTION

O. syriacum belongs to the Labiatae family, genus *Origanum*. Along with other species, *O. syriacum* is native to Palestine, Jordan and Syria [1]. In Lebanon, *O. syriacum* is spread as a wild subshrub in different areas, especially on west-facing slopes of Mount Lebanon mountain chain from north to south between 200 and 600 m above sea level. *O. syriacum* is characterized by its multiple benefits and diverse use. Their leaves and inflorescence contain essential oil rich in phenolic monoterpenoids mainly thymol and carvacrol [2-3-4-5-6-7]. They are used in traditional medicine as a remedy for cold, flu and cough and also as a food and flavor (spice). Extracts from *O. syriacum* show an antifungal activity [5] and they are used as an insecticidal agent against agricultural pests and diseases [8-9-10-11]. The production of *O. syriacum* from the wild habitat in Lebanon varies annually from 700 to 800 t (dry weight) [12]. This production is considered high compared to the small territory of Lebanon and may pose a serious threat to *O. syriacum* as a herbal wild plant. The domestication of *O. syriacum* was initialized several decades ago in response to the unsustainable over exploitation from its wild and also to its high market potential. The climate is characterized by the dry summer which constrains the cultivation of many crops unless the supply of irrigation water. Under natural habitats, one harvest is obtained annually. Without irrigation program, the cultivation of *O. syriacum* will fail to produce a high reasonable and sustainable yield under the prevailing hot and dry weather conditions during the summer period in Lebanon. Another limiting factor of irrigated *O. syriacum* is the weed management. Polyethylene mulch is the main treatments used to control weeds in agricultural crops. The use of chemical herbicides is not advisable on aromatic perennial plants since it is associated with series drawbacks [13]. However some herbicides have shown a selective weed control in *Origanum syriacum* [14]. Straw and pine needle mulch may offer an alternative method to the use of herbicides. The objective of this study is to investigate the effect of irrigation frequency and pine needle mulch on the herbage yield of *Origanum syriacum* under field conditions.

2. MATERIALS AND METHODS

2.1 SITE AND TREATMENTS

The study was conducted at the Lebanese agricultural research institute (LARI) in Lebaa station (33°32.681' North, 35°27.088' East, 354 m a.s.l.), south Lebanon, province of Jizzin, under open field condition. Cuttings of Rmeich ecotype were rooted in propagating units on December 2015 and then planted in the field on March 2016. Based on the soil analysis (Clay texture), the method of planting on raised ridges was adopted to avoid the water logging during rainy period. Plant spacing was 75cm between ridges, 40cm between plants. Drip irrigation system with flow meter was used to deliver the right amount of water to the plants. The irrigation of 1 and 0.6 l/plant/day at 5, 10 and 15 days

frequency were assigned to both organic mulch (pine needle and straw)and control plots(bare soil) respectively.

2.2 EXPERIMENTAL DESIGN AND STATISTICAL ANALYSIS

A total of 6 treatments were organized in complete block design with three replicates (blocks). The experimental unit consisted of 12 plants. Two-way analysis of variance was performed for testing the effect of treatments and their interaction on yield parameters by using SAS for windowsV8. A post hoc Tukey's test ($p < 0.05$) was used to discriminate among means of treatments. Wolcoxon signed-rank test was used for comparison two paired samples of spring and fall plant fresh weight.

2.3 DATA RECORDINGS

Plant weight was recorded on 10 plants cut at 10 cm above soil level .Four plants were assigned for measuring the plant proportion parts (leaves, flowers and stem).Plant height was measured from the soil level to the highest point of the plant. 6 plants from each experimental plot were photographed at 1.5 m height and then the images were processed through the JIamge program for evaluating canopy area. Weed biomass (fresh weight) were recorded at two harvesting dates.

3. DISCUSSION AND RESULTS

3.1 WEATHER DATA

Two cuts were obtained through the implementation of drip irrigation during the hot dry period which lasted from late spring to the mid of autumn. The average daily reference evapotranspiration (ET₀) was 4.1mm for the period from May till the end of September. Max reference evapotranspiration (ET₀) was recorded on June, July and August months (Fig.1).




Figure 1. Reference evapotranspiration (ET₀) and precipitation for the period from January to December of 2016 in Lebaa station

3.2 YIELD PARAMETERS

Analysis of variance showed a significant difference between treatments in regard to plant fresh weight in both spring ($F(5,172)=14.63, p < .0001, \eta^2=0.30$) and fall ($F(5,172)=17.40, p < .0001, \eta^2=34$) harvests respectively. (Table 1.). A Wilcoxon Signed-ranks test indicated that the plant fresh weight

was higher in fall harvest (Mn=142.2, SD=42.6) than in spring (Mn=78.4, SD=27.8), $p < .0001$. The longer growing period, the higher amount of irrigation water received and development of root system [15] caused the plants cut in fall to have a higher biomass as compared to the plants harvested in late spring. The results for two-way ANOVA indicated a significant main effect for mulch in both spring and fall seasons $F(1, 172) = 59.34, p < .0001, \eta^2 = .24$, $F(1, 172) = 77.71, p < .0001, \eta^2 = .30$ respectively. (Table 1).

Table 1. Two-way analysis of variance for the main effect of mulch, irrigation frequency and their interaction on the fresh plant weight and its average percent air dried weight

	Total	Average	Spring harvest	Fall Harvest	Plant Air Dried weight, %
Treatments					
PN5	261.6(50.1) ^a	130.1(42.4) ^a	104.9(28.3) ^a	153.6(39.9) ^a	36.1(5.9)
PN10	261.3(57.2) ^a	133.1(54.5) ^a	92.1(27.5) ^b	169.9(45.8) ^a	35.8(5.2)
PN15	262.3(35.4) ^a	131.7(52.2) ^a	87.6(25.3) ^b	175.9(29.7) ^a	36.4(4.3)
Ctrl5	198.6(46.3) ^b	101.2(39.5) ^b	73.9(23.1) ^c	125.7(35.0) ^b	37.4(8)
Ctrl10	188.7(31.1) ^b	92.0(36.3) ^b	66.0(18.4) ^c	121.9(27.8) ^b	35.9(5.2)
Ctrl15	174.2(34.4) ^b	85.4(52.2) ^b	62.6(18.4) ^c	111.9(28.3) ^b	37.1(4.8)
P	<.0001	<.0001	<.0001	<.0001	.85
Mulch(m)					
Pine needle	261.8(47.4) ^a	131.6(49.7) ^a	94.7(27.7) ^a	166.5(39.7) ^a	36.1(5.1)
Bare soil	186.7(38.6) ^b	92.5(37.0) ^b	67.0(20.2) ^b	119.8(30.8) ^b	36.8(6.1)
P	<.0001	<.0001	<.0001	<.0001	.40
(IF)					
5 days	229.5(57.4)	115.8(43.3)	89.7(30.1) ^a	139.6(40.0)	36.8(7)
10 days	224.3(58.3)	112.4(50.5)	78.2(26.4) ^b	147.2(45.0)	35.8(5.2)
15 days	217.5(56.3)	107.6(49.4)	74.1(25.0) ^b	143.9(43.2)	36.7(4.5)
P	.34	.45	.001	.61	.62
Interaction (M*IF)					
P	.29	.28	.76	.02	.85

Note. Columns with different letters are significantly different (Tukey's test, $p < 0.05$).

Means are reported with standard deviation (SD) in brackets.

Therefore the soil coverage with pine needle increased the average plant fresh weight as compared with bare soil regardless the implementation of irrigation frequency. This may be due to the positive effect of mulch on the total uptake of N P K [16]. Other studies also showed a significant effect of pine needle mulch on plant growth [17]. The results obtained by [18] showed the improving effect of organic mulches mainly pine needle mulch on the vegetative characters of squash. Regarding irrigation frequency, the two-way ANOVA showed no significant effect on the average and total plant fresh weight across two seasons $F(2, 349) = .80, p = .4522, \eta^2 = .0038, F(2, 159) = 1.07, p = .3460, \eta^2 = .007$ respectively (Table 1). The same results were obtained by Khaazaie, H.R and Uçan, K [19-20] which showed no significant effect of irrigation frequency on the herbal biomass. However, the results obtained by Gerami F [21] showed a negative effect of increasing irrigation intervals on morphological traits while it did not affect the oil production and yield of *Origanum vulgare*. In spring season, the observed variation in the fresh weight that was attributed to the irrigation frequency did not exceed 6% ($\eta^2 = .055$). Additionally, the results showed no significant interaction between mulch and irrigation frequency on the total $F(2, 159) = 1.22, p = .2969, \eta^2 = .009$, and average plant fresh weight $F(2, 349) = 1.25, p = .2891, \eta^2 = .006$ indicating that the effect of mulch does not depend on irrigation frequency. Very little variation in the plant fresh weight, (0.2%, $\eta^2 = .002$ – Spring and 3.1%-, $\eta^2 = .031$ fall) (Table 1) was attributed to the interaction of mulch and irrigation frequency. Air dried plant weight comprised in average about 36% of the total yield biomass and did not differ significantly between treatments. In a comparison between percent weight contribution of a different plant part, leaves formed the highest proportion compared to the inflorescence and stem (Table 2). In fall harvest, neither mulch nor irrigation frequency and their interaction significantly affected the percent weight contribution of the leaf, inflorescence and stem. In spring harvest, a two-way ANOVA indicated a significant effect of mulch on the percent contribution of leaves. The effect of mulch on percent contribution of inflorescence showed a marginal significance ($p = 0.0971$). Plants irrigated at 15 days frequency had a slightly more inflorescence as compared with plants irrigated at 5 days frequency (marginal significance $p = .0887$).

Table 2. Two-way analysis of variance for the main effect of mulch, irrigation frequency and their interaction on the percent weight contribution of plant parts of *Origanum syriacum* (Leaves, Inflorescence, and Stems) measured at spring and fall seasons.

	Spring Harvest			Autumn Harvest			Average		
	Leaves	Infl.	Stem	Leaves	Infl.	Stem	Leaves	Infl.	Stem
Treatmnts									
PN5	39.3(7.7)	30.3(7.0)	29.9(3.5)	57.5(3.6)	17.9(3.7)	24.7(2.0)	46.6(13.8)	26.1(12.1)	27.3(3.8)
PN10	37.7(7.1)	33.5(8.9)	28.8(3.3)	59.5(5.4)	15.6(8.1)	24.9(3.8)	48.6(12.9)	24.5(12.4)	26.8(3.9)
PN15	33.3(3.0)	38.6(1.5)	27.1(2.9)	63.0(6.7)	13.4(5.1)	23.5(3.6)	46.3(19.1)	28.4(17.8)	25.3(3.2)
Ctrl5	43.4(9.7)	28.1(7.6)	28.6(2.4)	55.5(5.2)	21.6(6.0)	23.0(1.0)	49.4(9.7)	24.8(7.3)	25.8(3.4)
Ctrl10	44.0(6.7)	27.9(7.5)	28.2(2.5)	59.6(6.6)	17.2(5.7)	23.1(2.6)	51.8(10.3)	22.5(8.4)	25.7(3.6)
Ctrl15	39.7(6.6)	33.5(8.7)	26.8(3.6)	60.7(8.8)	16.7(8.2)	22.6(2.8)	50.2(13.2)	25.1(11.9)	24.7(3.8)
P	.13	.14	.56	.47	.45	.64	.92	.92	.61
Mulch									
PN	36.8(6.5) ^b	34.1(7.2)	28.6(3.2)	60.0(5.6)	15.6(5.9)	24.4(2.8)	47.2(15.1)	26.3(14.0)	26.5(3.7)
CTRL	42.8(7.4) ^a	29.9(7.9)	27.8(2.8)	58.7(7.0)	18.3(6.7)	22.9(2.2)	50.5(11.0)	24.1(9.3)	25.4(3.5)
P	.02	.09	.48	.56	.22	.13	.32	.46	.21
Irrigation frequency									
5 days	41.3(8.5)	29.2(7.0)	29.3(3.0)	56.6(4.3)	19.6(5.0)	23.9(1.8)	47.9(11.9)	25.5(10.0)	26.6(3.7)
10 days	40.9(7.4)	30.7(8.4)	28.5(2.8)	59.5(5.7)	16.4(6.8)	24.0(3.2)	50.2(11.5)	23.5(10.4)	26.3(3.7)
15 days	36.8(6.0)	35.8(6.7)	26.9(3.1)	61.9(7.6)	15.1(6.7)	23.1(2.7)	48.3(16.2)	26.7(14.9)	25.0(3.4)
P	.22	.08	.20	.15	.24	.66	.83	.67	.31
Interaction(Mulch*Frequency)									
P	.70	.69	.95	.84	.83	.93	.98	.96	.96

Note. Columns with different letters are significantly different (Tukey's test, $p < 0.05$).

Means are reported with standard deviation (SD) in brackets

The effect of treatments was examined on the canopy area, plant height and number of shoots per plant. The results showed a significant difference between treatments on the three parameters. These parameters were positively affected by pine needle mulch. The irrigation frequency had a significant effect on the number of shoots/plant ($p=0.0222$) (Table3) whereas no significant effect was observed on the plant length and plant canopy area respectively ($p=0.5997$, $p=0.3351$). The highest number of shoots per plant was recorded in plots irrigated at 5 days frequency (Table3).

Table 3. Two-way analysis of variance for the main effect of mulch, irrigation frequency and their interaction on the canopy area, number of shoots and plant height of *Origanum syriacum*

	Canopy area ,cm ²	Number of shoots	Plant length ,cm
Treatments			
PN5	1727.4(360.3) ^a	39.0(12.7) ^a	55.7(8.1) ^a
PN10	1721.4(350.9) ^a	29.5(10.7) ^{bc}	55.5(7.2) ^a
PN15	1689.2(166.2) ^a	32.9(9.2) ^{ab}	54.4(5.4) ^a
Ctrl5	1248.8(340.4) ^b	26.6(9.3) ^c	49.5(7.4) ^b
Ctrl10	1318.5(268.6) ^b	26.0(13.1) ^c	49.7(5.5) ^b
Ctrl15	1128.1(144.6) ^b	23.6(9.4) ^c	50.3(7.0) ^b
P	.04	<.0001	.006
Irrigation frequency(if)			
5 days	1474(418.7)	32.8(12.7) ^a	52.6(8.3)
10 days	1531.8(368.9)	27.7(12.0) ^b	52.6(7.0)
15 days	1408.6(328.5)	28.3(10.3) ^b	(52.3(6.5)
P	.33	.02	.59
Mulch(m)			
Pine needle	1715.1(304.3)	33.8(11.5) ^a	55.2(6.9) ^a
Control(Bare soil)	1241.6(274.8)	25.4(10.7) ^b	49.8(6.6) ^b
P	.004	<.0001	.001
Interaction (m*if)			
P	.52	.003	.10

Note. Columns with different letters are significantly different (Tukey's test, $p < 0.05$).

Means are reported with standard deviation (SD) in brackets

3.3 WEEDS

In regards to the weed biomass (fresh weight/m²), the overall analysis of variance indicated a significant difference between treatments ($F(5, 66) = 6.52, P < .0001, \eta^2 = 0.33$) (Figure 2). Control plots accumulated significantly more weed biomass (Mn=328.7/m², SD=246.7) than plots mulched with pine needles (Mn=123.6/m², SD=69.7), ($F(1, 66) = 24.43, p < .0001, \eta^2 = .75$). Our results regarding the suppression effect of pine needle mulch on weed growth agree with the results obtained by Burkhard Nel [22]. Irrigation frequency showed a marginal significant effect on weed biomass. Plots irrigated at high frequency (5 days frequency) yielded a higher weed biomass (Mn=280.9, SD=257.4) as compared with plots irrigated at low frequency (15 days frequency) (Mn=164.8, SD=121.1), ($F(2, 66) = 2.64, p = .0792, \eta^2 = .16$).




Figure 2. Means of weed fresh biomass recorded from bare and mulched plots cropped with *O. syriacum* and irrigated at three frequencies (5, 10 and 15 days)

4. CONCLUSION

There was no significant effect between the three adopted frequencies on the fresh plant weight. Applying pine needle mulch enhanced most of the yield parameters except for the percent weight contribution of plant parts. No obvious trend of the interaction between pine needle mulch and irrigation frequency. Under calcareous clayey soil, *Origanum syriacum* could be irrigated at 15 days frequency without remarkable decrease in the yield.

Reference

1. Kokkini S. Taxonomy, diversity and distribution of origanum species.. In the IPGRI International Workshop on Oregano. 1996;CIHEAM, Valenzano, Bari, Italy.
2. Ali-Shtayeh M.S et al. Secondary treated effluent irrigation did not impact chemical composition, and enzyme inhibition activities of essential oils from *Origanum syriacum* var. *Syriacum*.. *Industrial Crops and Product*. 2018;111:775-786, DOI: 10.1016/j.indcrop.2017.11.055.
3. Bakkour Y, et al. Analysis of the essential oils of *Salvia Libanotica* and *Origanum Syriacum*. *Journal of Natural Products*. 2011;Volume 4:51-56, ISSN 0974 – 5211.
4. D'Antuono et al. Variability of Essential Oil Content and Composition of *Origanum vulgare* L. Populations from a North Mediterranean Area (Liguria Region, Northern Italy). *Annals of Botany*.2000;Volume 86, Issue 3, 1, Pages 471–478
5. Daouk K.R, et al.. Antifungal Activity of the Essential Oil of *Origanum syriacum* L. *Journal of food protection*.1995;58(10):1147-1149.
6. Farhat M, et al. Analysis of the essential oil compounds of *Origanum syriacum* L. In *Acta Facultatis Pharmaceuticae Universitatis Comeniana*, December 2012;DOI: 10.2478/v10219-012-0020-x
7. Hunsu Can Baser H. The essential oil of *Origanum syriacum* L. var. *sinaicum* (Boiss.) letsvaart. *Flavour and Fragrance Journal*.2003;18(2):98-99. DOI10.1002/ffj.1169
8. Abderrahmane K, Lahcen E. Insecticide Effect of Plant Extracts on Aphids of Watermelon. *Journal of Biology, Agriculture and Healthcare*. 2015; ISSN 2224-3208 (Paper) ISSN 2225-093X (Online) Vol.5, No.3.
9. Khalfi O, et al. Chemical composition and insecticidal properties of *Origanum glandulosum* (Desf.) essential oil from Algeria. *Journal of the science of food and agriculture*.2008; <https://doi.org/10.1002/jsfa.3251>
10. Abou Jawdah Y, et al. Antifungal activities of extracts from selected Lebanese wild plants against plant pathogenic fungi, *Phytopathol. Mediterr*.2004;43, 377–386.
11. Al Banna L, et al..Effect of plant extracts and essential oils on root-knot nematode. *Phytopathol. Mediterr*. 2003;42, 123–128.
12. GEF-UNDP-LARI. 2013.Conservation guideline for medicinal and aromatic plants in Lebanon.
13. Vouzounis NA,et al.Chemical control of weeds in the aromatic crops lavender, oregano and sage. *Technical bulletin*. 2003;218, ISSN 0070=2315. Available at <http://news.ari.gov.cy/publications/tb218-vouzounis.pdf>

14. Qasem RJ and Foy C. Selective Weed Control in Syrian Marjoram (*Origanum syriacum*) with Oxadiazon and Oxyfluorfen Herbicides. Weed technology. 2006;20(3): 670-676.
<https://doi.org/10.1614/WT-04-129R1.1>
15. Jaafar H, et al. Water productivity of *Origanum syriacum* under different irrigation and nitrogen treatments using an automated irrigation system. WIT Transactions on Ecology and The Environment. 2015; Vol 196, © 2015 WIT Press, www.witpress.com, ISSN 1743-3541 (on-line), doi:10.2495/WRM150181
16. Acharya CL and Charma PD.. Tillage and mulch effects on soil physical environment, root growth, nutrient uptake and yield of maize and wheat on an Alfisol in north-west India. Soil and tillage research. 1994; Volume 32 issue 4, Pages: 291-302. [https://doi.org/10.1016/0167-1987\(94\)00425-E](https://doi.org/10.1016/0167-1987(94)00425-E).
17. Bawang F.. Utilization of Benguet pine needles as mulch: their effects on germination, growth, yield and some postharvest qualities of pechay, snapbean and tomato. University Library of the Philippines at Los Banos. 1988; Available at <http://www.uplb.edu.ph>
18. Bhatt L, et al.. Effect of mulch materials on vegetative characters, yield and economics of summer squash (*Cucurbita pepo*) under rainfed mid-hill condition of Uttarakhand. Vegetable Science 2011;38(2): 165-168.
19. Khaazaie HR, et al.. Effect of irrigation frequency and planting density on herbage biomass and oil production of thyme (*Thymus vulgaris*) and hyssop (*Hyssopus officinalis*). Industrial crops and products. 2008; volume 27, issue 3, pages 315-321.
<https://doi.org/10.1016/j.indcrop.2007.11.007>
20. Uçan K, et al. Effect of irrigation frequency and amount on water use efficiency and yield of sesame (*Sesamum indicum* L.) under field conditions. Field crops research. 2007; Volume 101, Issue 3, Pages 249-258. <https://doi.org/10.1016/j.fcr.2006.11.011>.
21. Gerami F, Parviz RM, Ghorbani R, and Hassani A. Effects of irrigation intervals and organic manure on morphological traits, essential oil content and yield of oregano (*Origanum vulgare* L.). Annals of the Brazilian Academy of Sciences. 2016; 88(4): 2375-2385.
22. Burkhard N, Lynch D, Percival D.. Effects of Pine-Needle and Compost Mulches and Weeds on Nitrogen Dynamics in an Organically-Managed Highbush Blueberry Field., Proc. Organic Fruit Conference Eds, Acta Hort. 2010; 873, ISHS 2010.