

Review Paper

HEAVY METALS CONTAMINATION AND POTENTIAL HUMAN HEALTH RISK VIA CONSUMPTION OF VEGETABLES FROM SELECTED COMMUNITIES IN ONELGA, RIVERS STATE, NIGERIA

ABSTRACT

Background: Anthropogenic activities such as excessive oil exploitation, automobile emissions, agricultural activities tend to elevate the concentrations of heavy metals in the soil, these metals are absorbed by plant roots, thereby resulting to heavy metal contamination of such plants.

Objectives: This study evaluated the concentration of heavy metals namely: Lead (Pb), Cadmium (Cd), Nickel (Ni), Arsenic (As), Iron (Fe), and Copper (Cu) via consumption of vegetables from selected communities in Ogba/Egbema/Ndoni Local Government Area (ONELGA), Rivers State, Nigeria. **Materials and methods:** Sixteen different vegetable samples comprising of *Vernonia amygdalina*, *Talinum triangulare*, *Abelmoschus esculentus* and *Telfaria occidentalis* were digested and analyzed for heavy metals using Solar Thermo Elemental Flame Atomic Absorption Spectrophotometer (STEF-AAS). Results obtained from this study were used to estimate the human health risk of these heavy metals. **Results:** The results showed that the mean concentrations and (range) of total Pb, Cd, Ni, As, Fe, and Cu were 1.568 (0.164-4.925), 1.040 (0.359-3.000), 65.764 (2.764-10.301), 15.533 (0.152-2.640), 82.900 (61.576-101.525), and 29.266 (12.029-46.540) mg/kg, respectively. The concentration of Pb, Cd, Cu and Ni exceeded the permissible limit recommended by WHO/FAO, EC/CODEX, JECFA and NAFDAC respectively. The concentration of As exceeded the permissible limit by WHO/FAO. The concentration of Fe was below permissible limit by WHO/FAO and EC/CODEX. The Estimated Daily Intake (EDI) concentrations of Pb and Cd

exceeded the permissible tolerable daily intake of metals (TDI) but fall within the upper tolerable daily intake (UTDI) recommended by Institute of medicine, FDA. Ni, Fe, and Cu fall within the recommended TDI and the UTDI. As concentration exceeded the recommended TDI and the UTDI. The Target Hazard Quotient (THQ) of Pb, Cd, Cu and Ni were greater than 1. The Hazard Index (HI) values were greater than 1. The Life Cancer Risk (LCR) of Pb were within the range of permissible predicted lifetime risks for carcinogens as recommended by US EPA. **Conclusion:** The result from the present study may indicate that the exposed population may be at risk of Pb, Cd, Ni and Cu toxicity.

KEYWORDS: Heavy metals, anthropogenic food safety, , THQ, Risk assessment.

INTRODUCTION

Heavy metals are hazardous contaminants in food and the environment and they are non-biodegradable having long biological half-lives.¹The implications associated with heavy metals contamination are of great concern, particularly in agricultural production systems.^{2,3} due to their increasing trends in human foods and environment. Food safety has been a general concern in recent years.^{4,5}Vegetables are an important part of the human diet because they contain carbohydrates, proteins, vitamins, minerals as well as trace elements. These metals can pose as a significant health risk to humans, particularly in elevated concentrations above the very low body requirements.⁶ So, the metals must be controlled in food sources in order to assure public health safety.⁷Excessive amount of heavy metals in food cause a number of diseases, especially cardiovascular, renal, neurological, and bone diseases.⁸These metals could reach food chain through various biochemical process and ultimately biomagnified in various trophic levels and eventually threaten the health of human. Heavy metals enter into plants mainly via roots from the soil and travel along the food chain.⁹Once entered in the bodies of living organisms, heavy metals can pose serious threats due to their non-biodegradable nature, long biological half-lives and their potential to accumulate in different body parts, e.g., fatty tissues.¹⁰Moreover, most of the heavy metals are extremely toxic because of their solubility in water. Even low concentrations of heavy metals have damaging effects on human and animal populations^{11,12} by causing cardiovascular, kidney, nervous, mental impairment and bone diseases.¹³

The contamination of soil and vegetables by heavy metals is also a global environmental issue. They are ubiquitous in the environment through various pathways, due to natural and anthropogenic activities such as addition of manures, sewage sludge, fertilizers and pesticides to soils, oil exploitation, mining, railings, industrial wastes, agricultural runoff, occupational exposure, paints and treated timber and aerosols.¹⁴ Excessive accumulation of heavy metals in agricultural soils through the use of agrochemicals and by other sources may not only result in soil contamination but also lead to elevated heavy metal uptake by vegetables and thus affect food quality and safety.¹⁵

However, this study was designed to assess the level of heavy metals Pb, Cd, Ni, As, Fe, and Cu in four most grown and consumed vegetable species: Bitter leaf (*Vernonia amygdalina*), Water leaf (*Talinum triangulare*), Okra leaf (*Abelmoschus esculentus*), and Fluted pumpkin (*Telfaria occidentalis*) in selected communities in ONELGA and also to assess the potential health risk associated with their daily intake.

MATERIALS AND METHODS

Study area

The present study was carried out in Ogba/Egbema/Ndoni Local Government Area (ONELGA) in Rivers State located in the South-South region Nigeria as shown in Figure 1. Ogba/Egbema/Ndoni Local Government Area is located approximately between Longitudes 70 31 to 120 45 East of Greenwich meridian and Latitudes 60 71 to 100 40 north of equator. The population of ONELGA is about 200,000 people.¹⁶ with the capital located at Omoku. It is a major hub for oil and gas exploration by major international oil companies (IOCs) in the Niger Delta region.




Figure 1; Map showing research area and sampling points in ONELGA, Rivers State, Nigeria

Sample collection

Fresh samples each of the four selected vegetables namely, *V. amygdalina*, *T. triangulare*, *A. esculentus* and *T. occidentalis* were collected randomly from different farms from the selected communities in ONELGA, Rivers State. The samples were properly tagged according to communities in polythene bags and taken to the Herbarium of Plant Science and Biotechnology Department, University of Port Harcourt for identification, and subsequently taken to the laboratory for heavy metal analyses.

Sample preparation

Leafy vegetable samples were washed with tap water thoroughly to remove dust particles, soil, unicellular algae etc. The edible parts of the samples were further washed with distilled water and finally with deionized water. The washed vegetables were dried with blotting paper followed by filter paper at room temperature to remove surface water. The vegetables were immediately kept in desiccators to avoid further evaporation of moisture from the materials. After that the vegetables were chopped into small pieces they were oven dried at $(55 \pm 1) ^\circ\text{C}$. Then the vegetables were crushed into fine powder using a porcelain mortar and pestle. The

resulting powder was kept in air tight polythene packet at room temperature before being taken to the laboratory for digestion and metals analyses.

Digestion

A total volume of 100 ml of H_2SO_4 , HNO_3 , $HClO_4$ in the ratio of 40%:40%:20% was mixed together. A portion of (5g) of the samples was weighed and digested with a 2 ml of the mixed acid to each of the samples in a conical flask. The samples were then digested in a fume cupboard with hot plate until white fumes appeared. After that, the solution was then cooled, filtered and transferred into a 100 ml volumetric flask and made up to the 100 ml with distilled water.

Heavy metal health risk assessment

The potential health risks of heavy metal consumption through vegetables were assessed based on the Estimated daily intake of metals (EDI).¹⁷, the target hazard quotient (THQ).^{18,19} and Hazard index (HI).²⁰, The Estimated daily intake of metals (EDI) was calculated to averagely estimate the daily metal loading into the body system of a specified body weight of a consumer. This will inform the relative phyto-availability of metal.

Estimated daily intake (EDI)

$$EDI = \frac{C_{metal} \times D_{food\ intake}}{B_{Waverage}}$$

Where:

C_{metal} is the metal concentration in vegetable in mg/kg, D_{food} intake is the daily intake of food in kg/person and $B_{Waverage}$ is average body weight in kg/person.

An average daily consumption of 0.2 kg of vegetable was assumed in this study. This value was adopted because vegetable constitute a major part of the diet. Average adult body weight was considered to be 60 kg.

Non-carcinogenic health effect

Target hazard quotient: Non-carcinogenic risk estimation of heavy metals consumption was determined using THQ values.^{18,19} THQ is a ratio of the determined dose of a toxicant to a reference dose considered harmful.²¹ If the ratio is equal to or greater than 1, an exposed population is at risk. THQ values were calculated using the following formula below:

$$\text{THQ} = \frac{\text{Efr} \times \text{ED} \times \text{FIR} \times \text{C}}{\text{RfDo} \times \text{BWaverage} \times \text{ATn}} \times 10^{-3}$$

Where

Efr is exposure frequency in 156 days year⁻¹,

ED is exposure duration in 56 years equivalent to an average lifetime,

FIR is average daily consumption in Kg person⁻¹ day⁻¹,

C is concentration of metal in food sample in mg/kg

RfDo is reference dose in mg/Kg day⁻¹ and

ATn is average exposure time for non–carcinogens in days (156 x 56).

The following reference doses were used (Pb=0.0035 mg/kg Cd = 0.001 mg/kg, Ni=0.02mg/kg, As = 0.050mg/kg, Fe = 0.700mg/kg, Cu = 0.040mg//kg)

Calculation of hazard index:

Hazard index (HI) is used to evaluate the potential risk to human health when more than one heavy metal is involved. Hazard index was calculated as the sum of hazard quotients (HQs).

Since different pollutants can cause similar adverse health effects, it is often appropriate to combine HQs associated with different substances.

$$HI = \sum THQ (THQ_1 + THQ_2 + THQ_3 + \dots + THQ_n)$$

Carcinogenic Risk (CR)

Incremental lifetime cancer risk (ILCR) is the lifetime probability of an individual developing any type of cancer due to carcinogenic daily exposure to a contaminant over a life time. The ILCR is obtained using the Cancer Slope Factor (CSF) which evaluates the probability of an individual developing cancer from oral exposure to contaminant levels over a period of a lifetime as described by ATSDR in 2010 and it is contaminant specific. Ingestion cancer slope factors are expressed in units of (mg/kg/day).

The cancer risk was calculated using the equation below;

$$\text{Carcinogenic risk} = \text{EDI} \times \text{CSF}_{\text{ing}}$$

Where:

EDI is the estimated daily intake of each heavy metal (mg/kg/day)

CSF_{ing} is ingestion cancer slope factor (mg/kg/day)⁻¹ (Pb=0.0085, Cd=0.38, Ni=0.91, and As=1.5)

US EPA, 2011 states that 10⁻⁶ (1 in 1,000,000) to 10⁻⁴ (1 in 10,000) represent a range of permissible predicted lifetime risks for carcinogens. Chemical for which the risk factor falls below 10⁻⁶ may be eliminated from further consideration as a chemical of concern.

RESULTS

The results of different vegetable samples analyzed for Pb, Cd, Ni, As, Fe and Cu are shown in Table 1. The results from the present study showed that the mean concentrations and range of

total Pb, Cd, Ni, As, Fe, and Cu in edible portions of vegetables were 1.568 (0.164-4.925); 1.040 (0.359-3.000); 65.764 (2.764-10.301); 15.533 (0.152-2.640); 82.900 (61.576-101.525) and 29.266 (12.029-46.540) mg/kg, respectively. The average concentrations of heavy metals in all the analyzed vegetable samples were in the order of Fe > Ni > Cu > As > Pb > Cd. The coefficients of variation for the six heavy metals in the analyzed vegetables also followed the same order. The highest level of Pb was recorded in *T. triangulare* (4.925 mg/kg) from Elleta community, while the lowest level of Pb was recorded in *T. triangulare* (0.164 mg/kg) from Okposi community. The highest level of Cd was observed in *V. amygdalina* (3.000 mg/kg) from Obrikom community. However, *A. esculentus* from Obrikom community was below detectable limit (BDL < 0.001). The highest concentration of Ni was observed in *T. occidentalis* (10.301 mg/kg) from Elleta community, *T. triangulare* from Omoku community and *A. esculentus* from Elleta community were below detectable limit (BDL < 0.001). The highest concentration of As was recorded in *T. occidentalis* (2.640 mg/kg) from Elleta community. However, *A. esculentus* from Okposi community was below detectable limit (BDL < 0.001). The highest concentration of Fe was observed in *A. esculentus* (101.525 mg/kg) from Omoku community while the lowest level was recorded in *V. amygdalina* (61.576mg/kg) from Okposi community. The highest concentration of Cu was recorded in *T. triangulare* (46.540 mg/kg) from Omoku community while the lowest level was recorded in *A. esculentus* (12.029 mg/kg) from Okposi community.

TABLE 1:Concentration of heavy metals in mg/kg in selected vegetable samples in ONELGA

| Location | Sample | Pb | Cd | Ni | As | Fe | Cu |
|----------|------------------------|---------------|---------------|---------------|---------------|----------------|----------------|
| Obrikom | <i>V. amygdalina</i> | 1.634 ± 0.012 | 2.977 ± 0.02 | 3.848 ± 0.003 | 1.364 ± 0.02 | 86.902 ± 0.016 | 43.604 ± 0.005 |
| | <i>T. triangulare</i> | 3.824 ± 0.021 | 0.394 ± 0.05 | 2.963 ± 0.009 | 0.385 ± 0.005 | 78.766 ± 0.04 | 21.109 ± 0.01 |
| | <i>A. esculentus</i> | 0.773 ± 0.033 | BDL | 2.889 ± 0.01 | 1.782 ± 0.002 | 81.866 ± 0.05 | 19.526 ± 0.003 |
| | <i>T. occidentalis</i> | 1.375 ± 0.011 | 0.855 ± 0.013 | 4.158 ± 0.05 | 0.566 ± 0.047 | 90.494 ± 0.005 | 38.443 ± 0.006 |
| Omoku | <i>V. amygdalina</i> | 2.294 ± 0.005 | 1.816 ± 0.012 | 6.565 ± 0.04 | 0.656 ± 0.006 | 92.924 ± 0.003 | 36.109 ± 0.003 |
| | <i>T. triangulare</i> | 2.373 ± 0.001 | 0.825 ± 0.006 | BDL | 2.384 ± 0.006 | 84.374 ± 0.007 | 46.536 ± 0.006 |
| | <i>A. esculentus</i> | 0.567 ± 0.009 | 0.494 ± 0.02 | 3.456 ± 0.02 | 1.634 ± 0.01 | 96.617 ± 0.02 | 14.838 ± 0.005 |
| | <i>T. occidentalis</i> | 0.381 ± 0.007 | 1.454 ± 0.006 | 4.206 ± 0.01 | 0.374 ± 0.02 | 80.727 ± 0.006 | 23.244 ± 0.005 |

| | | | | | | | |
|---|------------------------|---------------|---------------|---------------|---------------|-----------------|----------------|
| Okposi | <i>V. amygdalina</i> | 0.413 ± 0.006 | 1.533 ± 0.003 | 8.177 ± 0.03 | 0.567 ± 0.01 | 61.589 ± 0.010 | 28.258 ± 0.005 |
| | <i>T. triangulare</i> | 0.168 ± 0.003 | BDL | 5.542 ± 0.06 | 1.478 ± 0.01 | 63.591 ± 0.007 | 42.394 ± 0.005 |
| | <i>A. esculentus</i> | 2.894 ± 0.005 | 0.724 ± 0.02 | 4.933 ± 0.007 | BDL | 72.127 ± 0.005 | 12.033 ± 0.004 |
| | <i>T. occidentalis</i> | 0.255 ± 0.005 | 1.345 ± 0.03 | 2.780 ± 0.02 | 0.782 ± 0.01 | 75.642 ± 0.008 | 26.247 ± 0.005 |
| Elleta | <i>V. amygdalina</i> | 1.860 ± 0.009 | 0.761 ± 0.009 | 8.166 ± 0.02 | 0.297 ± 0.006 | 101.520 ± 0.004 | 24.394 ± 0.005 |
| | <i>T. triangulare</i> | 4.908 ± 0.012 | 1.545 ± 0.003 | 7.441 ± 0.009 | 0.474 ± 0.01 | 88.955 ± 0.01 | 15.407 ± 0.003 |
| | <i>A. esculentus</i> | 0.234 ± 0.016 | 0.364 ± 0.006 | BDL | 0.157 ± 0.004 | 95.515 ± 0.01 | 33.492 ± 0.003 |
| | <i>T. occidentalis</i> | 1.128 ± 0.005 | 1.536 ± 0.003 | 10.241 ± 0.11 | 2.633 ± 0.006 | 74.538 ± 0.009 | 42.611 ± 0.007 |
| TOTAL MEAN | | 1.568 | 1.040 | 65.764 | 15.533 | 82.900 | 29.266 |
| Permissible limit of heavy metals for vegetables by health bodies (mg/kg) | | | | | | | |
| WHO/FAO, 2001 | | 2.0 | 1.0 | - | 0.43 | 425.00 | 73.00 |
| EC/CODEX, 2001 | | 0.3 | 0.2 | - | - | - | - |
| NAFDAC | | - | - | 2.7 | - | - | - |
| JECFA, 2005 | | - | - | - | - | - | 40 |
| CHINA, 1994 | | 0.2 | - | - | - | - | - |
| REFERENCE DOSES | | 0.0035 | 0.001 | 0.02 | 0.050 | 0.700 | 0.040 |

Estimated daily intake (EDI) of metals in (mg/kg) for adult population

The results of the EDI for adults are shown in Table 2. The result of Pb was within the range of 0.001 to 0.016mg/kg. The highest concentration was observed in *T. triangulare* (0.016mg/kg) from Elleta community while the lowest levels were recorded in *T. occidentalis* (0.001 mg/kg) from Omoku community, *V. amygdalina* (0.001 mg/kg), *T. triangulare* (0.001 mg/kg), *T. occidentalis* (0.001 mg/kg) all from Okposi community and *A. esculentus* (0.001 mg/kg) from Elleta community. The concentration value of Cd ranged between 0.001 to 0.010 mg/kg with the highest concentration recorded in *V. amygdalina* (0.010 mg/kg) from Obrikom community. However, concentration of *A.esculentus* from Obrikom community and *T.triangulare* from Okposicommunity were below detectable limits (BDL< 0.001). The result of Ni was within the range of 0.009 to 0.034 mg/kg with *T.occidentalis* fromElletacommunity recording the highest value (0.034 mg/kg) while *T.triangulare* from Omoku community and*A.esculentus* fromElletacommunity were below detectable limits (BDL < 0.001). The value of As ranged between 0.001 to 0.008 mg/kg. The highest concentration was observed in *T. triangulare* (0.008 mg/kg) from Omokucommunity and also, the concentration of *A.esculentus* from Okposi community was below detectable limit (BDL < 0.001). The value of Fe ranged between 0.205 to 0.338 mg/kg with the highest concentration recorded in *V. amygdalina* (0.338 mg/kg) from Elleta community while the lowest level was observed in *V.amygdalina*(0.205 mg/kg) from Okposi community.

The value of Cu ranged between 0.040 to 0.155 mg/kg with the highest concentration recorded in *T.triangularare* (0.155mg/kg) from Omoku Community while the lowest level was observed in *A. esculentus* (0.040 mg/kg) from Okposi community.

Table 2: Estimated daily intake (EDI) of heavy metals in mg/kg for adult (60kg) population in ONELGA via consumption of vegetables

| LOCATION | SAMPLES | Pb | Cd | Ni | As | Fe | Cu |
|----------|--|-------|--------|-------|------------------------|-------|-------|
| OBRIKOM | <i>V. amygdalina</i> | 0.005 | 0.010 | 0.013 | 0.005 | 0.290 | 0.145 |
| | <i>T. triangularare</i> | 0.013 | 0.0013 | 0.010 | 0.001 | 0.263 | 0.070 |
| | <i>A. esculentus</i> | 0.003 | 0 | 0.010 | 0.006 | 0.273 | 0.065 |
| | <i>T. occidentalis</i> | 0.005 | 0.003 | 0.014 | 0.002 | 0.302 | 0.128 |
| OMOKU | <i>V. amygdalina</i> | 0.008 | 0.006 | 0.022 | 0.002 | 0.310 | 0.120 |
| | <i>T. triangularare</i> | 0.008 | 0.002 | 0 | 0.008 | 0.281 | 0.155 |
| | <i>A.esculentus</i> | 0.002 | 0.002 | 0.012 | 0.006 | 0.322 | 0.050 |
| | <i>T. occidentalis</i> | 0.001 | 0.005 | 0.014 | 0.001 | 0.269 | 0.078 |
| OKPOSI | <i>V. amygdalina</i> | 0.001 | 0.005 | 0.027 | 0.002 | 0.205 | 0.094 |
| | <i>T. triangularare</i> | 0.001 | 0 | 0.018 | 0.002 | 0.212 | 0.141 |
| | <i>A. esculentus</i> | 0.010 | 0.002 | 0.016 | 0 | 0.240 | 0.040 |
| | <i>T. occidentalis</i> | 0.001 | 0.004 | 0.009 | 0.003 | 0.252 | 0.088 |
| ELLETA | <i>V. amygdalina</i> | 0.006 | 0.003 | 0.027 | 0.001 | 0.338 | 0.081 |
| | <i>T. triangularare</i> | 0.016 | 0.005 | 0.025 | 0.002 | 0.297 | 0.051 |
| | <i>A. esculentus</i> | 0.001 | 0.001 | 0 | 0.001 | 0.318 | 0.112 |
| | <i>T. occidentalis</i> | 0.004 | 0.005 | 0.034 | 0.009 | 0.249 | 0.142 |
| ❖ | TDI(mg day ⁻¹ person ⁻¹) | 0.00 | 0.000 | 0.50 | 6.3 x 10 ⁻⁸ | 20.5 | 0.9 |
| ❖ | UTDI(mg day ⁻¹ person ⁻¹) | 0.240 | 0.064 | 1.00 | 0.002 | 45.0 | 10.00 |

- ❖ Recommended tolerable daily intake (TDI) and upper tolerable daily intake (UTDI) levels of heavy metals in food stuffs (FDA, 2001; Garcia – Rico, 2007)

Estimated daily intake (EDI) of metals in (mg/kg) for children population

The results of the EDI for children are shown in Table 3. The concentration of Pb ranged between 0.001 to 0.041 mg/kg. Calculated EDI for Lead in children was highest in *T. Triangularare* (0.041 mg/kg) from Elleta community. The value of Cd ranged from 0.003 to 0.025 mg/kg with *V. amygdalina* from Obrikom community recording the highest value (0.025 mg/kg). Calculated EDI for Ni ranged between 0.023 to 0.085 mg/kg with the highest value recorded in *T. occidentalis* (0.085 mg/kg) from Elleta community. Calculated EDI for Fe in children ranged between 0.513 to 0.846 mg/kg with the highest

value recorded in *V.amygdalina* (0.846 mg/kg) from Elleta community. The EDI for Cu in children ranged between 0.100 to 0.388 mg/kg. Calculated EDI for Cu in children was highest in *T. triangulare* from Omoku community.

Table 3; Estimated daily intake (EDI) of heavy metals mg/kg for children(24kg) population in ONELG via consumption of vegetables

| LOCATION | SAMPLES | Pb | Cd | Ni | As | Fe | Cu |
|----------|------------------------|-------|-------|-------|----------------------|-------|-------|
| OBRIKOM | <i>V. amygdalina</i> | 0.014 | 0.025 | 0.032 | 0.011 | 0.724 | 0.363 |
| | <i>T. triangulare</i> | 0.032 | 0.003 | 0.025 | 0.003 | 0.656 | 0.176 |
| | <i>A. esculentus</i> | 0.006 | 0 | 0.024 | 0.015 | 0.683 | 0.163 |
| | <i>T. occidentalis</i> | 0.012 | 0.007 | 0.035 | 0.005 | 0.754 | 0.320 |
| OMOKU | <i>V. amygdalina</i> | 0.019 | 0.015 | 0.055 | 0.005 | 0.774 | 0.301 |
| | <i>T. triangulare</i> | 0.020 | 0.007 | 0 | 0.020 | 0.703 | 0.388 |
| | <i>A. esculentus</i> | 0.005 | 0.004 | 0.029 | 0.014 | 0.805 | 0.124 |
| | <i>T. occidentalis</i> | 0.003 | 0.012 | 0.023 | 0.003 | 0.673 | 0.194 |
| OKPOSI | <i>V. amygdalina</i> | 0.003 | 0.013 | 0.068 | 0.005 | 0.513 | 0.236 |
| | <i>T. triangulare</i> | 0.001 | 0 | 0.046 | 0.012 | 0.530 | 0.353 |
| | <i>A. esculentus</i> | 0.024 | 0.006 | 0.041 | 0 | 0.601 | 0.100 |
| | <i>T. occidentalis</i> | 0.002 | 0.011 | 0.023 | 0.006 | 0.630 | 0.219 |
| ELLETA | <i>V. amygdalina</i> | 0.016 | 0.006 | 0.068 | 0.003 | 0.846 | 0.203 |
| | <i>T. triangulare</i> | 0.041 | 0.013 | 0.062 | 0.004 | 0.741 | 0.128 |
| | <i>A. esculentus</i> | 0.002 | 0.003 | 0 | 0.001 | 0.796 | 0.279 |
| | <i>T. occidentalis</i> | 0.009 | 0.013 | 0.085 | 0.022 | 0.621 | 0.355 |
| ❖ | TDI(mg/day/person) | 0.00 | 000 | 50 | 6.3×10^{-8} | 20.5 | 0.9 |
| ❖ | UTDI (mg/day/person) | 0.240 | 064 | 00 | 0.002 | 45.0 | 10.00 |

- ❖ Recommended tolerable daily intake (TDI) and upper tolerable daily intake (UTDI) levels of heavy metals in food stuffs (FDA, 2001; Garcia – Rico, 2007)

Target hazard quotient (THQ) and hazard index (HI) of heavy metals for adult and children population

The THQ value greater than 1 indicates health concern. The non-carcinogenic risk of six heavy metals exposure to consumption of vegetables from four towns in Ogba/Egbema/Ndoni Local Government Area of Rivers State are presented in Tables 4 and 5. The THQ value for Pb ranged from 0.156 to 4.690 mg/kg with the highest value recorded in *T. triangulare* (4.690 mg/kg) from Elleta community in adult whereas calculated THQ value for Pb in children ranged between 0.390 to 11.726 mg/kg with *T. triangulare* (11.726 mg/kg) from Elleta community recording the highest. THQ value of Cd ranged between 1.205 to 9.937mg/kg and 3.012 to 24.838 mg/kg for adults and children respectively. However, *V. amygdalina*

from Obrikom community recorded the highest THQ values of 9.937 and 24.838 mg/kg for adults and children respectively. The THQ value for Ni ranged between 0.461 to 1.701 mg/kg and 1.151 to 4.252 mg/kg for adults and children respectively. However, the highest values in adults and children were seen in *T. occidentalis* (1.701 mg/kg and 4.252 mg/kg) both from Elleta community. The THQ value of As was within the range of 0.010 to 0.175 mg/kg and 0.025 to 0.438 mg/kg for adults and children respectively. The highest value was recorded in *T.occidentalis* (0.175 and 0.438 mg/kg) in both adults and children population respectively from Elleta community. The THQ value of Fe ranged between 0.293 to 0.483 mg/kg and 0.733 to 1.209 mg/kg for adults and children population respectively. The highest value was recorded in *V.amygdalina* (0.483 and 1.209 mg/kg) both from Elleta community in adults and children respectively. The THQ value of Cu was within the range 1.003 to 3.878 mg/kg and 2.507 to 9.693 mg/kg for adults and children population respectively. The highest value was recorded in *T. triangulare*(3.877 and 9.693 mg/kg) from Omoku community in both adults and children population respectively. On the other hand, the result of the Hazard Index (HI) of adults and children population from all the communities shown in Tables 4 and 5 were in the range 3.343 to 16.259 mg/kg and 8.357 to 40.645 mg/kg respectively. The highest Hazard Index (HI) value in adults was recorded in *V. amygdalina* (17.26 mg/kg) whereas the highest Hazard Index value in children was recorded in *V. amygdalina* (40.646 mg/kg) both from Obrikom community.

Table 4: Target Hazard Quotient (THQ) and Hazard index (HI) for adults exposed to vegetables contaminated with heavy metals in ONELGA

| LOCATION | SAMPLES | Pb | Cd | Ni | As | Fe | cu | Hazard Index (HI) |
|----------|------------------------|-------|-------|-------|-------|-------|-------|-------------------|
| OBRIKOM | <i>V. amygdalina</i> | 1.544 | 9.937 | 0.641 | 1.090 | 0.414 | 3.634 | 17.26 |
| | <i>T. triangulare</i> | 3.640 | 1.314 | 0.493 | 0.026 | 0.375 | 1.758 | 7.606 |
| | <i>A. esculentus</i> | 0.724 | 0 | 0.482 | 0.119 | 0.390 | 1.627 | 3.343 |
| | <i>T. occidentalis</i> | 1.314 | 2.804 | 0.695 | 0.040 | 0.431 | 3.203 | 8.487 |
| OMOKU | <i>V. amygdalina</i> | 2.185 | 6.001 | 1.089 | 0.043 | 0.443 | 3.009 | 12.775 |
| | <i>T. triangulare</i> | 2.259 | 2.763 | 0 | 0.159 | 0.402 | 3.877 | 9.460 |
| | <i>A. esculentus</i> | 0.535 | 1.645 | 0.577 | 0.108 | 0.460 | 1.236 | 4.562 |
| | <i>T. occidentalis</i> | 0.364 | 4.869 | 0.699 | 0.024 | 0.384 | 1.937 | 8.277 |
| OKPOSI | <i>V. amygdalina</i> | 0.389 | 5.098 | 1.357 | 0.037 | 0.293 | 2.355 | 9.529 |
| | <i>T.triangulare</i> | 0.156 | 0 | 0.919 | 0.098 | 0.303 | 3.533 | 5.009 |
| | <i>A. esculentus</i> | 2.757 | 2.405 | 0.821 | 0 | 0.343 | 1.003 | 7.330 |

| | | | | | | | | |
|--------|------------------------|-------|-------|-------|-------|-------|-------|--------|
| | <i>T.occidentalis</i> | 0.245 | 4.351 | 0.461 | 0.051 | 0.360 | 2.187 | 7.655 |
| ELLETA | <i>V. amygdalina</i> | 1.774 | 2.544 | 1.357 | 0.020 | 0.483 | 2.033 | 8.211 |
| | <i>T. triangulare</i> | 4.690 | 5.131 | 1.235 | 0.031 | 0.424 | 1.284 | 12.795 |
| | <i>A. esculentus</i> | 0.232 | 1.205 | 0 | 0.010 | 0.455 | 2.791 | 4.694 |
| | <i>T. occidentalis</i> | 1.075 | 5.122 | 1.701 | 0.175 | 0.355 | 3.551 | 11.979 |

THQ > 1 indicates health concerns whereas, THQ < 1 pose no risk to the health of any given population

Table 5; Target hazard quotient (THQ) and hazard index (HI) for children exposed to vegetables contaminated with heavy metals in ONELGA

| LOCATION | SAMPLES | Pb | Cd | Ni | As | Fe | Cu | Hazard Index (HI) |
|----------|------------------------|--------|--------|-------|-------|-------|-------|-------------------|
| OBRIKOM | <i>V. amygdalina</i> | 3.860 | 24.838 | 1.603 | 0.226 | 1.035 | 9.084 | 40.646 |
| | <i>T. triangulare</i> | 9.091 | 3.285 | 1.232 | 0.064 | 0.937 | 4.396 | 19.006 |
| | <i>A. esculentus</i> | 1.811 | 0 | 1.206 | 0.297 | 0.975 | 4.068 | 8.357 |
| | <i>T. occidentalis</i> | 3.284 | 7.009 | 1.739 | 0.091 | 1.077 | 8.008 | 21.208 |
| OMOKU | <i>V. amygdalina</i> | 5.462 | 14.999 | 2.722 | 0.107 | 1.106 | 7.522 | 31.918 |
| | <i>T. triangulare</i> | 5.647 | 6.907 | 0 | 0.397 | 1.004 | 9.693 | 23.649 |
| | <i>A. esculentus</i> | 1.338 | 4.113 | 1.443 | 0.270 | 1.150 | 3.091 | 11.405 |
| | <i>T. occidentalis</i> | 0.901 | 12.171 | 1.747 | 0.051 | 0.961 | 4.842 | 20.672 |
| OKPOSI | <i>V. amygdalina</i> | 0.972 | 12.742 | 3.392 | 0.093 | 0.733 | 5.887 | 23.820 |
| | <i>T. triangulare</i> | 0.390 | 0 | 2.297 | 0.245 | 0.757 | 8.832 | 12.522 |
| | <i>A. esculentus</i> | 6.894 | 6.012 | 2.052 | 0 | 0.859 | 2.507 | 18.323 |
| | <i>T.occidentalis</i> | 0.613 | 10.875 | 1.151 | 0.128 | 0.900 | 5.467 | 19.135 |
| ELLETA | <i>V. amygdalina</i> | 4.434 | 6.359 | 3.393 | 0.050 | 1.209 | 5.081 | 20.525 |
| | <i>T. triangulare</i> | 11.726 | 12.826 | 3.088 | 0.077 | 1.059 | 3.210 | 31.986 |
| | <i>A. esculentus</i> | 0.581 | 3.012 | 0 | 0.025 | 1.137 | 6.978 | 11.734 |
| | <i>T. occidentalis</i> | 2.688 | 12.803 | 4.252 | 0.438 | 8.888 | 8.878 | 29.946 |

THQ > 1 indicates health concerns whereas THQ < 1 pose no risk to the health of any given population

Carcinogenic risk (CR) for adult and children population

Carcinogenic Risk (CR) also known as Lifetime cancer risk (LCR) is the lifetime probability of an individual developing any type of cancer due to carcinogenic daily exposure to a contaminant over a life time. The LCR is obtained using the Cancer Slope Factor (CSF) which evaluates the probability of an individual developing cancer from oral exposure to contaminant levels over a period of a lifetime as described by ATSDR in 2010 and it is contaminant specific.²² Ingestion cancer slope factors are expressed in units of (mg/kg/day). US EPA, 2011 states that 10^{-6} (1 in 1,000,000) to 10^{-4} (1 in 10,000) represent a range of permissible predicted lifetime risks for carcinogens. Chemical

for which the risk factor falls below 10^{-6} may be eliminated from further consideration as a chemical of concern. The results in tables 6 and 7 shows the Life Cancer Risk of vegetables exposed to heavy metals contamination over a life time for adult and children population respectively. The LCR value for Pb, Cd, Ni, and As in adult population ranged from 8.5E-06 to 1.11E-04 mg/kg/day, 7.6E-04 to 0.004 mg/kg/day, 0.008 to 0.031 mg/kg/day, and 0.002 to 0.014 mg/kg/day respectively having their highest values at *T. triangulare* (1.11E-04 mg/kg day), *V. amygdalina* (0.004 mg/kg/day) both from Obrikom community, *T. occidentalis* (0.031 mg/kg/day), and *T. occidentalis*(0.014 mg/kg/day) both from Elleta community respectively. Meanwhile, the LCR for Pb, Cd, Ni, and As in children population ranges from (8.5E-06 to 1.02E-04 mg/kg/day), (0.001 to 0.095 mg/kg/day), (0.021 to 0.077 mg/kg/day), and (0.002 to 0.033 mg/kg/day) respectively having their highest values at *T. occidentalis* (1.02E-04 mg/kg/day), *V.amygdalina* (0.0095 mg/kg/day) both from Obrikom community, *T. occidentalis* (0.077 mg/kg/day), and *T. occidentalis* (0.033 mg/kg/day) both from Elleta community respectively.

TABLE 6: Life Cancer Risk (LCR) Table for Adult Population

| LOCATIONS | SAMPLES | Pb | Cd | Ni | As | Fe | Cu |
|-----------|------------------------|----------|----------|-------|-------|----|----|
| OBRIKOM | <i>V. amygdalina</i> | 4.25E-05 | 0.004 | 0.012 | 0.008 | - | - |
| | <i>T.triangulare</i> | 1.11E-04 | 4.94E-04 | 0.009 | 0.002 | - | - |
| | <i>A. esculentus</i> | 2.55E-05 | 0 | 0.009 | 0.009 | - | - |
| | <i>T. occidentalis</i> | 4.25E-05 | 0.001 | 0.013 | 0.003 | - | - |
| OMOKU | <i>V. amygdalina</i> | 6.8E-05 | 0.002 | 0.020 | 0.003 | - | - |
| | <i>T. triangulare</i> | 6.8E-05 | 7.6E-04 | 0 | 0.012 | - | - |
| | <i>A. esculentus</i> | 1.7E-05 | 7.6E-04 | 0.011 | 0.009 | - | - |
| | <i>T. occidentalis</i> | 8.5E-06 | 0.002 | 0.013 | 0.002 | - | - |
| OKPOSI | <i>V. amygdalina</i> | 8.5E-06 | 0.002 | 0.025 | 0.003 | - | - |
| | <i>T. triangulare</i> | 8.5E-06 | 0 | 0.016 | 0.003 | - | - |
| | <i>A. esculentus</i> | 8.5E-05 | 7.6E-04 | 0.015 | 0 | - | - |
| | <i>T.occidentalis</i> | 8.5E-06 | 0.002 | 0.008 | 0.005 | - | - |

| | | | | | | | |
|--------|------------------------|----------|---------|-------|-------|---|---|
| ELLETA | <i>V. amygdalina</i> | 5.1E-05 | 0.001 | 0.025 | 0.002 | - | - |
| | <i>T. triangulare</i> | 1.36E-04 | 0.002 | 0.023 | 0.003 | - | - |
| | <i>A. esculentus</i> | 8.5E-06 | 3.8E-04 | 0 | 0.002 | - | - |
| | <i>T. occidentalis</i> | 3.4E-05 | 0.002 | 0.031 | 0.014 | - | - |

Ingestion Cancer Slope Factor 0.0085 0.38 0.91 1.5000

US EPA, 2011 states that 10^{-6} (1 in 1,000,000) to 10^{-4} (1 in 10,000) represent a range of permissible predicted lifetime risks for carcinogens.

TABLE 7: Life Cancer Risk (LCR) Table for Children Population

| LOCATIONS | SAMPLES | Pb | Cd | Ni | As | Fe | Cu |
|-----------|------------------------|----------|--------|-------|-------|----|----|
| OBRIKOM | <i>V. amygdalina</i> | 1.19E-04 | 0.0095 | 0.029 | | | |
| | <i>T. triangulare</i> | 2.72E-04 | 0.001 | 0.023 | 0.005 | - | - |
| | <i>A. esculentus</i> | 5.1E-05 | 0 | 0.022 | 0.023 | - | - |
| | <i>T. occidentalis</i> | 1.02E-04 | 0.003 | 0.032 | 0.008 | - | - |
| OMOKU | <i>V. amygdalina</i> | 1.62E-04 | 0.006 | 0.050 | 0.008 | - | - |
| | <i>T. triangulare</i> | 1.7E-04 | 0.003 | 0 | 0.03 | - | - |
| | <i>A. esculentus</i> | 4.25E-05 | 0.002 | 0.026 | 0.021 | - | - |
| | <i>T. occidentalis</i> | 2.55E-05 | 0.005 | 0.021 | 0.005 | - | - |
| OKPOSI | <i>V. amygdalina</i> | 2.55E-05 | 0.005 | 0.062 | 0.008 | - | - |
| | <i>T. triangulare</i> | 8.5E-06 | 0 | 0.056 | 0.018 | - | - |
| | <i>A. esculentus</i> | 2.04E-04 | 0.002 | 0.037 | 0 | - | - |
| | <i>T. occidentalis</i> | 1.7E-05 | 0.004 | 0.021 | 0.009 | - | - |
| ELLETA | <i>V. amygdalina</i> | 1.36E-04 | 0.002 | 0.062 | 0.005 | - | - |
| | <i>T. triangulare</i> | 3.49E-04 | 0.005 | 0.056 | 0.006 | - | - |
| | <i>A. esculentus</i> | 1.75E-05 | 0.001 | 0 | 0.002 | - | - |
| | <i>T. occidentalis</i> | 7.65E-05 | 0.005 | 0.077 | 0.033 | - | - |

Ingestion Cancer Slope Factor 0.0085 0.38 0.91 1.5000

US EPA, 2011 states that 10^{-6} (1 in 1,000,000) to 10^{-4} (1 in 10,000) represent a range of permissible predicted lifetime risks for carcinogens.

Discussion

Heavy metals contamination in soil from various anthropogenic sources such as Industrial wastes, waste water irrigation, air deposition, oil spillage from oil exploitation, automobile emissions, mining activity, and agricultural practices are believed to easily accumulate in the top soil which results to bioaccumulation, biomagnification and geoaccumulation of the heavy metals in different parts of the plants especially the vegetables.^{3,23,24} As a result of sewage water irrigation, high level of oil exploitation, application of phosphate fertilizers, sewage sludge application, etc. Accumulation of heavy metals in vegetables may be increased.²⁵ This can easily enter the food chain thereby causing different diseases. The intake of these edible vegetables contaminated with heavy metals may result to intoxication that can be described as acute or when the disease appears after a potential period of time and long term or chronic intoxications.⁹ It was described in previous studies that leafy vegetables have higher concentration of metals as compared to bulbs and tuber vegetables due to smoke emit from industries present in vicinity of vegetables.²⁶ Ingestion of contaminated food is the main route of human exposure to these toxic compounds accounting for more than ninety percent compared to other routes.^{11,12} Higher concentration of these heavy metals is the outcome of industrial pollution, oil exploitation, road runoff, food, wear of tires, hospital waste, atmospheric deposition and burning of fuels.²⁷ The present study has highlighted the contamination profile of vegetables commonly consumed from selected communities in Ogba/Egbema/Ndoni Local Government Area in Rivers State, Nigeria and the possible public health implications. The concentrations of Pb in *T. triangulare* (3.824 mg/kg) from Obrikom community, *V. amygdalina* (2.294 mg/kg), *T. triangulare* (2.373 mg/kg) from Omoku community, *A. esculentus* (2.894 mg/kg) from Okposi community and *T. triangulare* (4.908 mg/kg) from Elleta community, respectively were higher than the 2.0 mg/kg value as recommended by FAO/WHO.²⁸ Almost all samples analyzed for Pb were above the 0.3 mg/kg as recommended by European communities.²⁹ All the analyzed samples of Pb were higher than the 0.2 mg/kg.³⁰ except for *T. triangulare* (0.164 mg/kg) from Okposi community. These were similar to the results obtained by Asdeo and Loonker,³¹ where the Pb accumulation in vegetables was found to be within the range of 2.32 to 5.76 mg/kg. The high levels of Pb in these vegetables may probably be

attributed to pollutants in irrigation water, farm soil (farm site) contaminated by crude oil from oil exploitation or due to pollution from the highways traffic.^{32,23} Besides, high concentration of Pb could be due to increase in mobile metal fraction of Pb and high level of oil exploitation.³³ With higher metal fraction and high level of oil exploitation, uptake and accumulation of metal in vegetable increase. There were lower levels of Pb in *T. triangulare* (0.168 mg/kg), *T. occidentalis* (0.255 mg/kg) both from Okposi community and *A. esculentus* (0.234 mg/kg) from Elleta community as reported by Agrawal.³⁴ Lower levels of Pb have been reported in *T. occidentalis* by Akinola and Ekiyoyo.³⁵ Lead (Pb) is known to induce reduced cognitive development and intellectual performance in children and cardiovascular disease and increased blood pressure in adults.²⁹ Generally, Pb contaminations occur in vegetables grown on contaminated soils, through air deposition, through sewage sludge/waste water application or through oil exploitation.²³ In human body, Pb causes neurological, hematological and physiological disorders.²⁷ Lead (Pb) influences the nervous system, slowing down nervous response. This influences learning abilities and behaviour.²⁴ The concentration of Cd in all the samples from the four communities exceeded the permissible limit of 1.0 mg/kg and 0.2 mg/kg.^{29,36} except for *A. esculentus* and *T. triangulare* from Obrikom and Okposi community respectively, which were below detectable limit (BDL < 0.001). Lower values have been reported for leafy vegetables cultivated along road sides (0.27 mg/kg) and leafy vegetables obtained from four markets (0.20 mg/kg) in Lagos State.^{23,37} Low level of cadmium may be attributed to low or no use of phosphate fertilizers by the farmers on the planted vegetables. The higher level of cadmium in the present study may be attributed to high use of phosphate fertilizers by the farmers in the study area which tend to increase the accumulation of cadmium in the soil. Cadmium is a heavy metal with high toxicity and it is a non-essential element in foods and natural waters and it accumulates principally in the kidneys and liver.^{38,24} Accumulation of Cd in human body leads to certain disorder including cardiovascular diseases, liver and nervous system.^{38,24} The most common sources of Cd in plants and vegetables are sewage sludge application, deposition from fossil fuel combustion, phosphate fertilizers etc.²⁴. Cadmium accumulates especially in the kidneys leading to dysfunction of the kidney with increased secretion of e.g. proteins in urine (proteinuri) and other effects.³⁹ The highest level of Ni was recorded in *T. occidentalis* (10.241 mg/kg) from Elleta community while *T. occidentalis* (2.780 mg/kg) from

Okposi community recorded the least Ni concentration. In this study, Ni concentration in all the analyzed samples exceeded the permissible limit of 2.7 mg/kg recommended by NAFDAC except *T. triangulare* and *A. esculentus* from Omoku and Elleta communities respectively which were below detectable limit (BDL < 0.001). Also Ni concentration in vegetables were lower than the estimated maximum guideline set by USFDA of 70-80mg/g.⁴⁰ This was contrary to Adedokun, *et al.*³⁷ who reported a lower level of Ni, this could be attributed to lack of oil exploitation in that area as compared with the high rate of oil exploitation by oil companies in ONELGA. Inhaled Ni carbonyl, a carcinogenic gas that results from the action of nickel with heated carbon monoxide from cigarette smoke, car exhaust, and some industrial wastes is very toxic and Ni allergy can also cause systematic reactions.⁴¹ The concentration of the metalloid, As in over 80% of the samples from all the selected communities exceeded the permissible limit of 0.43mg/kg recommended by WHO/FAO.²⁸ This was in line with Jun Yang, *et al.*⁴² who reported high level of As concentration in the soil of Daye city of Hubei Province, China. This may be attributed to the contamination of the soil by oil spillage caused by excessive oil exploitation by oil companies, pipeline vandalization etc. Lower levels of As were obtained from *T.triangulare* (0.385 mg/kg), *T.occidentalis* (0.374 mg/kg), *V.amygdalina* (0.297 mg/kg) and *A.esculentus* (0.157 mg/kg) from Obrikom, Omoku and Elleta communities respectively, while *A.esculentus* from Okposi community was below detectable limit (BDL < 0.001). Arsenic has been classified as a human carcinogen, with chronic ingestion associated with lung cancer.⁴³ It has been suggested that exposure to arsenic-rich food increases the risk of cardiovascular disease, hypertension and diabetes mellitus.⁴⁴ Iron is essential for the synthesis of chlorophyll and activates a number of respiratory enzymes in plants. The deficiency of Fe results in severe chlorosis of leaves in plants. High levels of exposure to Fe dust may cause respiratory diseases such as chronic bronchitis and ventilation difficulties. The Fe contents of these vegetables in the present study were lower than the FAO/WHO (2001) safe limit of 425.00 mg/kg. These vegetables could be good supplement for Fe. From the six heavy metals analyzed, the concentration of Fe in *V. amygdalina* (101 mg/kg) was the highest, suggesting that Fe is more susceptible to the interference of vegetable species, pollution and other external factors from the present study. The concentration of Cu in all the

samples in this study area within the permissible level of 73.00 mg/kg by the FAO/WHO in vegetables. Also 80% of all the samples from the four communities were within the suggested safe limits of 40mg/kg recommended by joint Expert Committee for Food Additives.⁴⁵ for foodstuffs except for *V.amygdalina* (43.604 mg/kg) from Obrikom community, *T. triangulare* (46.536 mg/kg) from Omoku community, *T. triangulare* (42.394 mg/kg) from Okposi and *T. occidentalis* (42.611 mg/kg) from Elleta communities which were slightly higher than the suggested safe limits of 40mg/kg as recommended by JECFA, (2005).⁴⁵ However the Cu levels obtained from this present study were higher than that of Adedokun, *et al.*,³⁷ where the concentration of Cu ranged between (2.338 mg/kg) in *T. occidentalis* to (14.075 mg/kg) in *T.triangulare* in the vegetables they worked with. Copper is essential to human life as metalloproteins and function as enzymes, however, critical doses leads to health risks such as anemia, diabetes, inflammation, kidney and liver dysfunction and vitamin c deficiency.⁴⁶ Although toxicity is rare, it's metabolism is enhanced by Molybdenum and Zinc constituents in the body.⁴⁷.

Estimated daily intake (EDI) of metals in (mg/kg) for adult population

The EDI results in adults shown in Table 2 were compared with the recommended tolerable daily intake of metals (TDI) and the upper tolerable daily intake level (UTDI) established by the Institute of medicine for people between the ages of 18 to 70 years.⁴⁸ It is very clear from Table 2 which represents the EDI of these metals in vegetables species in adult in Pb (0.001 to 0.016 mg/kg) and Cd (0.001 to 0.005 mg/kg) exceeded the recommended tolerable daily intake of metals (TDI) but fall within the upper tolerable daily intake (UTDI). However, Ni (0.009 to 0.034 mg/kg), Fe (0.205 to 0.338 mg/kg) and Cu (0.040 to 0.155 mg/kg) were lower than the recommended tolerable daily intake of metals (TDI) and the upper tolerable daily intake level (UTDI). Arsenic (0.001 to 0.009 mg/kg) exceeded the recommended tolerable daily intake of metals (TDI) and of the analyzed samples from different locations such as *V. amygdalina* (0.005 mg/kg), *A.esculentus* (0.006 mg/kg) both from Obrikom community, *T. triangulare* (0.008 mg/kg), *A. esculentus* (0.006 mg/kg) both from Omoku community, *T. occidentalis* (0.003 mg/kg) from Okposi community and *T. occidentalis* (0.009 mg/kg) from Elleta community exceeded the upper

tolerable daily intake level (UTDI) whereas the other analyzed samples fall within the upper tolerable daily intake level (UTDI).

Estimated daily intake (EDI) of metals in (mg/kg) for children population

From Table 3, which represents the EDI of these metals in vegetables species in children for Pb (0.001 to 0.041 mg/kg) and Cd (0.003 to 0.025 mg/kg) exceeded the recommended tolerable daily intake of metals (TDI) but fall within the upper tolerable daily intake level (UTDI). However, just like the EDI in adults, Ni (0.023 to 0.85 mg/kg), Fe (0.513 to 0.84 mg/kg), and Cu (0.100 to 0.388 mg/kg) were lower than the recommended tolerable daily intake (TDI) and the upper tolerable daily intake level (UTDI), except for *A. esculentus* (0.001 mg/kg) from Elleta community which falls within the upper tolerable daily intake level (UTDI).

Target hazard quotient (THQ) of heavy metals for adult and children population

The THQ is a ratio between the measured concentration and the oral reference dose, weighted by the length and frequency of exposure; amount ingested and body weight.⁵⁰ From the results in the present study for adult population, the concentrations of Pb in most of the samples in adults most notably in *T. triangulare* (4.690 mg/kg) from Elleta community, Cd in almost all the samples were greater than 1 except for *A. esculentus* from Obrikom community which was below detectable limit (BDL < 0.001) and it is highest in *V. amygdalina* (9.937 mg/kg) from Obrikom community. Nickel concentrations were mostly less than 1 but highest in *T. occidentalis* (1.701 mg/kg) from Elleta community. The concentrations of As and Fe were less than 1 which indicates that the population may not be at risk of As and Fe contamination and toxicity. The concentrations of Cu in all the analyzed samples were all greater than 1 most notably in *V. amygdalina* (3.634 mg/kg) from Obrikom community in adult population. The result for THQ for the children population in the present study shows that Pb concentrations were mostly greater than 1 and is highest in *T. triangulare* (11.726 mg/kg) from Elleta community. The concentrations of Cd in most of the samples were greater than 1 most notably in *V. amygdalina* (24.838 mg/kg) from Obrikom community. The concentrations of Ni in the analyzed samples were mostly greater than 1 and highest in *T. occidentalis* (4.252 mg/kg) from Elleta community. The concentrations of As

in the analyzed samples were all less than 1 which indicates that the population may not be at risk of As contamination from the consumption of vegetables in the present study. The concentrations of Fe in all the analyzed samples were mostly greater than 1 and a good portion of Fe concentrations in the samples were less than 1 and was highest in *V.amygdalina* (1.209 mg/kg) from Elleta community. The concentrations of Cu in the analyzed samples were all greater than 1 and is highest in *T. triangulare* (9.693 mg/kg) from Omoku community. From the result of the present study for adult population, the concentrations of Cu, Cd, and Pb in all the analyzed samples were mostly greater than 1, which indicates that the adult population of Ogba/Egbema/Ndoni Local Government Area in Rivers State may be at risk of either Cu, Cd, and Pb contamination and toxicity. The concentrations of As and Fe in the analyzed samples were all less than 1 which indicates that the adult population may not be at risk of As and Fe contamination and toxicity in the present study, also Ni concentrations in most of the analyzed samples were mostly less than 1, which also indicates that Ni may not be a metal of concern for the adult population in the present study. The adult and children population were highly exposed to health risks associated to these metals in the order Cu>Cd>Pb>Ni>Fe>As for the adult population and Cu>Cd>Ni>Pb>Fe>As for the children population in the present study. In the present study, the result of the THQ in children population shows that the concentrations of Cu, Cd, Ni, Pb, and slightly Fe were mostly greater than 1, which indicates level of concern that the children population may be at risk of either Cu, Cd, Ni, Pb, and slightly Fe contamination and toxicity from vegetables consumption. The concentrations of As in all the samples were all less than 1 which indicates that there may not be health concern of As contamination and toxicity for the children in the present study. Generally, the present study has shown that both the adult and children population have a THQ greater than 1 which indicates that there may be health concerns for the adult and children population. Similar THQ for Pb, Cd, and Ni were reported by Singh *et al.*³ in vegetables from waste water irrigated area. Similar THQ for Pb and Antimony (Sb) smelter in Nanning, China.. In conclusion, this study has highlighted the various concentrations of the heavy metals of interest in various vegetable samples and it has shown that out of the six metals analyzed, only the metalloid, As in both adults and children population have their Target Hazard Quotient (THQ) less than 1 which indicates that the exposed population may not be at risk of As contamination and toxicity. The other five metals namely: Pb, Cd, , Ni, Fe and Cu concentrations in the

samples were all greater than 1 which indicates that the exposed population in Ogba/Egbema/Ndoni Local Government Area in Rivers State may be at risk of Cu, Cd, Ni, Pb and Fe contamination and toxicity due to the frequent consumption of these edible vegetables.

Hazard index (HI) of metals for adult and children population

The HI in all the vegetable samples from all the locations ranges from (3.343 to 17.26 mg/kg) and (8.357 to 40.646 mg/kg) in adults and children population respectively. From this study, the Hazard Index (HI) is far greater than 1 ($HI > 1$). Generally $HI < 1$ means that the exposed population is safe from metals health risk while $HI > 1$ means the population is in danger or at risk of heavy metals contamination and toxicity.⁴⁹ The population is therefore at greater risk of Pb, Cd, Ni, As, Fe, and Cu as also reported by Tsafe *et al.*⁵⁰ Generally, It was observed that the health risk for children was higher than the adults which could be due to body weight.

Life Cancer Risk (LCR) for adult population

The Life Cancer Risk (LCR) result in adult population shown in table 6, shows that the LCR of Pb which ranged from (8.5×10^{-6} to 1.11×10^{-4} mg/kg/day) were all within the range of permissible predicted lifetime risks for carcinogens as suggested by USEPA, this indicates that there may be risk of developing any type of cancer due to carcinogenic ingestion and exposure to Pb over time. The LCR of Cd which ranges from (7.6×10^{-4} to 0.004 mg/kg/day) shows that almost all the analyzed vegetable samples from the four different locations exceeded the range of permissible predicted lifetime risks for carcinogens as suggested by USEPA which indicates that there may not be risk of developing any type of cancer due to the carcinogenic ingestion and exposure to Cd over time in the analyzed samples from these locations except for *T. triangulare* (4.94×10^{-4} mg/kg/day) from Obrikom community, *T. triangulare* (7.6×10^{-4} mg/kg/day), *A. esculentus* (7.6×10^{-4} mg/kg/day) from Omoku community, *A. esculentus* (7.6×10^{-4} mg/kg/day) from Okposi community and *A. esculentus* (3.8×10^{-4} mg/kg/day) from Elleta community were within the range of permissible predicted lifetime risks for carcinogens as suggested by USEPA, this indicates that there may be risk of developing any type of cancer due to carcinogenic ingestion and exposure to Cd over time from these analyzed from these locations. The LCR of Ni and As all exceeded

the range of permissible predicted lifetime risks for carcinogens as suggested by USEPA which indicates that there may not be risk of developing any type of cancer due to carcinogenic ingestion and exposure to Ni and As over time due to carcinogenic ingestion and exposure to Ni and As overtime.

Life Cancer Risk (LCR) for Children population

The Life Cancer Risk (LCR) result in children population shown in table 7, shows that the LCR of Pb which ranged from (8.5 E-06 to 1.02 E-04 mg/kg/day) were all within the range of permissible predicted lifetime risks for carcinogens as suggested by USEPA, this indicates that there may be risk of developing any type of cancer due to carcinogenic ingestion and exposure to Pb over time. The LCR of Cd, Ni, and As all exceeded the range of permissible predicted lifetime risks for carcinogens as suggested by USEPA which indicates that there may not be risk of developing any type of cancer due to carcinogenic ingestion and exposure to Cd, Ni, and As over time due to carcinogenic ingestion and exposure to Cd, Ni, and As overtime.

References

1. Heidarieh, M., Maragheh, M.G., Shamami, M.A., Behgar, M., Ziaei F., Alvaro, Z (2013). *Evaluation of heavy metal concentration in shrimp (Penaeus semisulcatus) and crab (Portunus pelagicus) with INAA method. Springerplus 2: 72.*
2. Kachenko, A.G., Singh, B. (2006). *Heavy metals contamination in vegetables grown in urban and metal smelter contaminated sites in Australia. Water Air Soil Pollut 169: 101-123.*
3. Singh, A., Sharma, R.K., Agarwal, M., Marshall, F.M. (2010). *Risk assessment of heavy metal toxicity through contaminated vegetables from waste water irrigated area of Varanasi, India. Trop Ecol 51: 375-387.*
4. Hansen, B; Alroe, HF; Kristensen, ES; Wier, M (2002), *Assessment of food safety in organic farming, Danish Research Centre for Organic Farming (DARCOF), Working papers, (52):1-24*
5. Gyawali, R; Ibrahim, SA; Abu Hasfa, SH; Smqadri, SQ; Haik, Y (2011), *Antimicrobial activity of copper alone and in combination with lactic acid against Escherichia coli O157:H7 in laboratory medium and on the surface of lettuce and tomatoes. Journal of Pathogens, 11: 1 - 9.*
6. Gupta, U.C., Subhas, C., Gupta, M.D. (2008). *Selenium in soils and crops, its deficiencies in livestock and humans: Implications for management. Commun Soil Sci Plant Anal 29: 1791-1807.*
7. WHO (1995). *Inorganic lead. Geneva, World Health Organization, International Programme on Chemical Safety. Environmental Health Criteria 165.*
8. Chailapakul, O., Korsrisakul, S., Siangproh, W., Grudpan, K.(2007). *Fast and simultaneous detection of heavy metals using a simple and reliable microchip-electrochemistry route: An alternative approach to food analysis. Atalanta 74: 683-689.*

9. Shahid, M., Dumat, C., Pourrut, B., Abbas, G., Shahid, N., Pinelli, E. (2015a). Role of metal speciation in lead-induced oxidative stress to *Vicia faba* roots. *Russian Journal of Plant Physiology* 62(4), 4484-454.
10. Shahid, M., Khali, S., Abbas, G., Shahid, N., Nadeem, M., Sabir, M., Aslam, M., Dumat, C. (2015b). Heavy Metal Stress and Crop Productivity. In: Hakeem, K. R., (Ed.), *Crop Production and Global. Environmental Issues. Springer*, pp. 1-25.
11. Uzu, G., Sauvain, J.J., Baeza-Squiban, A., Riediker, M., Sánchez, Sandoval Hohl M., Val, S., Tack, K., Denys, S., Pradere, P., Dumat, C. (2011a). In vitro assessment of the pulmonary toxicity and gastric availability of lead-rich particles from a lead recycling plant. *Environmental Science & Technology* 45(18), 7888-7895.
12. Uzu, G., Sobanska, S., Sarret, G., Sauvain, J.J., Pradere, P., Dumat, C. (2011b). Characterization of lead-recycling facility emissions at various workplaces: Major insights for sanitary risks. assessment. *Journal of Hazardous Materials* 186(2), 1018-1027.
13. Yargholi, B., Azim, A., Baghvand, A., Liaghat, A., Fardi, G. (2008). Investigation of cadmium absorption and accumulation in different parts of some vegetables. *American-Eurasian Journal of Agricultural & Environmental Sciences* 3(3), 357-364.
14. Wilson, B., Pyatt, F.B. (2007). Heavy metal dispersion, persistence, and bioaccumulation around an ancient copper mine situated in Anglesey, UK. *Ecotoxicol Environ Saf* 66: 224-231.
15. Muchuweti, M., Birkett, J.W., Chinyanga, E., Zvauya, R., Scrimshaw M.D., Lester, J. N. (2006). Heavy metals content of vegetables irrigated with mixture of waste water and sewage sludge in Zimbabwe: Implication for human health. *Agricul Ecos Environ* 112: 41-48.
16. Ahiamadu, A. (1993). *Ogba kingdom population*
17. Chary, N.S., Kamala, C.T., RAJ, D.S.S. (2008). Assessing risk of heavy metals from consuming food grown on sewage irrigated soils and food chain transfer. *Ecotoxicology and Environmental Safety*, 69: 513-524.
18. Wang, X., Sato, T., Xing, B., Tao, S. (2005). Health risks of heavy metals to the general public in Tianjin, China via consumption of vegetables and fish. *Science and Total Environment*, 350: 28-37
19. Storelli, MM (2008). Potential human health risks from metals (Hg, Cd, and Pb) and polychlorinated biphenyls (PCBs) via seafood consumption: Estimation of target hazard quotients (THQs) and toxic equivalents (TEQs). *Food Chemistry and Toxicology*, 46: 2782-2788.
20. Jan, F.A., Ishaq, M., Khan, S., Ihsanullah, I., Ahmad, I., Shakirullah, M. (2010). A comparative study of human health risks via consumption of food crops grown on wastewater irrigated soil (Peshawar) and relatively clean water irrigated soil (lower Dir). *Journal of Hazard Materials*, 179: 612-621.
21. Gerba, CP (2000). Risk Assessment: Environmental Microbiology. *Academic Press, London*
22. Baird, C. 2002. *Environmental Chemistry. Bookman, Porto Alegre, RS, Brazil (in Portuguese)*.
23. Oluwole, SO; Makinde, OSC; Yusuf, KA; Fajana, OO; Odumosu, AO (2013), Determination of Heavy Metal Contaminants in Leafy Vegetables Cultivated By the Road Side, *International Journal of Engineering Research and Development*, 7(3):1-5.
24. Adesuyi, AA; Njoku, K.L; Akinola, MO (2015). Assessment of Heavy Metals Pollution in Soils and Vegetation around Selected Industries in Lagos State, Nigeria. *Journal of Geoscience and Environment Protection*, 3: 11-19.
25. Devkota, B., Schmidt, G. (2000). Accumulation of heavy metals in food plants and grasshoppers from the Taigetos Mountains, Greece. *Agriculture, ecosystems & environment* 78(1), 85-91.
26. Mahmood A. Malik RN. (2014) Human health risk assessment of heavy metals via consumption of contaminated vegetables collected from different irrigation sources in Lahore, Pakistan. *Arabian Journal of Chemistry* 7(1), 91-99.
27. Sörme, Lagerkvist R. (2002). Sources of heavy metals in urban wastewater in Stockholm. *Science of the Total Environment* 298 (1), 131-145.

28. FAO/WHO (2001). *Report on the 32nd Session of the Codex Committee on Food and Contaminants, ALINORM 01/12, Beijing, China, 20–24 March 2000.*
29. Codex Alimentarius Commission (2001). *Food additives and contaminants Joint FAO/WHO Food Standard programme; ALINORM 01/12A:1-289.*
30. Chinese Department of Preventive Medicine. *Threshold for Food Hygiene. Beijing: China Standard Press; 1994.* (in Chinese)
31. Anjula Asdeo and Sangeeta Loonker, 2011. A Comparative Analysis of Trace Metals in Vegetables. *Research Journal of Environmental Toxicology*, 5: 125-132.
32. Qui XX, Huang DF, Cai SX, Chen F, Ren ZG, Cai YC (2000). Investigations on vegetables pollution and pollution sources and its control in Fuzhou, Fujian Province. *Fujian J. Agric. Sci.* 15: 16-21.
33. Siebe, C., 1995. Heavy metal availability to plants in soil irrigated with waste water from Mexico city. *Wat. Res.* 32 (12), 29–34.
34. Agrawal, G.D., 1999. Diffuse agricultural water pollution in India. *Wat.Sci. Technol.* 39 (3), 33–47.
35. Akinola, MO; Ekiyoyo, TA (2006). Accumulation of lead, cadmium and chromium in some plants cultivated along the bank of river Ribila at Odo-nla area of Ikorodu, Lagos state, Nigeria. *Journal of Environmental Biology*, 27(3):597-599.
36. Maleki, A; Zarasvand, MA (2008), Heavy Metals in Selected Edible Vegetables and Estimation of their Daily Intake in Sanandaj, Iran, *The Southeast Asian Journal of Tropical Medicine and Public Health*, 39(2): 335 – 340.
37. Adedokun, A. H; Njoku, K. L; Akinola, M. O; Adesuyi, A. A; Jolaoso, A. O. (2016). *Environmental Biology Research Unit, cell Biology and Genetics Department, University of Lagos, Aloha, Lagos, Nigeria.*
38. Divrikli, U; Horzum, N; Soylak, M; Elci, L (2006) Trace heavy metal contents of some spices and herbal plants from western Anatolia, Turkey. *International Journal of Food Science and Technology*, 41: 712-716
39. Waalkes, MP (2000). Cadmium carcinogenesis in review, *Journal of Inorganic Biochemistry*, 79: 240-244.
40. Iwegbue, CMA (2010). Composition and Daily Intakes of Some Trace Metals from Canned Beers in Nigeria, *Journal of Institute of Brewing and Distilling*, 116(3): 312–315.
41. World Health Organization (2004). Evaluation of certain food additives and Contaminants. In: *Sixty-First Report of the Joint FAO/WHO Expert on Food Additives. WHO, Geneva, Switzerland. (WHO Technical Series, 922).*
42. Jun, Yang; Silu, Ma; Jingcheng Zhou; Yongwei Song; Fei, Li; (2017). Heavy metal contamination in soil and vegetables and health risk assessment of inhabitants in Daye, China, Research center for environment and health, 124452 Hongnan University of Economics and Law2 school of information and safety. *Journal of International Medical Research*. 0 (0) 1-14
43. Kottutt, Jepkoech; "Arsenic Biosand filters in developing countries" (2015). *Honors projects*. 464. https://scholarworks.gvsu.edu/honors_projects/464
44. Liu, Li; Zhou, Hang; Wen-Tao, Yang; Jiao-Feng, Cui and Bo-Han, Liao. Arsenic in vegetables poses a health risk in the vicinity of a mining area in the Southern Human Province, China, Human and Ecological Risk Assessment : *An International Journal* (2017)
45. JECFA (2005), Joint FAO/WHO Expert committee on Food additives, 64th Meeting, JECFA/64/SC, *Codex Standard* 193-1995.
46. Lokeshappa, B; Shivpuri, K; Tripathi, V; Dikshit, AK (2012), Assessment of Toxic Metals in Agricultural Produce, *Food and Public Health*, 2(1): 24-29.
47. Oladele, A.T; Fadare, OO (2015). Heavy metals and proximate composition of forest leafy vegetables in oil producing area of Nigeria, *Ethiopia Journal of Environmental Studies and Management*, 8(4):451-463.
48. Institute of Medicine, Food and Nutrition Board (2001) [Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc.](#) *National Academy Press. Washington, USA.*

49. Khan, S; Lin, A; Zhang, S; Huc, Q; Zhu, Y. (2008), Accumulation of polycyclic aromatic hydrocarbons and heavy metals in lettuce grown in the soils contaminated with long-term waste water irrigation. *Journal of Hazardous Material*, 152: 506-515.
50. Tsafe A. I; Hassan, LG; Sahabi, D. M; Alhassan, Y; Bala B. M (2012), Evaluation of Heavy Metals Uptake and Risk Assessment of Vegetables Grown in Yargalma of Northern Nigeria. *Journal of Basic and Applied Science Research*, 2(7): 6708-6714