

## Original research papers

# Evaluation of Galantamine, Phenolics, Flavonoids and Antioxidant Content of *Galanthus* Species in Turkey

---

### ABSTRACT

**Aims:** The aim of the present study was to determine the total phenolic and flavonoid content and antioxidant activities in *Galanthus* species (*Galanthus woronowii*, *Galanthus nivalis*, and *Galanthus elwesii*) indigenous to Turkey.

**Study design:** The plant materials used in the study, *Galanthus elwesii* Ibradi samples were collected in Antalya province, *Galanthus nivalis* samples were collected in Istanbul province, and *Galanthus woronowii* samples were collected in Çaykara, Trabzon province in September 2018..

**Place and Duration of Study:** Plant samples were stored in Herbarium Material Warehouse at Afyon Kocatepe University. The plant leaves and grated bulbs were dried in an incubator at 60°C. The bulb and leaf samples were then pulverized to 80 mesh particle size for analysis.

**Methodology:** Total phenolic content was determined spectrophotometrically with Folin-Ciocalteu procedure and calculated as gallic acid equivalent (GAE). Total flavonoid content was determined with aluminum chloride colorimetric method and calculated as catechin equivalent (CAE). Antioxidant activities were determined with TEAC (Trolox equivalent antioxidant capacity) and DPPH (diphenyl-p-picrylhydrazyl radical) methods. The phenolic acid and galantamine content were determined by reversed phase HPLC.

**Results:** The highest total flavonoid content was determined as 33 mg CAE/g DW in *Galanthus woronowii* leaves and as 27 mg CAE / g DW in bulbs. DPPH removal activity was 77% in 500 µg/mL *Galanthus woronowii* leaf extract concentration and 93% in the ascorbic acid control group. The highest antioxidant content was observed in the leaves of *Galanthus woronowii* as 23 µmol Trolox/100 g DW and as 21 µmol Trolox/100 g DW in the bulbs. Higher galantamine content was determined in aerial parts (leaves) when compared to the underground parts (bulbs). The galantamine content in the leaves of all three *Galanthus* species was about 0.070%. The galantamine content in the bulbs of all three species was about 0.040%. Gallic, protocatechic, vanilic, caffeic, syringic, rosmarinic acid and catechin were identified in the leaves and bulbs of the three species with HPLC phenolic acid analysis. It was determined that the major phenolic acid was gallic acid.

**Conclusion:** The present study findings demonstrated that *Galanthus* species has antioxidant capacity. *Galanthus* spp. leaves had higher antioxidant activity when compared to the bulbs. *Galanthus woronowii* species exhibited the highest antioxidant activity among the scrutinized species.

*Keywords: Galanthus spp, Phenolic, Flavonoid, Antioxidant.*

## 1. INTRODUCTION

The interest in natural antioxidants, especially those found in fruits and vegetables, have been increasing among consumers and scientists during recent years. Epidemiological studies demonstrated that frequent fruit and vegetable intake is associated with a lower age-related coronary heart disease [1] and cancer risks [2,3]. Several natural nutrients contain dietary antioxidants that could scavenge free radicals. Certain studies demonstrated that phenolics such as flavonoids, phenolic acids, and tannins are more powerful antioxidants when compared to vitamins C and E [4]. These phenolic compounds also serve various biological functions, including anti-inflammatory, anti-carcinogenic and anti-atherosclerotic activities, which may be related to their antioxidant activities [5]. Several studies demonstrated a high degree of correlation between the total antioxidant activity of certain fruits and their phenolic content [6].

In addition to fruit and vegetable-based antioxidants, another important source of antioxidants is traditional medicinal plants, which could exhibit stronger antioxidant activities when compared to traditional dietary plants [7,8]. The Amaryllidaceae family is the most important plant family among 20 plant families that contain alkaloid, and it includes 85 genera and about 1100 perennial bulb species. These plants are distributed in the warm climates and tropical regions in the world [9]. It was reported that *Galanthus* (Amaryllidaceae) genus has about 14 species and 1 hybrid (15 taxa) indigenous to Turkey [10]. The images of the three species in genus *Galanthus* are presented in Figure 1.

*Galanthus woronowii* Losinsk., is one of the fourteen *Galanthus* L. species (fifteen taxa) that are indigenous to Turkey [11, 12]. It is prevalent in Caucasus, Trans-Caucasus, southern Russia, Georgia and northeastern Turkey. This species grows in low-to medium altitudes between 20 and 1500 m, however they are usually observed between 200 and 600 m. *G. woronowii* has broad green leaves and is an attractive gardening plant [13, 14]. Among the *Galanthus* species indigenous to Turkey, the bulbs of *G. elwesii* Hook. And the bulbs of *G. woronowii* are exported [15, 16]. *Galanthus woronowii* Losinsk (*Woronowii* snowdrop) and *Galanthus nivalis* L. (common snowdrop) are bulbous plant that flowers in spring and grown for its ornamental properties in gardens, which is also used in medicine. *Galanthus nivalis*, common snowdrop, is a perennial bulbous plant indigenous to Europe and southwest Asia. *Galanthus nivalis*, snowdrop or common snowdrop is the most well-known and most common among the 20 species of genus *Galanthus*. Snowdrops are among the first bulbs to bloom in the spring and they can reflect an impressive white feature in natural areas or where they were naturalized. *Galanthus woronowii*, which is indigenous to Turkey, Russia and Georgia, was named to honor Russian botanist and plant collector Georg Woronow (1874-1931).

The herbal pharmaceuticals developed from *Galanthus* L. genus plants contain several active biological compounds: Amaryllidaceae alkaloids [17-19], flavonoids, organic and hydroxycinnamic acids [20]. Alkaloids are biologically active substances that reflect the strong pharmacological activities of medicinal plants [21, 22]. *G. nivalis* and *G. elwesii* are two of the best known and most commonly grown bulbous plants. Their popularity is due to their beauty and longevity. They have numerous varieties and clones (Davis, 1999). Every year Turkey exports several bulbs collected from the nature. In early 1980s, this trade increased and millions of *G. elwesii* bulbs were exported to the Netherlands. The large volume of *Galanthus* bulb trade caused great concern, since it was unclear whether such a large bulb collection was sustainable. Thus, *Galanthus* was included in CITES Annex II in 1990. The harvest of wild *G. elwesii* bulbs is now controlled and monitored, and annual

export quotas are determined. Certain wild snowdrop species are endangered, and harvesting bulbs in the wild is illegal in several countries. However, CITES allows the limited trade of the wild bulbs of only three species in Turkey (*G. nivalis*, *G. elwesii*, and *G. woronowii*) [23].

*Galanthus elwesii* Hook. (Amaryllidaceae) is an easily recognized species with broad yellowish leaves, large flowers and thick marks on the inner parts. It has a relatively wide distribution in the eastern regions of former Yugoslavia, northern Greece, eastern Aegean Islands, southern Ukraine, Bulgaria and Turkey. This species has the widest distribution among other species in Turkey and it is indigenous to northwest, western and southern Anatolia [24, 25]. *Galanthus elwesii* (snowdrop) is a small bulbous plant distributed throughout South-Eastern European countries and Eurasia [26], and the plant is cultivated for its elegant flowers. Earlier research on *Galanthus elwesii* led to the isolation of a large variety of Amaryllidaceae alkaloids [27-29]. It was found that many of these compounds exhibited strong acetylcholinesterase inhibitory, cytotoxic and antiviral activities among others [30].



Figure 1. A.

*Galanthus woronowii*



Figure 1. B.

*Galanthus elwesii*



Figure 1. C.

*Galanthus nivalis*

Although the *Galanthus* species have been partially studied, most reported phytochemical studies demonstrated alkaloid structure diversity and no bioactivity studies were conducted. Thus, it is necessary to determine antioxidant capacity, phenolic and flavonoid content of this species.

## 2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

**2.1 Plant Material** The plant materials used in the study, *Galanthus elwesii* Ibradi samples were collected in Antalya province, *Galanthus nivalis* samples were collected in Istanbul province, and *Galanthus woronowii* samples were collected in Çaykara, Trabzon province in September 2018. The plant was collected and identified by Mustafa KARGIOĞLU. Plant samples were stored in Herbarium Material Warehouse at Afyon Kocatepe University. The plant leaves and grated bulbs were dried in an incubator at 60°C. The bulb and leaf samples were then pulverized to 80 mesh particle size for analysis.

**2.2. Chemical Material** Only analytical purity chemicals and solvents were used in all experiments. All chemicals used for chromatographic purposes were in HPLC purity. All solvents were filtered with a 0.45 µm filter (Millipore, Bedford, MA, USA). Galantamine hydrobromide, Gallic, protocatechuic, vanilic, caffeic, syringic, rosmarinic acids and catechin standard were procured from Sigma Chemical Co.

**2.3. Ultrasonic Assisted Plant Extraction** Ultrasonic assisted extraction was conducted with Bandelin Sonorex ultrasonic bath at a frequency of 50 kHz. 1 g leave and bulb samples of the dried plants were weighed and each plant sample was extracted separately with 30 ml 70% methanol for 30 minutes. After the extraction, the mixture was filtered with Whatman brand white band filter paper and stored in a +40C refrigerator until the analysis. Total phenolic, flavonoid, and phenolic acid content, galantamine analysis and antioxidant capacity were determined using the plant extracts.

**2.4. Determination of Galantamine Content with HPLC** Extract galantamine content was determined with the analysis method specified in the USP 40-NF 35 monograph. All analyses were conducted with an Agilent 1260 HPLC system equipped with a UV detector. The analytical column was an Agilent Zorbax extended C18 (5 µm, 150 mm × 5 mm) with a mobile phase that included a mixture of solvent A (acetonitrile) and B (water with 4.0 g/l Potassium dihydrogen phosphate) and employed the isocratic elution (10/90, v/v) at a flow rate of 1.2 mL/min (Table 1). The column temperature was maintained at 30°C and the detection wavelength was set to 288 nm for galantamine. The solvent was filtered through a 0.22 µm filter and degassed. The sample injection volume was 20 µL [31].

Table 1. Analytical conditions of HPLC for Galantamine analysis.

Parameters	Conditions
Column	Zorbax extended-C18 (C18, 4.6 mm × 150 mm, 5 µm)
Flow rate	1.2 mL/min
Injection volume	20 µL
UV detection	288 nm
Run time	12 min

**2.5 Total Phenolic Substance Determination** Extract phenolic content was determined with the Folin-Ciocalteu method modified by Elzaawely and Tawata [32]. 7250 µl deionized water, 500 µl extract, and 250 µl Folin-Ciocalteu reagent were added to 10 mL tubes and mixed and stored in a dark environment for 5 minutes. After adding 2000 µl (7.5%) Na<sub>2</sub>CO<sub>3</sub>, the volume was completed to 10 mL and incubated for 30 min. After incubation, the absorbance values of the samples were measured with a Shimadzu brand UV-1800 spectrophotometer at 765 nm wavelength. Gallic acid was used as the standard and the findings were expressed in mg gallic acid equivalent (GAE)/1 g dry weight (DW).

**2.6 Total Flavonoid Substance Determination** The total extract flavonoid content was determined with aluminum chloride colorimetric method [33]. 50 µl extract was transferred into 10 ml test tube and 950 µl methanol and 6400 µl deionized water were added, followed

by 300 µl 5% NaNO<sub>2</sub> solution. Then, 300 µl 10% AlCl<sub>3</sub> solution was added to the mixture and remixed. After 5 minutes of incubation, 2000 µl 1 M NaOH solution was added and the total volume of the mixture was completed to 10 mL. Mixture was incubated for 15 m. and the absorbance value was measured at 510 nm with a Shimadzu brand UV-1800 spectrophotometer. Catechin was used as the standard and the total flavonoid content was expressed as mg catechin equivalent per dry weight.

## 2.7 Antioxidant Activity Analysis

**2.7.1. 1,1-Diphenyl-2-Picryl-Hydrazyl Test** The extract antioxidant activity was determined with the 1,1-diphenyl-2-picryl-hydrazyl (DPPH) test as previously described in certain previous modifications [34]. Briefly, 200 µL of each extract (100 µL/mL) was mixed with 3.8 mL DPPH solution and incubated for 1 hour at room temperature in the dark. The absorbance of the mixture was measured at 517 nm. Ascorbic acid was used as the positive control. The ability of the sample to remove the DPPH radical is determined using the following formula:

$$\% \text{ DPPH free radical} = [(A_{\text{blank}} - A_{\text{sample}})/A_{\text{blank}}] \times 100$$

A<sub>blank</sub> is the absorbance of the control and A<sub>sample</sub> is the absorbance of the test compound.

The sample concentration that provided % 50 inhibition (IC<sub>50</sub>) was calculated by plotting the inhibition rates against the sample concentrations.

**2.7.2. Inhibition of the ABTS<sup>•+</sup> Radical Cation (TEAC) Test** The sample antioxidant activities were determined with Trolox equivalent antioxidant capacity (TEAC) method [35]. This method is based on the inhibition of ABTS radical cation (ABTS<sup>•+</sup>), produced by the oxidation of 2,2'-azinobis (3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) and potassium persulfate (K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>), with the addition of antioxidants to the medium. The antioxidant activity is determined by the reduction of the absorbance of the radical that has blue-green color at 734 nm wavelength for 6 minutes. Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid) was used as the positive control. The sample antioxidant activity figures were expressed as µmol Trolox/100 g.

**2.8. Phenolic Acid Determination with HPLC** The phenolic acids were separated with an Agilent 1260 series HPLC system equipped with C18 (4.6 mm × 150 mm, 5 µm) column. The flow rate of the mobile phase was kept at 0.5 mL/min. Mobile phase A was water containing 0.02% TFA, and phase B was methanol containing 0.02% TFA. The gradient conditions were as follows: 0-5 min, 25% B; 5-10 min, 25-30% B; 10-16 min, 30-45% B; 16-18 min, 45% B; 18-25 min, 45-80% B; 25-30 min, 80% B; 30-40 min, 80-25% B. The temperature of the column was controlled at 25°C. Injection volume was 10 µL. The detection wavelengths of DAD were set at four positions: 254, 275, 305, and 320 nm [36].

## RESULTS AND DISCUSSION

### 3.1. Determination of Galantamine Content with HPLC

The galantamine content determined in the leaves and bulbs of all three *Galanthus* species are presented in Table 2 below. The galantamine content was higher in the aerial parts (leaves) when compared to the underground parts (bulbs). Galantamine content of the samples are presented in Table 2.

Table 2. *Galanthus* spp. galantamine content

Galanthus Species	Aerial and Underground parts	Galantamine Content %
<i>Galanthus nivalis</i>	Leaf	0.075
	Bulbs	0.038
<i>Galanthus elwesii</i>	Leaf	0.082
	Bulbs	0.045
<i>Galanthus woronowii</i>	Leaf	0.078
	Bulbs	0.040

**3.2. Total Phenolic Content** The absorbance values for different concentrations of gallic acid standard solutions were measured at 765 nm with a Shimadzu brand UV-1800 spectrophotometer. The gallic acid standard curve was plotted.




Figure 2: Standard gallic acid curve

The total phenolic content in *G. nivalis*, *G. elwesii* and *G. Woronowii* leaves and bulbs are presented in Table 3 below.

Table 3: Total phenolic content analysis findings

Galanthus Species	Aerial and Underground parts	Total Phenolic Content mg GAE / gr. D.W.

<i>Galanthus nivalis</i>	Leaf	36
	Bulbs	32
<i>Galanthus elwesii</i>	Leaf	34
	Bulbs	30
<i>Galanthus woronowii</i>	Leaf	44
	Bulbs	36

A relatively high total phenolic content was determined in the leaves of all three *Galanthus* species when compared to the bulbs. The total phenolic content of the *Galanthus woronowii* species leaves and bulbs were higher when compared to the other species. The highest total phenolic content (22 mg GAE/g DW) was determined in *Galanthus woronowii* bulbs.

### 3.3. Total Flavonoid Content

The absorbances for five concentrations of catechin standard solutions were measured at 510 nm wavelength with a Shimadzu brand UV-1800 spectrophotometer. Catechin standard curve was plotted.




Figure 3: Standard catechin curve

The total flavonoid content in *G. nivalis*, *G. elwesii* and *G. Woronowii* leaves and bulbs are presented in Table 4 below.

Tablo 4: Total flavonoid analysis findings

<b>Galanthus Species</b>	<b>Aerial and Underground Parts</b>	<b>Total Flavonoid Content mg CAE / gr. D.W.</b>
<i>Galanthus nivalis</i>	Leaf	32
	Bulbs	24
<i>Galanthus elwesii</i>	Leaf	22
	Bulbs	19
<i>Galanthus woronowii</i>	Leaf	33
	Bulbs	27

A relatively higher total flavonoid content was determined in all three Galanthus species bulbs when compared to the leaves. The total flavonoid content in Galanthus woronowii leaves and bulbs were higher when compared to the other species. The highest total flavonoid content was determined in the bulbs of the Galanthus woronowii (5.3 mg GAE/g DW) species.

### 3.4. Antioxidant Activity

**3.4.1. 1,1-Diphenyl-2-Picryl-Hydrazyl Test DPPH** radical scavenging method was used to investigate the antioxidant capacity of the methanol extracts of three *Galanthus* species leaves and onions against free radicals. The DPPH radical is commonly used in the analysis of free radical scavenging activity due to its reaction facility. In this method, primarily the proton transfer reaction with the DPPH free radical due to the antioxidant leads to a decrease in absorbance at 510 nm. This process is based on monitoring the visible area with the spectrophotometer until the absorbance is constant. DPPH scavenging activity was 77% in the 500  $\mu\text{g/mL}$  *Galanthus woronowii* leaf extract concentration and 93% in the ascorbic acid control group (Figure 4). The percentage inhibition of species extracts are presented in Figure 4 below.




Figure 4: Free radical scavenging activity of methanolic extracts of *Galanthus* spp. Ascorbic acid was included as a positive control.

### 3.4.2. Inhibition of the ABTS<sup>•+</sup> Radical Cation (TEAC) Test Findings

The inhibition curve developed with different concentrations of the standard Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid) solutions is presented in Figure 5.




Figure 5. ABTS<sup>•+</sup> radical removal activity by Trolox standards

*G. nivalis*, *G. elwesii* and *G. Woronowii* species leaf and bulb antioxidant content are presented in Table 5 below. The sample antioxidant activities were expressed as  $\mu\text{mol}$  Trolox/100 g.

Table 5: Antioxidant activities

<b>Galanthus Species</b>	<b>Aerial and Underground Parts</b>	<b>Antioxidant Activity <math>\mu\text{mol}</math> Troloks/100 g D.W.</b>
<i>Galanthus nivalis</i>	Leaf	20
	Bulbs	19
<i>Galanthus elwesii</i>	Leaf	20
	Bulbs	17
<i>Galanthus woronowii</i>	Leaf	23
	Bulbs	21

**3.5. Chromatographic Analysis of Phenolic Acids** The highest phenolic acid content was observed in gallic acid content in the extracts, while the lowest phenolic acid content was observed in the acidic acid content. Following the gallic acid, significant phenolic acid protocatechic acid content were determined in extracts. Sample phenolic acid content are presented in Table 6 below.

Table 6. Sample phenolic acid content

Galanthus Species	Aerial/Underground Parts	Gallic Acid (ppm)	Protocatechic Acid (ppm)	Vanilic Acid (ppm)	Caffeic Acid (ppm)	Syringic Acid (ppm)	Coumaric Acid (ppm)	Rosmarinic Acid (ppm)	Catechin (ppm)
<i>Galanthus nivalis</i>	Leaf	17	12	2	5	7	4	12	8
	Bulbs	10	5	1	2	6	3	10	6
<i>Galanthus elwesii</i>	Leaf	15	10	3	4	5	3	14	7
	Bulbs	8	6	1	3	4	3	11	5
<i>Galanthus woronowii</i>	Leaf	18	11	3	3	6	5	14	6
	Bulbs	12	4	2	3	5	2	8	5

#### 4. CONCLUSION

Most previous phytochemical studies on *Galanthus* species have reported alkaloid structure diversity and no bioactivity studies have been conducted. The present study is the first in this area. Plants contain a significant number of phytochemical components, most of which are known to be biologically active and responsible for various pharmacological activities. Some of these secondary plant metabolites are preferred natural antioxidant sources against synthetic ones due to safety concerns. It was demonstrated that bioactive secondary metabolites reduce the risk and slow the progression of diseases such as cancer, cardiovascular and neurodegenerative diseases, etc. by scavenging free radicals through various biological mechanisms [37]. Phenols have the ability to remove the radicals since they contain hydroxyl groups. These important plant components release the hydrogen atoms from the hydroxyl groups to the radicals to form stable phenoxyl radicals. Therefore, they play an important role in antioxidant activity. Thus, it is very important to determine the

plant phenolic compound content to determine the antioxidant capacity of plant extracts [38]. Excess free radical formation leads to cellular damage and several functional disorders in humans such as atherosclerosis, myocardial infarction, cancer and neurodegenerative diseases. However, natural antioxidant compounds are beneficial in repairing cellular free radicals and managing various chronic diseases. Antioxidant tests are highly specific and sensitive to temperatures and incubation periods. Furthermore, the physicochemical properties of the sample are very important in the analysis of antioxidant properties [39]. Thus, the present study may serve as a guide for future researchers in pharmacology to conduct further studies on these plants by providing different perspectives.

## REFERENCES

- [1] Scarborough P, Allender S, Rayner M, Goldacre M. Validation of model-based estimates (synthetic estimates) of the prevalence of risk factors for coronary heart disease for wards in England. *Health Place*. 2009;(1):54-60.
- [2] Skuladottir H, Tjoenneland A, Overvad K, Stripp C, Olsen, J.H. Does high intake of fruit and vegetables improve lung cancer survival? *Lung Cancer*. 2006;(3):267-273.
- [3] Ling LT, Radhakrishnan AK, Subramaniam T, Cheng HM, Palanisamy UD. Assessment of antioxidant capacity and cytotoxicity of selected Malaysian plants. *Molecules*. 2010;(15): 2139-2151.
- [4] Vinson JA, Dabbagh YA, Serry M.M, Jang, J. Plant flavonoids, especially tea flavonols, are powerful antioxidants using an in vitro oxidation model for heart disease. *J. Agric. Food Chem*. 1995;(43):2800-2802.
- [5] Amarowicz R, Estrella I, Hernández T, Robredo S, Troszyńska A, Kosińska A, Ronald B. (Free radical-scavenging capacity, antioxidant activity, and phenolic composition of green lentil (*Lens culinaris*). *Food Chem*. 2010;(3):705-711.
- [6] Velioglu YS, Mazza G, Gao L, Oomah B.D. Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. *J. Agric. Food Chem*. 1998;(10):4113-4117.
- [7] Yoo KM, Lee CH, Lee H, Moon B, Lee CY. Relative antioxidant and cytoprotective activities of common herbs. *Food Chem*. 2008;(3):929-936.
- [8] Távare L, Carrilho D, Tyagi M, Barata D, Serra AT, Duarte CMM, Duarte RO, Feliciano R.P, Bronze MR, Chicau P, Espírito-Santo MD, Ferreira RB, Santos CN. (Antioxidant capacity of Macaronesian traditional medicinal plants. *Molecules*. 2010;(15):2576-2592.
- [9] Willis JC. *Amaryllidaceae. A dictionary of the flowering plants and ferns*. 8th ed. Cambridge: Cambridge University Press. (1988).
- [10] World Checklist of Selected Plant Families (WCSP). Royal Botanic Gardens, Kew. Available from: <http://apps.kew.org/wcsp/home.do>. [Last cited on 2018 April 25].
- [11] Davis AP, Galanthis L. *Flora of Turkey and the East Aegean Islands*. Edinburgh University Press, Edinburgh. (2000;(11):265-270.

- [12] Davis AP, Özhatay N. *Galanthus trojanus*: a new species of *Galanthus* (Amaryllidaceae) from north-western Turkey. *Bot J Linn Soc.* 2001;(137):409-412.
- [13] Davis AP. The genus *Galanthus*. In: *A Botanical Magazine Monograph*. Timber Press Inc. Oregon. 1999;165-170.
- [14] Davis AP. The genus *Galanthus*-snowdrops in the wild In: *Snowdrops a monograph of cultivated Galanthus*. Griffin Press Publishing Ltd. Cheltenham. 2006;44-47.
- [15] Özhatay N. (). Diversity of bulbous monocots in Turkey with special reference chromosome numbers. *Pure Appl Chem*; 2002;(74):547-555.
- [16] Tarım ve Köy İşleri Bakanlığı, Doğal Çiçek Soğanlarının 2010 Yılı İhracat Listesi Hakkında Tebliğ (2009/55), Resmi Gazete, Sayı: 27404.
- [17] Bokov DO, Samylina IA, Malinkin AD, Nikolov S. Application of HILIC-UV method in analysis of medicines containing Amaryllidaceae alkaloids. *Russ J Biopharm.* 2017;(9):52-58.
- [18] Sarikaya BB, Kaya GI, Onur MA, Bastida J, Somer NU. Phytochemical investigation of *Galanthus woronowii*. *Biochem Syst Ecol.* 2013;276–279.
- [19] Berkov S, Codina C, Viladomat F, Bastida J. Alkaloids from *Galanthus nivalis*. *Phytochemistry.* 2007;(68):1791-8.
- [20] Bokov DO, Malinkin AD, Samylina IA, Bessonov VV. Hydroxycinnamic and organic acids of snowdrops (*Galanthus L.*) *J Appl Pharm Sci.* 2017;(7):36-40.
- [21] Prakasia PP, Nair AS. Evaluation of in vitro antioxidant potential of the total crude alkaloid extract of *Glycosmis pentaphylla* (Retz.) Correa leaves. *Int J Pharm Pharm Sci.* 2016;(8):85-91.
- [22] Debnath B, Uddin MD J, Patari P, Das M, Maiti D, Manna K. Estimation of alkaloids and phenolics of five edible cucurbitaceous plants and their antibacterial activity. *Int J Pharm Pharm Sci.* 2015;(7):223-7.
- [23] Berkov S, Codina C, Bastida J. The Genus *Galanthus*: A Source of Bioactive Compounds. *Phytochemicals – A Global Perspective of Their Role in Nutrition and Health.* 2012;236-254.
- [24] Bishop M, Davis AP, Grimshaw JP. *Snowdrops: a monograph of cultivated Galanthus*. Cheltenham: Griffin Press. 2006;32-36.
- [25] Yüzbaşıoğlu S. Morphological variations of *Galanthus elwesii* in Turkey and difficulties on identification. *Bocconea.* 2012;(24):335-339.
- [26] Webb DA. In: *Flora Europaea*. Tutin TG, Heywood VH, Burges NA, Moore D, Valentine DH, Walters SM, Webb DA. Cambridge University Press; Cambridge, U.K. 1980(5):77.
- [27] Boit H, Horst E. Amaryllidaceae alkaloids VIII. Alkaloids from *Sprekelia formosissima*, *Galanthus elwesii*, *Zephyranthes candida* and *Crinum powellii*. *Chem. Ber.* 1955;(88):1590–1594. doi: 10.1002/cber.19550881019.

- [28] Cherkasov O. Plant sources of galanthamine. *Khim.-Farm. Zh.* 1977;(11):84–87.
- [29] Latvala A, Önür M, Gözler T, Linden A, Kivçak B, Hesse M. Alkaloids of *Galanthus elwesii*. *Phytochemistry*. 1995(39):1229–1240. doi: 10.1016/0031-9422(95)00187-C.
- [30] Bastida J, Lavilla R, Viladomat F. In: *The Alkaloids*. Cordell G. Elsevier Inc. Amsterdam, The Netherlands. 2006;(63):87–179.
- [31] The Galantamine Extended-Release Capsules Revision Bulletin supersedes the currently official monograph. The Revision Bulletin will be incorporated in USP 40–NF 35. (2016).
- [32] Elzaawely AA, Tawata S. Antioxidant activity of phenolic rich fraction obtained from *Convolvulus arvensis* L. leaves grown in Egypt. *J. Crop. Sci.* 2012;(4):32–40.
- [33] Chang C, Yang M, Wen H, Chern J. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J. Food Drug Anal.* 2002;(10):178-18.
- [34] Villano D, Fernandez-Pachon MS, Moya ML, Troncoso AM, Garcia-Parrilla MC. Radical scavenging ability of polyphenolic compounds towards DPPH free radical. *Talanta*, 2007;(71):230.
- [35] Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine*. 1999;(26):1231-1237.
- [36] Dawei W, Chen Chen L, Hao D, Yiping L, Huwei L. A universal HPLC method for the determination of phenolic acids in compound herbal medicines. *J. Agric. Food Chem.* 2005;(53):6624–6629
- [37] Al-Owaisi M, Al-Hadiwi N, Khan SA. GC/MS analysis, determination of total phenolics, flavonoid content and free radical scavenging activities of various crude extracts of *Moringa peregrina* (Forssk.) Fiori leaves. *Asian Pac J Trop Med.* 2014;4(12):964-970.
- [38] Sengul M, Yildiz H, Gungor N, Cetin B, Eser Z, Ercisli S. Total phenolic content, antioxidant and antimicrobial activities of some medicinal plants. *Pak J Pharm. Sci.* 2009;22 (1):102-106.
- [39] Kuppusamy P, Yusoff MM, Parine NR, Govindan N. Evaluation of invitro antioxidant and antibacteria properties of *Commelina nudiflora* L. extracts prepared by different polar solvents. *Saudi J Biol Sci.* 2015;22(3):293-30.