

Original Research Article

Plasmid Profile and Antibiotic Resistance Pattern of Bacteria from Abattoirs in Port Harcourt city, Nigeria

ABSTRACT

Several activities occur in abattoirs including receiving, slaughtering and sale of cattle for meat protein. In rearing of these animals, antibiotics and vaccines are incorporated as part of their meals as well as in the treatment of their infections. The regular use of antibiotics leading to the increased occurrence of antibiotic resistant organisms worldwide and also from abattoir samples, has prompted the determination of plasmid profile in these microorganisms as the plasmids act as a faster means of transmission of resistance genes. Several samples including swabs from the tables, cow blood, faecal matter and service water were collected from the Iwofe, Rumuodumaya and Trans-Amadi abattoirs located within Port Harcourt City. The plasmid profile of the multiple antibiotic resistant microorganisms was determined using standard microbiological procedures. From the results, Gram-positive isolates of the genera *Bacillus* and *Staphylococcus* exhibited 100% resistance to Cefuroxime, Ceftazidime and Augmentin while they exhibited 100% sensitivity to Ofloxacin. The Gram negative isolates including *E. coli*, *Pseudomonas* spp. and *Proteus* spp. exhibited 100% resistance to cefuroxime alone as well as 100% sensitivity to Ofloxacin. *Escherichia coli* strain 2017C-4109, *Bacillus amyloliquefaciens* strain WU-12, *Bacillus flexus* isolate Murraya and *Klebsiella pneumoniae* strain K20 lacked plasmids while *Pseudomonas* sp. strain 6174, *Escherichia coli* strain SAMA_EC, *Staphylococcus aureus* strain NCIM2654, *Klebsiella* sp. strain EIKU11 and *Proteus mirabilis* strain 46X4 all had at least one plasmid. The absence of plasmids in some of the isolates may indicate that the resistance exhibited by them may be chromosome-mediated and not plasmid-mediated. The occurrence of plasmids in multi-resistant microorganisms, poses a serious public health threat as other susceptible organisms may become resistant to the regularly used antibiotics over time.

Keywords: Plasmid profile, bacteria, antibiotic resistance, abattoir.

1. INTRODUCTION

The increased demand for cow meat and other by-products from cow has led to the establishment of abattoirs in almost every locality within Port Harcourt. As an advantage, this has not only provided ease of meat purchase but also reduced cost of the meat. Also, it serves as source of employment for the increasing population [1]. However, with the increase in the establishment of abattoirs, there is proportional increase in the generation of wastes resulting from the rearing and slaughter of the animals. In most cases, these wastes are channeled through tributaries into major streams or rivers such as that of the Trans-Amadi abattoir in Port Harcourt [2]. These wastes which are either in solid or liquid form often contain microorganisms which may be pathogenic and in some cases resistant to regularly prescribed antibiotics. These microorganisms include rotaviruses, *Cryptosporidium parvum*, *Giardia lamblia*, hepatitis E virus, *Yersinia enterocolitica*, *Campylobacter* spp., *Vibrio* spp., *Salmonella* spp. and *E.coli* 0157:H7 [3]; [2]. Conversely, the slaughter men, meat sellers and buyers are exposed to these microorganisms. In some instances, the water used in washing the slaughtered animal may not be portable and could as well be a source of microbial contamination of the meat and possibly the handlers. The faecal material from the animals, which often contain pathogenic organisms, may act as a vehicle for the transmission of zoonotic infections [4].

Zoonotic infections can be spread through hand contact and have over time been a public health concern [5]. The reduction of these diseases in animal farms has been effective using antibiotics; however, awareness on the use of antibiotics has been on the increase

[6]. In the US, it was reported that about 40% of the produced antibiotics were added to livestock feeds [7]. According to studies, penicillins and cephalosporins which are β -lactam inhibitors are among the antibiotics regularly prescribed, worldwide [8]. This scenario has led to increased resistance of bacteria to these antibiotics. This rapid growth in the occurrence of antibiotic resistance has been made possible through different means including the acquisition of plasmids. These plasmids are extra-chromosomal materials which are able to effect the production of β -lactamases. Some scholars have suggested that these β -lactamases have not only been active against β -lactam drugs but also on aminoglycosides and quinolones which are non- β -lactam drugs; this have more destroying effect on patients who are immunocompromised and makes treatment of illnesses difficult [9].

Plasmids are often implicated in increasing drug resistance as they are able to transfer the genes both within species and between different species [10]; [11]. Mechanisms of antibiotic resistance include structural modification of the target, degradation of the drug by enzymes and efflux of antibiotics [12]. Conversely, the genes responsible for resistance are either located on the chromosome or on the plasmid. This provides a medium for the quick spread of resistance genes than mutation and vertical evolution [13]. Plasmid profiling has proved to be relevant in the epidemiologic study of drug resistance as this explains the pattern, occurrence and likely future picture of the resistance when linked with some parameters [14]. It also aids in surveillance in relating strains with outbreaks and their spread [15]. The occurrence of Multiple Drug Resistance in some microorganisms isolated from abattoirs has also been reported [16]. This study was therefore aimed at determining the antibiotic susceptibility pattern and plasmid profile of isolated microorganisms from some abattoirs in Port Harcourt as this provides information on the potential health and environmental effects of the presence of the microorganisms.

Materials and Method

Sampling Locations

The locations where the samples were collected included Iwofe, Rumuodumaya and Trans-Amadi abattoirs. All three abattoirs are located within Port Harcourt City in Rivers State, Nigeria. The Trans-Amadi and Rumuodumaya abattoirs are however usually busy as it is located within popular markets. Trans-Amadi abattoir is located at longitude 04 48.442 N and latitude 007 2.303E; Rumuodumaya abattoir at longitude 04 '52' 48.0 N and latitude 7'58'20.0 E while Iwofe abattoir is located at latitude 4 59'14.0N and longitude 7 16' 12.0 E.

Table a: Sampling points, GPS coordinates and sample types

Sampling stations	Sampling points	Sampling coordinates		Type of samples
		Northing	Easting	
Iwofe	i	004° 48.598'	006° 58.517'	Blood
	ii	004° 48.592'	006° 58.501'	Swab
	iii	004° 48.601'	006° 58. 525'	Water
	iv	004° 48.594'	006° 58.518'	Faeces
Rumuodumaya	i	004° 52.118'	006° 59.580'	Blood
	ii	004° 52.102'	006° 59.571'	Swab
	iii	004° 52.124'	006° 59. 602'	Water
	iv	004° 52.120'	006° 59.582'	Faeces
Trans-Amadi	i	004° 48.442'	007° 02.303'	Blood
	ii	004° 48.434'	007° 02.293'	Swab
	iii	004° 48.456'	007° 02.319'	Water
	iv	004° 48.444'	007° 02.301'	Faeces




Fig 1: Map of Port Harcourt Metropolis showing the sampling points
 (Source: Rivers State Ministry of Lands and Survey, Port Harcourt)

Sample Collection

Sample Collection

Samples used in this study were cow blood, table swab, service water and cow faeces from the abattoirs. All samples were collected under aseptic conditions using sterile syringes and swab sticks while service water was collected by allowing the tap flow for 1 minute into sterile containers and faecal samples were collected by scooping surface faeces of the cow using sterile sample containers. All collected samples were put in ice packed coolers and transported to the Microbiology laboratory immediately for analyses.

Isolation of Microorganisms

Microbiological analyses were done according to the method by [17]. The swabs were dipped in 5 ml sterilized and cooled bacteriological peptone and allowed to stand for 30 minutes to resuscitate the microorganisms; after which 1 ml was transferred to 9 ml normal for serial dilution. One milligram of faecal material from the slaughtered animal was added to sterile 9 ml peptone water from where serial dilution was done. One millilitre of the collected blood and water samples each were transferred to 9 ml of normal saline (diluent) and 10-fold serial dilution was carried out. Aliquots (0.1 ml) of the samples were plated on sterilized and cooled Nutrient agar plates, spread using a sterile glass bent rod and incubated at 37 °C for 24 - 48 hours. Biochemical and molecular characterization of the isolates were carried out to identify the isolates.

Antibiotic susceptibility testing

Antibiotic sensitivity testing was carried out using standard Gram Negative and Gram Positive discs (Abtek Biologicals, Liverpool, UK). Each ring was embedded with 8 antibiotics. Antibiotics used include Gentamicin (10µg), Ceftriaxone (30 µg), Erythromycin (5 µg), Cloxacillin (5 µg), Ofloxacin (5 µg), Augmentin (30 µg), Ceftazidime (30 µg), Cefuroxime (30 µg), Nitrofurantoin (300 µg), Cefixime (5 µg) and Ciprofloxacin (5 µg). Mueller-Hinton (MH) agar was prepared according to manufacturers instructions, allowed to cool and solidify. Cultures of the organisms were prepared by diluting the isolate in 5 ml of sterile distilled water to 0.5 Macfarland's standard. Sterile swab stick was used to pick the culture and streaked on MH agar medium ensuring even distribution of the culture, allowed to stay at room temperature for 30 minutes and then the antibiotic ring containing the discs were aseptically placed on the plate using sterilized forceps. The plates were incubated for 18 - 24 hours at 37 °C. The zones of inhibition of the antibiotic visible as a clear halo around the discs were measured and recorded in millimeters. The percentage sensitivity of the isolates was determined per antibiotic using [18].

Extraction of Plasmid DNA

Isolates that exhibited Multiple Drug Resistance (MDR) to at least three antibiotics were used for plasmid analyses. The plasmids were extracted using the Zyppy™ Plasmid Miniprep Kit (Zymo Research, Irvine, California). Pure isolates were inoculated on Luria-Bertani broth and incubated at 37 °C overnight. 60µl of the bacterial culture from LB was transferred to a 1.5 ml microcentrifuge tube; 100 µl of 7X lysis buffer was added to the above tube and mixed by inverting the tube 6 times. The setup was incubated for 1-2 minutes. Complete cell lysis was evident by the change of colour from opaque to blue; 350 µl of cold neutralization buffer was added and mixed thoroughly. Complete neutralization was evident by the change of colour to yellow; The mixture was centrifuged for 2-4 minutes at 14000 rpm; The supernatant was transferred to a Zymo-Spin™ IIN column placed in a Collection Tube; and spun for 15 seconds at 14000 rpm; The flow-through was discarded and column returned to the Collection tube; 200 µl of Endo-Wash buffer was added to the column and spun for 30 seconds at 14000 rpm; 400 µl of Zyppy™ Wash buffer was added to the column and spun for 1 minute at 14000 rpm; The column was transferred to a clean sterile 1.5 ml microcentrifuge tube and then 35 µl of Zyppy™ Elution buffer was added directly to the column matrix and incubated for 1 minute at room temperature; The setup was spun for 30 seconds at 14000 rpm to extract the plasmid DNA. The quantity of the extracted plasmid DNA was determined using a Nanodrop 1000 spectrophotometer and after which 1% agarose gel electrophoresis was used to verify the integrity of the extracted plasmids.

Gel Electrophoresis

Plasmids were separated by electrophoresis in 1% agarose (Sigma Aldrich, USA) at a voltage of 4.5 V/cm; buffer used was TAE (Tris-Acetate-EDTA) for 3 hours. Following electrophoresis, the gels were stained for 15 minutes with ethidium bromide solution (1.0 µg/ml EtBr in 0.5 x Tris-Acetate-EDTA (TAE)), and then observed under UV light. The image was registered and analyzed using Quantity One software, version 4.1 [19].

Results

Biochemical and Genomic characterization of the isolates revealed that 10 distinct microorganisms including *Escherichia coli* strain 2017C-4109, *Pseudomonas* sp. strain 6174, *Bacillus amyloliquefaciens* strain WU-12, *Klebsiella pneumoniae* strain K20, *Bacillus flexus* isolate Murraya, *Escherichia coli* strain SAMA_EC, *Staphylococcus aureus* strain NCIM2654,

Klebsiella sp. strain EIKU11 and *Proteus mirabilis* strain 46X4 were isolated from the samples (blood, swab, water and faecal matter). Table 1 shows the weight of the isolated plasmid DNA of each of the organisms in ng/μl. Figure 2 shows the sensitivity test for *Escherichia coli*. The clear zones show the antibiotics which were active against the isolates. Figure 3 shows that the gram-positive isolates (*Bacillus* spp. and *Staphylococcus* spp.) exhibited 100% resistance to Cefuroxime (30 μg), Ceftazidime (30 μg) and Augmentin (30 μg) while they exhibited 100% susceptibility to Ofloxacin (5 μg). Cloxacillin (5 μg) and Gentamicin (10 μg) were however effective against the test isolates with 78% and 52% respectively. The Gram-negative isolates (*E. coli*, *Pseudomonas* spp. and *Proteus* spp.) were resistant to Cefuroxime with 100%, Augmentin with 97.1 while they were susceptible to Ofloxacin with 100% (Fig. 4). The gram-negative bacteria were susceptible to Nitrofurantoin and Ceftazidime with 25.2% each, while they were susceptible to Ciprofloxacin with 69.8%. Figure 5 shows the plasmid profile of the 10 isolates with the lines showing the isolates with plasmids. The figure shows that isolates 2 (*Pseudomonas* sp. strain 6174), 7 (*Escherichia coli* strain SAMA_EC), 8 (*Staphylococcus aureus* strain NCIM2654), 9 (*Klebsiella* sp. strain EIKU11) and 10 (*Proteus mirabilis* strain 46X4) all had plasmids. Whereas isolates *Klebsiella* sp. strain EIKU11 and *Proteus mirabilis* strain 46X4 had multiple plasmids, isolate *Pseudomonas* sp. strain 6174 had only 1 plasmid.



Fig 2: Multiple antibiotic sensitivity test for *Escherichia coli*

Key: GEN- Gentamicin (10μg), OFL- Ofloxacin (5 μg), AUG- Augmentin (30 μg), CAZ- Ceftazidime (30 μg), CRX-Cefuroxime (30 μg), NIT- Nitrofurantoin (300 μg), CXM- Cefixime (5 μg) and CPR-Ciprofloxacin (5 μg)




Fig 3: Percentage Sensitivity of Gram-positive isolates to tested antibiotics

Key: S - sensitive, I - Intermediate, R- Resistant, GEN- Gentamicin (10µg), OFL- Ofloxacin (5 µg), AUG- Augmentin (30 µg), CAZ- Ceftazidime (30 µg), CRX-Cefuroxime (30 µg), NIT- Nitrofurantoin (300 µg), CXM- Cefixime (5 µg) and CPR-Ciprofloxacin (5 µg)




Fig 4: Percentage Sensitivity of Gram-Negative isolates to tested antibiotics

Key: S - Sensitive, I - Intermediate, R- Resistant, GEN- Gentamicin (10µg), OFL- Ofloxacin (5 µg), AUG- Augmentin (30 µg), CAZ- Ceftazidime (30 µg), CRX-Cefuroxime (30 µg), NIT- Nitrofurantoin (300 µg), CXM- Cefixime (5 µg) and CPR-Ciprofloxacin (5 µg)

Table 1: Nanodrop Quantification of Plasmid DNA

S/N	Microorganism	Quantity (ng/μl)
1	<i>Escherichia coli</i> strain 2017C-4109	25.64
2	<i>Pseudomonas</i> sp. strain 6174	27.24
3	<i>Bacillus amyloliquefaciens</i> strain WU-12	11.70
4	<i>Klebsiella pneumoniae</i> strain K20	6.41
5	<i>Klebsiella pneumoniae</i> strain K20	14.25
6	<i>Bacillus flexus</i> isolate <i>Murraya</i>	14.04
7	<i>Escherichia coli</i> strain SAMA_EC	11.19
8	<i>Staphylococcus aureus</i> strain NCIM2654	18.27
9	<i>Klebsiella</i> sp. strain EIKU11	15.44
10	<i>Proteus mirabilis</i> strain 46X4	17.14

Table 2: Resistance pattern of the selected isolates

Bacteria	Antibiotics
<i>Escherichia coli</i> strain 2017C-4109	CAZ
<i>Klebsiella pneumoniae</i> strain K20	CAZ
<i>Pseudomonas</i> sp. strain 6174	CAZ
<i>Escherichia coli</i> strain SAMA_EC	CAZ
<i>Klebsiella</i> sp. strain EIKU11	CAZ
<i>Proteus mirabilis</i> strain 46X4	CAZ
<i>Klebsiella pneumoniae</i> strain K20	CAZ
<i>Bacillus amyloliquefaciens</i> strain WU-12	CAZ, CRX, AUG
<i>Staphylococcus aureus</i> strain NCIM2654	CAZ, CRX, AUG
<i>Bacillus flexus</i> isolate <i>Murraya</i>	CAZ, CRX, AUG

Key: CAZ- Ceftazidime (30 μg), CRX-Cefuroxime (30 μg), AUG- Augmentin (30 μg)




Fig 5: Agarose gel electrophoresis showing plasmids profile of the bacterial isolates. M represents the 1kb ladder

Discussion

Majority of the isolates such as *E. coli*, *Pseudomonas* spp. and *Proteus* spp. were coliforms and according to [20], these organisms are commonly encountered in aquatic environments and soil; these are possible sources in the meat products or abattoir environments. Plasmids have been known to be extra-chromosomal elements that can replicate on their own. They are distinct from chromosomal DNA in that they can exist independent of the host [21]. In most cases, the plasmids are not required for the day to day survival of the bacteria; but confer additional features for survival especially in harsh conditions such as in the case of antibiotic resistance. The link between Multiple Drug Resistance (MDR) and plasmid profile of bacteria gives a hint on the important role played by plasmids in the spread of MDR among bacterial species [22]. Over the years, plasmid profile has been used to determine the presence, size, type and number of plasmids in a bacteria [23]. Multidrug-resistance mediated by plasmids has also, been a constraint in the treatment of infectious diseases [24].

The antibiotics sensitivity pattern of the isolates showed that all the Gram negative isolates were resistant to Cefuroxime while all the Gram positive isolates exhibited high level of resistance to Augmentin, Cefuroxime and Ceftazidime. The high rate of resistance of the isolates to Cefuroxime, Ceftazidime and Augmentin is of concern as most physicians are quick to prescribe these drugs especially Augmentin in the treatment of bacterial infections. Resistance of bacteria to Cefuroxime has been reported by [25] who investigated the susceptibility of some microorganisms to antibiotics such as cefuroxime. Augmentin resistance by microorganisms of the genera *Klebsiella*, *Escherichia*, *Proteus* and *Pseudomonas* from abattoir sources has been reported by [26]. This resistance may be caused by mechanisms such as the synthesis of low-affinity β -lactams binding proteins and the production of penicillinase [27]. Although gentamicin resistance of 63.3% and 52% resistance has been reported among the Gram negative and positive isolates respectively from abattoirs [28]. The occurrence of gentamicin resistance has been linked to transferable genetic elements which included plasmids [29]. The high percentage susceptibility of the test isolates to Ofloxacin is in agreement with reports by other authors on multiple-drug resistance bacteria [30]. In this study, it was observed that drugs with higher MIC values such as Augmentin were not as effective against the isolates as those with lower MIC such as Ofloxacin (5 μ g). This may be due to the frequency of usage of the drug. Multiple antibiotic resistances among bacterial isolates from abattoir sources have been attributed to the increased use of antibiotics in cattle farms in Nigeria for prophylaxis and treatment infections in cows. Antibiotics are also used in producing animal food [31]. Excretion of quantities of administered drugs from the animals to the environment either as metabolized or non-metabolized compounds undergo transformation in the environment into their active forms [32]. Also, the inappropriate use of antimicrobials by individuals has promoted the presence of strains with resistance plasmids (Nsofor and Iroegbu, 2013a). According to Nsofor and Iroegbu (2013b), resistant *E. coli* strains can be transmitted to humans from animals through food and can act as an in vivo source of transmission of resistance plasmids to strains that are drug sensitive in the intestine of the animal through conjugation.

The plasmid profile of 10 multi-resistant isolates were determined. Gel electrophoresis of the plasmid DNA showed that all the isolates except *Escherichia coli* strain 2017C-4109, *Bacillus amyloliquefaciens* strain WU-12, *Klebsiella pneumoniae* strain K20, *Klebsiella pneumoniae* strain K20 and *Bacillus flexus* isolate *Murraya* had at least one plasmid. The two strains of *Bacillus* isolated lacked plasmid DNA while one strain of *E. coli* (*Escherichia coli* strain 2017C-4109) lacked plasmid DNA but the other (*Escherichia coli* strain SAMA_EC) possessed plasmid. The presence of plasmid in *Pseudomonas* sp. especially *Pseudomonas aeruginosa* has been reported by [33] who studied the plasmid profile of *P. aeruginosa* isolated from wound infections in South West, Nigeria. The absence of plasmids in *Escherichia coli* strain 2017C-4109, *Bacillus amyloliquefaciens* strain WU-12, *Klebsiella pneumoniae* strain K20 and *Bacillus flexus* isolate *Murraya* implies that the resistance may not be plasmid-mediated and may invariably be chromosome-mediated and as such may be caused by mechanisms other than plasmid-mediation [34]; [35]. This study recorded the presence of *Klebsiella pneumoniae* strain K20 with a weight of 6.41 kbp. [36] reported that strains containing larger plasmids were mainly from non-human sources while the smaller ones of about 4.7 to 10.8 kbp were mostly from humans. Also, this difference in plasmid sizes may be attributed to the presence of different resistance plasmids (R plasmids) responsible for multiple antibiotic resistances [37]. A study carried out by Nsofor and Iroegbu (2013a), also reported the presence of plasmids in strains of *E. coli* isolated from cattle.

Conclusion

The location of two of the abattoirs within markets not only exposes the meat handlers to these pathogenic resistant organisms but also other traders who may come in contact with the meat handlers or the runoffs from the slaughterhouse. The findings of this study points to the presence of plasmid-containing strains which serve as reservoirs for resistance plasmids that may be transferred to otherwise susceptible bacteria making them resistant, thus increasing the occurrence of antibiotic resistance among these pathogenic organisms. This causes difficulty in the treatment of infections caused by them. It is strongly recommended that the frequent and indiscriminate use of antibiotics be discouraged by both humans and animals. The service water from the abattoirs be treated regularly to reduce the transfer of resistant microorganisms from the water source to the animals as they consume these abattoir waters too. The abattoir wastes should be properly disposed to reduce the persistence of these resistance microorganisms in the environment.

REFERENCES

1. Nafarnda et al, I.E., ShawuNafarnda, W.D., Ajaylu, J.C., Kawe, M.S., Omeiza, G.K., Sani, N.A., Tenuche, O.Z., Dantong, D.D. and Tags, S.Z. (2). Bacteriological quality of abattoir effluents discharged into water bodies in Abuja, Nigeria. *Veterinary Science*. 515689.

2. Ogbonna. D.N. (2014). Distribution of Microorganisms in Water, Soils and Sediment from Abattoir Wastes in Southern Nigeria. *International Journal of Current Microbiology and Applied Sciences*. 3 (9): 1183-1200.
3. Sobsey MD, Handzel T, Venczel L (2003). Chlorination and safe storage of household drinking water in developing countries to reduce waterborne diseases. *Water Science and Technology*. 47(3): 221-228.
4. Adeyemo OK, Ayodeji IO, Aiki-raji CO (2002). The water quality and sanitary conditions in a major abattoir (Bodija) in Ibadan, Nigeria, *Afr. J. Biomedical Res.* 5: 51-55.
5. Tsakul, P.A., Ehinmidu, J.O. and Mohammed, S.A. (2017). Antibiotic Susceptibility and Plasmid Profile of *Escherichia coli* from Door Handles in Two Tertiary Institutions in Nasarawa State, Nigeria. *Journal of Advances in Microbiology*, 3(3): 1-13.
6. Van der Auwera, G.A., Krol, J.E., Suzuki, H., Foster, B., Van Houdt, R., Brown, C.J., Mergeay, M. and Top, E.M. (2009). Plasmids captured in *C. metallidurans* CH34: defining the PromA family of broad-host-range plasmids. *Antonie van Leeuwenhoek* 96:193-204.
7. NOT LISTED (Carraminana et al) 2009
8. Mbim, E.N., Mboto, C.I. and Edet, U.O. (2016). Plasmid Profile Analysis and Curing of Multidrug-resistant Bacteria Isolated from Two Hospital Environments in Calabar Metropolis, Nigeria. *Asian Journal of Medicine and Health*. 1(1): 1-11.
9. Amaya, E., Reyes, D., Paniagua, M., Calderon, S., Rashid, M.U., Colque, P., Kuhn, I., Carraminana, J.J., Rota, C., Agustin, I. and Herrera, A. (2004). High prevalence of multiple resistance to antibiotics in *Salmonella* serovars isolated from a poultry slaughterhouse in Spain. *Vet. Microbiol.* 104: 133-139.
10. Grohmann, E., Muth, G. and Espinosa, M (2003). Conjugative plasmid transfer in gram positive bacteria. *Microbiology Molecular Biology Reviews*. 67(2):277-301.
11. Kalpana, C. and Harish, K.B. (2015). Plasmid mediated methicillin and vancomycin resistant *Staphylococcus aureus* isolated from Northern India. *Journal of Agricultural and Biological Science*. 10(3):1990-6145
12. Lowy, F.D. (2003). Antimicrobial resistance: The example of *Staphylococcus aureus*. *Journal Clinical Investment*. 111: 1265-1273.
13. Akindele, PO and Afolayan CO (2017). Plasmid Profile of Multidrug Resistant Bacteria Isolated from Wound Swabs from Hospital Patients in Akure, Nigeria. *Asian Journal of Medicine and Health* 2(3): 1-13.
14. Wilberforce, M, Tucker S, Abendstern M (2013). Membership and management: structures of inter-professional working in community mental health teams for older people in England. *Int. Psychogeriatr.* 25(9): 1485-92.
15. Tatsuya, A. and Mary, C.S. (2010). Populations of antibiotic resistance coliform bacteria change rapidly in a wastewater effluent dominated stream. *Science of Total Environment*, 48: 6192-6201.
16. Odeyeme, T.O., Awokunmi, E.E., Adebayo, AA, Fasoranti, FO (2015). Plasmid profile of multi-drug resistance bacteria isolated from available water sources and leachate samples from dumpsite at Ebira communities in Ekiti North senatorial district, ekiti state, Nigeria. *European Journal of Advanced Research in Biological and Life Sciences* Vol. 3 No. 1
17. Cheesbrough, M. District laboratory Practice in Tropical Countries. Part 2. Cambridge University Press; 2006, pp 39-40.
18. Microsoft Excel (2010).
19. Molina-Aja, A., Garcia-Gasca, A., Abreu-Grobois, A., Bolan-Mejia, C., Roque, A. and Gomez-Gil, B. (2002). Plasmid profiling and antibiotic resistance of *Vibrio* strains isolated from cultured penaeid shrimp. *FEMS Microbiol. Lett.*, 213: 7-12.

20. Okonko, I.O., Adejoye, O.D. and Ogunnusi, T.A. (2008). Microbiological and physicochemical analysis of different water samples used for domestic purposes in Abeokuta and Ojota, Lagos State, Nigeria. *African Journal of Biotechnology*. 7(5): 617-621.
21. Raghada ,S.M., Munera, C.I. and Ayad, M.A.F. (2013). Detection of plasmid DNA profile in bacterial and fungal isolates from catheterized patients and its relation with antibiotic susceptibility. *Journal of Biotechnology Research Center*. 7: 51-60.
22. Christopher, A.F., Hora, S. and Ali, Z. (2013). Investigation of plasmid profile, antibiotic susceptibility pattern, and multiple antibiotic resistance index calculation of *Escherichia coli* isolates obtained from different human clinical specimens at tertiary care hospital in Bareilly-India. *Annals of Tropical Medicine and Public Health*. 6:285-9.
23. Pattnaik, S., Pradhan, M. and Narayani, M. (2013). Resistance Pattern acquired by nosocomial pathogens: A plasmid profile study. *Journal of Applied Pharmaceutical Science*. 3:057-059.
24. Parisa, K., Mehran, H. and Arezoo, T. (2011). Multimetal resistance study of bacteria highly resistant to mercury isolated from dental clinic effluent. *African Journal of Microbiological Research*, 5(7): 831-837
25. Harrison, C.J. and Bratcher, D. (2008). Cephalosporins: a review. *Pediatr. Rev.* 29(8): 264-267.
26. Adesoji, A.T., Abubakar, F. and Ipinlaye, J.S (2016). Occurrence of Antibiotic Resistant Bacteria in Faeces from Abattoir Waste, Processing Water and Products from Dutsin-Ma, Katsina State, Nigeria. *Journal of Bacteriology and Mycology*. 3(1): 1022.
27. Chambers, H.F. (2001). The changing epidemiology of *Staphylococcus aureus*? *Emerging infectious diseases* 7: 178.
28. Ogunshe, A., Niemogha, M., Azum, G. and Odikagbue, A. (2012). Microbiological evaluation of antibiotic resistance in bacterial flora from skin wounds. *Journal of Pharmaceutical and Biomedical Sciences* 22: 1-7.
29. Heuer, H., Krögerrecklenfort, E., Wellington, E., Egan, S., Van Elsas, J., Van Overbeek, L., Collard, J.M., Guillaume, G., Karagouni, A. and Nikolakopoulou, T. (2002). Gentamicin resistance genes in environmental bacteria: prevalence and transfer. *FEMS microbiology ecology*. 42: 289-302.
30. Odeyemi, AT, Ajayi, AO and Igbalajobi, O. A. (2013a). Plasmid Profile of isolated Bacteria from Arinta Waterfall in Ipole-Iloro Ekiti. *Journal of Microbiology Research* 3(1): 32-38.
31. Alliance for the prudent use of antibiotics (2010). Antibiotics in food animal production: a forty year debate. *APUA Newsletter*. 28(2):1-27.
32. Kümmerer, K. (2003). Significance of antibiotics in the environment. *J Antimicrob Chemother*. 52: 5-7.
33. Akingbade, S.A., Balogun,., Ojo, D.A, Afolabi, R.O., Motayo, BO, Okerentugba, PO and Okonko IO (2012). Plasmid Profile Analysis of Multidrug Resistant *Pseudomonas aeruginosa* Isolated from Wound Infections in South West, Nigeria. *World Applied Sciences Journal* 20 (6): 766-775, 2012
34. Normark, B.H. and Normark, S. (2002). Evolution and spread of antibiotic resistance. *Journal of Internal Medicine*. 252: 91-106.
35. Ezeokoli, O., Ezeamaramu, L.A.F., Ovioma, G., Oladele, J., Fowora, M., Ugo-Ijeh, M. and Iyile, J. (2010). Antibiotics Susceptibility Pattern and Plasmid Profile of Bacteria Isolated from Public Motorcycle Helmets Obinna . *American Journal of Microbiological Research*. 4(4): 126-131
36. Agbagwa, Icggen, B., Guerakan, G.C. and Oezcengiz, G. (2002). Characterization of local isolates of Enterobacteriaceae from Turkey. *Microbiol. Res*. 157: 233-238.
37. O.E. and Jirigwa, C.E. (2015). Antibiotics Resistance and Plasmid Profile of *Staphylococcus aureus* from Wound Swabs in Port Harcourt Nigeria. *Current Research in Bacteriology*, 8 (3): 70-76.

38. Alliance for the Prudent Use of Antibiotics (APUA) (2010). The Cost of Antibiotic Resistance to U.S Families and the Health Care System.
39. Nsofor, C.A. and Iroegbu, C.U. (2013a). Antibiotic Resistance Profile of *Escherichia coli* Isolated from Five Major Geopolitical Zones of Nigeria. *J. Bacteriol. Res.* 5(3):29-34.
40. Nsofor, C.A. and Iroegbu, C.U. (2013b). Plasmid profile of antibiotic resistant *Escherichia coli* isolated from domestic animals in South-East Nigeria. *Journal of Cell and Animal Biology.* 7(9): 109-115.
41. Nwoire, A., Madubuko, E.F., Eze, U.A., Oti-Wilberforce, R.O., Azi, S.O. and Ibiam, G.A. (2013). Incidence of *Staphylococcus aureus* in clinical specimens in Federal Teaching Hospital. Abakaliki, Ebonyi State. *Merit Research Journal of Medicine and Medical Sciences.* 1(3):043-046.
42. Weintraub, R.A. and Nord, C.E. (2012). Antibiotic resistance patterns of *Escherichia coli* isolates from different aquatic environment sources in leon, Nicaragua. *Clinical Microbiological Infections.* 18:347-54.