

Original Research Article

EVALUATIONS OF RED CABBAGE (*BRASSICA OLERACEA* L. VAR. *CAPITATA* F. *RUBRA* DC.) EXTRACT BEHAVIOR UNDER HIGH DOSE GAMMA IRRADIATION

ABSTRACT

Aqueous solutions of different pH from red cabbage (*Brassica oleracea* L. var. *capitata* f. *rubra* DC.) extracts were prepared. Effects of different doses/dose rates of gamma irradiation on the red cabbage aqueous solutions were investigated. Solutions were exposed to 1- 10 kGy of gamma irradiation doses at different rates ranged between 0.317 to 38.883 kGy/h. The absorption change due to gamma irradiation exposures was determined by a spectrophotometry. Results demonstrate that under gamma irradiation exposure variations (doses and dose rates), the best linearity ($R^2 > 0.97$) of red cabbage solution extract was at a pH=3.3. Results could be of great importance and clearly indicated the efficacy of a natural red cabbage dye as dosimetric indicators.

Keywords: Red cabbage, Absorption, Gamma Irradiation, Fitting, Dosimetry Indicator, pH.

1. Introduction

Ionizing radiation is an important technology that has a wide arrays of beneficial applications in different trailed such as agriculture (preservation of food), medicine (sterilization of medical and pharmaceuticals products), environment, and industrial improvement. Application of ionizing radiation is an essential method for food irradiation especially in developing countries and recently for food decontamination from environmental pollutants such as polycyclic aromatic hydrocarbons (PAHs) without changing in food proprieties, and moreover increase the storage period of food and make food more inoffensive [1, 2]. It is well known, that ionizing radiation can alter physical, chemical, and biological properties of materials depending on absorbed doses; these processes could be an important indicator related to many radiation dosimetry applications. Consequently, the dosimetry is considered a confidence key parameter of a quality assurance for radiation processes and irradiated product controls. Dosimetry is devoted for many applications in numerous areas including environmental protection, nuclear safety, food sterilization and aliments security, medicine and health care [3].

Nowadays, world current trend gives important ethics to natural sources particularly in food and medical applications [4], including natural coloured compounds namely 'dyes' such as anthocyanins, carotenoids, and betalains [5, 6], these dyes are not only used in food applications, but also for reducing some important diseases such as diabetes, obesity and cancer [7, 8].

Several natural dyes are used also in dosimetry measurement since natural dyes are biodegradables, health preserving and have no clearance problem for the environment. Contrary to synthetic dyes, which may be carcinogenic, and non-biodegradables, in addition, the potential side effect influences of the chemicals used in food synthesis colorants and the existence of environmental pollutants [9]. Therefore, world attention to natural

dyes as visual or dosimetric indicators increased [10]. Red cabbage contains the most sensitive pigment, which we can meet in food, called as anthocyanin (Fig. 1), this pigment changes colour dependent on aqueous solution pH of the red cabbage extract. Thus, red cabbage is used to monitor the level of hydrogen ions in the solution [11].




Fig. 1. Structure of cyanidin, type of anthocyanin in red cabbage [12]

The present study investigates the impact of dose and dose rate effects of ionizing radiation on the aqueous solution extract of a natural red cabbage. In addition, radiation effects on different ranges of pH on the dosimetric responses were also reported.

2. Materials and methods

2.1. Sample preparation

The red cabbage extract was prepared in aqueous solution with different pH. The red cabbage samples were ground into small pieces and about 15g of ground red cabbages was add to the solution and left for 24h at room temperature before filtration (B₁, B₂, N, A₁, A₂), while the two samples A_{1-H}, B_{2-H} were obtained by heating the red cabbage pieces in the solution for one hour at 60 °C. The different pH solutions were prepared using glacial acetic acid and sodium hydroxide (CARLO ERBA, France). Different pH values used in this study are listed in Table (1).

Table 1. Extracted colours description

| | B ₁ | B _{2-H} | B ₂ | N | A ₁ | A _{1-H} | A ₂ |
|-----------------|----------------|------------------|----------------|-----|----------------|------------------|----------------|
| pH | 12.1 | 12.1 | 13 | 6.9 | 3.3 | 3.3 | 3.6 |
| Color | yellow | yellow | yellow | red | red | red | red |
| λ_{max} | 325 | 325 | 318 | 528 | 531 | 531 | 530 |

2.2. pH measurement

pH values of the solutions were determined using an HI 8521pH meter (Hanna Instruments, Woonsocket, RI, USA). The pH meter was calibrated using a solution of known pH (4, 7 and 9).

2.3. Irradiation treatment

The gamma irradiation processes were performed under general conditions, (air atmosphere and ambient temperature) and by using two different ways of gamma irradiation. The first, was by means of a gamma cell

(dose rate 0.425130kGy/h, trans. dose 2.05 Gy) and the second, was in a ^{60}Co Gamma irradiator (Russian Type: ROBO) (200 kCi). Samples were exposed to 1, 2, 5, 10 kGy doses with a different dose rates (2.2470, 9.2585, 20.9714, 31.5389, 38.8829kGy/h) of gamma irradiation. The two gamma facilities were calibrated by Fricke using the standard (ISO/ASTM 51026, 2005. Standard Practice for Using the Fricke reference Standard Dosimetry System) [13].

3. Results and discussion

3.1. Absorption spectra

Absorption spectra of unirradiated and irradiated solutions of red cabbage extracts were measured using a UV-VIS spectrophotometer at wavelength range from 400-700 nm (resolution: 1 nm, reference: solution of same pH without red cabbage). It is known that, red cabbage contains dyes that can react with the gamma irradiation and produce some change in the intensity of the colour similar to artificial dyes [14-16].

The result of red cabbage in alkaline solution did not give a good correlation between absorption and gamma irradiation doses, and only red cabbage in acidic solution that has a maximum of absorption at 531 nm corresponding to the red colour of the anthocyanin that gave good correlation. This result was in agreement with result reported by Shahid et al. [11].

The result of irradiation in alkaline solution (basic conditions) shows that no changes were found and turbidity in the formed solution has been seen very clearly. As shown in Fig. 1, the anthocyanin type structure presented in red cabbage has five acidity hydrogens ($\text{pK}_a = 3.8$). The interpretation for that behaviour under basic conditions is due the formation of anions, which are stabilized by resonance and having negative charge on highly electromotive atoms. Therefore the formed salt after the addition of sodium hydroxide is very stable, whereas. the formed five RO^- groups are stabilizing the formed complex as the result of irradiation under basic conditions which is sort of acid-base equilibrium.

Fig. 2 shows the changes in the intensity of absorption peak at 531 nm due to the irradiation process. The intensity of absorption peak at 531 nm decreased gradually with the increase of the dose of Gamma radiation without changing the position of the peak of absorption. The increase of gamma irradiation doses is probably due to the red cabbage degradation dye by the effect of irradiation. Moreover, this decrease was also affected by the dose rate; the decrease was higher (same absorbed dose) at high dose rate, while the absorption was lower at low dose rate.




Fig. 2. Variation of absorption spectrums of red cabbage solution (pH=3.3) with respect to gamma radiation dose (from 1 to 10 kGy) at different dose rate: 10kGy/h (left), 38 kGy/h (right)

3.2. Evaluation the relation between the absorption at 531 nm and the irradiation dose rate

The behaviour of this solution under different irradiation dose rate, which varies between 0.43 and 38.88 kGy/h, was found to go through a linear tendency of photo- degradation (absorption decrease) [14], as in the Eq. (1)

$$A_0 - A_{d.r.} = \theta . D_r + \vartheta \quad (1)$$

Where: A_0 : Absorption of un-irradiated sample

$A_{d.r.}$: Absorption of irradiated sample at specific dose rate.

D_r : Gamma irradiation dose rate

θ, ϑ : constants

Fig. 3 demonstrates that, the same trend was also observed at different absorbed doses (1, 5, 10 kGy). Table (2) displays the value of the slope of linear fitting and the crossing point. The correlation coefficient (R^2) of the linear regression model has a value > 0.99 . The negative values of slope indicate that the absorption decreased with the increase of applied dose rates.

Table (2) displays the value of the slope of linear fitting and the crossing point. The correlation coefficient (R^2) of the linear regression model has a value > 0.99 . The negative values of slope indicate that the absorption decreased with the increase of applied dose rate.




Fig. 3. Variation of absorption of red cabbage extract in acidic solution A2 with irradiation dose rate, Dose rate 0.43, 2, 10, 20, 38 kGy/hat irradiation dose 1, 5, 10 kGy

Table 2. Values of the constant of the linear fitting

| Dose (kGy) | θ | ϑ | R^2 |
|------------|----------|-------------|-------|
| 1 | -0.0035 | 0.161 | 0.994 |
| 5 | -0.0046 | 0.437 | 0.998 |
| 10 | -0.0027 | 0.470 | 0.994 |

3.3. Evaluation the relation between absorption at 531 nm and irradiation dose at different dose rate

Sample solution exposed to different dose rates was found to be discoloured with the increase of the absorbed dose. Fig. 4 shows the relation between the absorbed dose and the variation of the absorption at 531 nm. The relation between the absorbed dose and the absorbance follows an exponential model that can be expressed in the following Eq.(2):

$$A = A_{\infty D.r.} + \gamma_0 \cdot e^{-R_0 \cdot D} \quad (2)$$

Where: D: absorbed dose,

$A_{\infty D.r.}$, γ_0 , R_0 : apparent constants

This model was found to be appropriate for all studied dose rates (Fig. 4). The correlation coefficient (R^2) shows a strong relation with a range between (94.4- 99.7).

Constant value of the model was listed in the Table (3). The value of the constant of γ_0 had a constant value for all dose rates of 0,456. While the constants of R_0 were correlated and had a linear relation by the dose rate effects. The coefficient of correlation had a value of 0.954, which shows a good relation between these constants. Moreover, Fig. 5 clearly shows that R_0 was not a real constant, but it is correlated to both the irradiation dose rates.




Fig. 4. Variation of absorption of red cabbage in acidic solution A2 with irradiation dose at different dose rate
2.2470, 9.2585, 20.9714, 31.5389, 38.8829 kGy/h

Table 3. Values of the constant of the fitting

| Dose rate (kGy/h) | $A_{\infty D,r.}$ | γ_0 | R_0 |
|-------------------|-------------------|-------------------|-------------------|
| 38.88289 | 0.082 ± 0.051 | 0.464 ± 0.048 | 0.174 ± 0.050 |
| 31.53895 | 0.120 ± 0.032 | 0.427 ± 0.036 | 0.276 ± 0.074 |
| 20.9714 | 0.102 ± 0.010 | 0.442 ± 0.015 | 0.318 ± 0.027 |
| 9.25847 | 0.071 ± 0.007 | 0.467 ± 0.016 | 0.379 ± 0.030 |
| 2.24695 | 0.063 ± 0.005 | 0.483 ± 0.024 | 0.461 ± 0.051 |
| 0.42513 | 0.053 ± 0.018 | 0.454 ± 0.059 | 0.442 ± 0.103 |
| average | | 0.456 | |




Fig. 5. Linear relation between the constant R_0 for the exponential model and irradiation dose at different dose rate

In addition, the constants $A_{\infty D,r.}$ of the model at different dose rate were also correlated via linear model as demonstrated in the Fig. 6.

The abovementioned correlations allow modifying the proposed exponential model, correlated the red cabbage absorbance with the absorbed dose, and rewrite the equation as follow in the Eq. (3) and Eq. (4):

$$A = (a_{\infty, D_r} + \beta_{\infty, D_r} \cdot D_r) + 0.456 \times e^{-(0.456 + \beta_{D_r} \cdot D_r) \cdot D} \quad (3)$$

Where:

$$a_{\infty, D_r} = 0.0567(24)$$

$$\beta_{\infty, D_r} = 0.0020(2)$$

$$\beta_{D_r} = 0.0068(7)$$

Then

$$A = (0.0567 + 0.0020 \cdot D_r) + 0.456 \times e^{-(0.456 + 0.0068 \cdot D_r) \cdot D} \quad (4)$$

The relative equation above shows that the absorbance of red cabbage solution is simultaneously subjects to the variation in absorbed dose and dose rate of gamma irradiation.




Fig. 6. Linear relation between the constants A_0 of the exponential model between absorption and irradiation dose at different dose rate

Moreover, the dose rate and the absorbed dose have a relation depending on the time of the radiation process as follow in Eq. (5):

$$D[kGy] = D_r \left[\frac{kGy}{hour} \right] * \frac{t[s]}{3600} \quad (5)$$

By consequence, the absorption of red cabbage extract measurements gives a direct indicator about the irradiation dose during a specified time as shown in the following equation Eq. (6):

$$A = (0.0567 + 0.0020 \cdot D_r) + 0.456 \times e^{-\frac{t}{3600} \cdot D_r \cdot (0.456 + 0.0068 \cdot D_r)} \quad (6)$$

3.4. Effect of irradiation on pH of the acidic solution of red cabbage

The value of pH of irradiated and un-irradiated acidic solution of red cabbage was determined at different doses and different dose rate. Table (4) shows a slight variation, which indicate that, no effect of gamma irradiation on acidic property of the solution and the acidity rest stable during the whole process of irradiation.

Table 4. Variation of pH vs. irradiation dose and irradiation dose rate

| Dose kGy | 0 | | | 1 | | | 5 | | | 10 | | | | | | |
|------------------|-----|-----|-----|------|------|------|-----|-----|------|------|------|-----|-----|------|------|------|
| Dose rate kGy/h | - | 2.2 | 9.3 | 21.0 | 31.5 | 38.9 | 2.2 | 9.3 | 21.0 | 31.5 | 38.9 | 2.2 | 9.3 | 21.0 | 31.5 | 38.9 |
| A ₁ | 3.3 | 3.3 | 3.3 | 3.2 | 3.3 | 3.4 | 3.2 | 3.3 | 3.2 | 3.2 | 3.2 | 3.1 | 3.1 | 3.1 | 3.4 | 3.3 |
| A ₂ | 3.6 | 3.6 | 3.7 | 3.6 | 3.7 | 3.7 | 3.7 | 3.6 | 3.7 | 3.6 | 3.6 | 3.7 | 3.6 | 3.7 | 3.7 | 3.7 |
| A _{1-H} | 3.3 | 3.3 | 3.3 | 3.0 | 3.3 | 3.5 | 3.2 | 3.3 | 3.2 | 3.1 | 3.1 | 3.2 | 3.5 | 3.2 | 3.3 | 3.2 |

4. Conclusions

In this study, pH of different red cabbage (*Brassica oleracea* L. var. *capitata* f. *rubra* DC.) extract aqueous solutions were prepared. Acidic solution with pH= 3.3 showed good relations between absorbance and absorbed dose or dose rate during the irradiation process by gamma rays. An equation was concluded that correlates absorption and irradiation dose in the dose range from 1 to 10 kGy and dose rates from 0.317 to 38.883 kGy/h. The equation shows a possible application of red cabbage extract as a dosimetry indicator. Further studies are required on the extract stability of red cabbage solutions with the time and temperature variations, before a firm conclusion could be drawn for red cabbage solution possible application as a natural dosimeter indicator.

References

1. Khalil A, Al-Bachir M. The deployment of γ -irradiation for reducing polycyclic aromatic hydrocarbons and microbial load in wheat kernels. *Toxicol Environ Chem.* 2015;97: 857-867.
2. Khalil A, Al-Bachir M (2017) Decontamination of polycyclic aromatic hydrocarbons in pea seeds by gamma irradiation. *J Food Measur Charact.* 2017; DOI:10.1007/s11694-017-9493-6.
3. Abboudi M, Al-Bachir M, Koudsi Y, Jouhara H. Combined Effects of Gamma Irradiation and Blanching Process on Acrylamide Content in Fried Potato Strips. *Inter J Food Prop.* 2016; 19:1447-1454.

4. Lucera A, Costa C, Conte A, Del Nobile MA. Food applications of natural antimicrobial compounds. *Front Microbiol.* 2012; 3:287.
5. Delgado Vargas, F, Jimenez, AR, Paredes Lopez, O. Natural pigments: carotenoids, anthocyanins, and betalains-characteristics, biosynthesis, processing, and stability. *Crit Rev Food SciNutr.* 2000; 40(3): 173-289.
6. Cortez R, Luna-Vital DA, Margulis D, Gonzalez de Mejia E. Natural pigments: Stabilization Methods of Anthocyanins for Food Applications. *Compr Rev Food Sci Food Saf.* 2017; 16: 180–198.
7. Li XD, Li J, Wang M, Jiang H. Copigmentation effects and thermal degradation kinetics of purple sweet potato anthocyanins with metal ions and sugars. *App Biological Chem.* 2016; 59(1):15–24.
8. Rodriguez EB, Vidallon MLP, Mendoza DJR, Reyes CT. Health-promoting bioactivities of betalains from red dragon fruit (*Hylocereuspolyrhizus* (Weber) Britton and Rose) peels as affected by carbohydrate encapsulation. *J Sci Food Agric.* 2016; 96(14):4679–89.
9. Carocho M, Barreiro MF, Morales P, Ferreira ICFR. Adding molecules to food, pros and cons: A review on synthetic and natural food additives. *Compr Rev Food Sci Food Saf.* 2014; 13:377–99.
10. Silva-Pereira M C, Teixeira J A, Pereira-Júnior V A, Stefani R. Chitosan/corn starch blend films with extract from *Brassica oleraceae* (red cabbage) as a visual indicator of fish deterioration. *LTW- Food Sci Tech.* 2015; 61:258-262.
11. Shahid MAK, Mubashir A, Bashir B, Mansoor N. Spectrophotometric analysis and gamma irradiation effects on dosimetric properties of *Brassica oleracea* dye aqueous solutions. *Int J Chem Mater Sci.* 2013; 1(8): 195-200.
12. Akram NG, Bhutto WA, Nawaz Sharif I. A study on the response of natural dye to Gamma radiation as a dosimeter. *Afr J Chem.* 2016; 3(3): 182-187.
13. AMERICAN STANDARD TESING AND METHODS, ASTM, (1994) Using the Fricke References Standard Dosimetry System, Designation. 1994; E 1026-92,
14. Gobara M, baraka A. Tartrazine solution as dosimeter for gamma radiation measurement. *Int Lett ChemPhys Astron.* 2014; 14(1):106-117
15. Akhtar S, Hussain T, Shahzad A, Ul-Islam Q, Hussien M Y, Akhtar N. Radiation induced decoloration of reactive dye in PVA for film dosimetry. *J Bas Appl Sci.* 2013; 9:416-419
16. Ebraheem s, El-Kelany M. Dosimeter film based on Ethyl violet- bromophenol blue dyed poly (vinyl alcohol). *Open J Poly Chem.* 2013; 3:1-5