

1
2
3
4
5
6
7
8
9

***Cyperus esculentus*: Moisture Absorption Behaviour, Extract Yields and Sensory Evaluation of Soaked Tubers in Nigeria**

10
11

ABSTRACT

This work reports the comparison in the moisture absorption behavior, the extract yields and sensory evaluation of brown and yellow varieties of tiger nut (*Cyperus esculentus*) tubers, a rhizome cultivated in Nigeria, when soaked in water for tiger nut beverage production. For each tuber variety, 3g of it was steeped in 30ml of distilled water for 5 days and periodically re-weighed at 24-hour intervals until the tubers had attained saturated moisture content. Furthermore, 200g of each variety was steeped in 800ml of sterile distilled water for periods of 0 hour (control), 24 hours, 48 hours and 96 hours, respectively. At the end of each time interval, the tubers were removed, ground in 800ml sterile distilled water, sieved, and the beverage liquid filtrate obtained measured as the percentage extract yield for the tuber sample. The 9-point Hedonic scale was used for sensory evaluation. The brown tubers showed a significantly higher moisture absorption behavior than the yellow tubers; the brown and yellow had the highest rate of moisture absorption ability of 52.22% and 35.56%, respectively, occurring after soaking for 24 hours. At same 24 hours of the soaking period, the resultant extracts obtained from the brown and yellow tubers were at a significant peak yields of 92% and 89.5%, respectively. Extracts from the brown tubers were preferred in taste and colour to those from the yellow variety. Water absorption potentials of tiger nut tubers during soaking process, has potential effects on the beverage extraction and quality.

12
13
14

Keywords: tiger nuts, brown variety, yellow variety, water absorption, sensory evaluation, extract yield

15

1. INTRODUCTION

16
17
18
19
20
21
22
23
24
25
26
27
28

Tiger nut is an underutilized crop of the family *Cyperaceae*, which produces rhizomes from the base and tubers that are somewhat spherical [1]. Tiger nut is commonly known as earth almond, chufa, yellow nut sedge and Zulu nuts. [2]. In Nigeria, some of its native names include 'Aya' in Hausa, 'Imumu' in Yoruba, 'Ofio' or 'akihausa' in Igbo [3]. Furthermore, two major varieties predominates namely; yellow and brown varieties which are readily available in the market especially in the Northern part of Nigeria with the yellow variety being the most popular because of its bigger size, attractive colour and fleshier body [3,4]. The tubers are widely consumed raw or unprocessed. They could be dried, mixed with groundnut or soaked in water for varied time-lengths of about 3 days [5]. They are particularly savoured for the production of dairy-like beverage extract which is non-alcoholic and consumed as soon as it is produced during leisure hours in family gatherings, local bars or in social functions like wedding/naming ceremonies. In recent times, it has increasingly become popular in many local social functions as substitute to industrially produced conventional carbonated drinks, perhaps probably due to its low price, and wide acceptance by the people.

29
30
31
32
33
34
35
36
37
38

Traditionally, the consumed beverage extract is prepared by steeping the tubers in water for different periods of time. This may range from a few hours to a full day; or for about three to four days, after which the harvested tubers could be consumed directly or sold in the public or hawked as delicacies. Little information is available on the effects of periods of steeping in water in terms of the moisture absorption characteristics of the soaked tubers and the beverage extract yield of the derived beverage. The information acquired will shore up efforts towards moving this traditional process into an industrial scale. This research aims to determine and compare the moisture absorption behavior and the extract yields from the two varieties of tiger nut tubers, locally available in the market, when soaked in water.

2. MATERIAL AND METHODS

Collection of Samples:

The yellow and brown varieties of tiger nuts tubers, locally identified as the 'Big' and 'small' varieties, were purchased directly from local markets in Biliri (Gombe State) and used for the study. These were labeled "BBF" and "VSF," respectively, brought into the laboratory, and separately sorted as described by [6], in order to remove foreign materials, broken and damaged tubers. The sorted tubers were used in every experimental stage in the course of the study

Moisture absorption characteristics of tiger nut tubers steeped at ambient temperature

The steeping procedure described by [6], for the determination of rate of moisture absorption of tubers was used but with slight modification. For each variety, a weight of 3g of tuber was steeped in 30ml of distilled water contained in a McCartney bottle with cover, at room temperature for a period of five days. The tubers were removed, superficially dried with tissue paper, and weighed using an electronic balance (Adams AQT 1500, UK), at 24-hour intervals. Similarly, the volume of liquor left at the same time interval was also measured using a measuring cylinder. The tubers and the steeping liquor were returned into the McCartney bottles. The experiment was terminated when the tubers had attained saturation moisture content. Six replicates of this procedure were conducted.

A graph of the mean of the recorded weights of the tubers, at specified soaking times, was plotted. Also, the moisture contents of the tuber samples at each time interval was calculated based on the increase in tuber mass at corresponding times.

Water absorption rate = $(W_t - W_o)/W_o \times 100$

Where, W_t = weight of soaked tiger nut tubers at various time points, and W_o = weight of original tiger nut tubers

Determination of extract yields

Determination of the extract yield from the tiger nut tubers were as described by [7]. A total of three 200g sample of each variety of tiger nut tubers which had been washed several changes of sterile distilled water, were steeped in 800ml sterile distilled water for periods of 0 hour (control), 24 hours, 48 hours, 72 hours, and 96 hours, respectively (Figure 1).




Figure 1: Determination of beverage extract yield process from soaked tiger nut tubers

At the end of each time interval mentioned, the steeping water was discarded and the tubers rinsed very well in sterile distilled water. The sieving of each blended sample in 800ml of sterile distilled water resulted in the removal of the chaff. What was left was the quantity of 'beverage' that was extracted. The quantity of the tiger nut beverage obtainable at the point of packaging into plastic bottles was recorded as the percentage extract yield for that tuber sample. The experiment was triplicated.

Sensory evaluation

Extracts from the different varieties were subjected to organoleptic assessment by a 43-member panel. Each panelist was requested to taste the samples one after the other and to indicate their degree of likeness based on the questionnaire provided. The samples were evaluated for colour,

99 taste, aroma and general acceptability. They were required to score each parameter on a 9-point
100 Hedonic scale, with 9 indicating 'Like extremely', and 1 indicating 'Dislike extremely'.

101 **Statistical Analyses**

102 All the different experimental data obtained, except those of sensory evaluation, were analyzed by
103 two-way analysis of variance (ANOVA) using the Genstat Release 7.22 DE, version 2008 (VSN,
104 International Limited). All sensory evaluation data were analyzed using the one-way ANOVA of the
105 same software. Probability values ($P \leq 0.05$) were considered as statistically significant.

106


107 **3. RESULTS AND DISCUSSION**

108

109 **Moisture absorption characteristics of tubers**

110 The graph of the weight-change in the two varieties of tubers steeped in water at ambient temperature
111 is shown in Figure 2.

112



113

114

KEY:

115

VSF-Brown variety

116

BBF- Yellow variety

117

Figure 2: Weight-change in tubers of tiger nuts soaked in water at ambient temperature

118

119


120

121 It was observed that the moisture absorption rate for the yellow and brown tubers was
122 highest within the first 24 hours during which 35.56% and 52.22% of the total moisture
123 absorption took place, respectively. These changes were significant. Additional increases of
124 15.00%, 5.83% and 1.39% occurred at 48 hours, 72 hours, and 96 hours, respectively, for
125 the yellow tubers, while 3.34% and 5.55% were the additional increases recorded for the
126 brown tubers at 48 hours and 72 hours, respectively. The moisture absorption changes were
127 also significant at 48 hours and 72 hours in the yellow and brown tubers, respectively. At 96
128 hours, the brown tubers had had a 1.11% loss in weight.

129 It is worth noting that both the brown and yellow tuber curves became horizontal at 72 hours
130 for brown tubers and 96 hours for yellow tubers when the weights of their soaked tubers
131 were about 1.58 and 1.51 times, respectively, of the initial dry weight. This was when the
132 peak weight-change had occurred, with weight change increases of 61.11% and 57.78%,
133 respectively. The values obtained are less than the 2.2 times the original weights and 120%
134 times their weights in water obtainable in soyabeans [8], after soaking in water for eight to
135 ten hours. The traditional method of preparing retailed soaked tubers for eating purpose
136 involves soaking the tubers in water at ambient temperature for about four to five days. Once
137 every day, the steeping water is decanted and replaced with a fresh one. The understanding
138 of the water absorption abilities of tiger nut tubers during soaking is of practical importance
139 because the water absorbed governs the beverage extraction and quality. Generally, the
140 brown tubers apparently showed a higher moisture absorption characteristic than the yellow.
141 Soaking, essentially renders dried tiger nut, edible with ease and ensures acceptable
142 sensory quality [9]. The amount of water absorbed in tiger nut tubers, at room temperature,
143 has been reported to be directly related to the tuber size; the bigger the tuber, the higher the
144 quantity of water absorbed [10]. Nevertheless, water absorption properties of plant materials
145 are enhanced or hindered by some chemical components such as protein, starch, lipid and
146 seed or tuber coat components [9]. The presence of one or some of these properties in the
147 brown tubers might be the reason these tubers displayed improved moisture absorption
148 property than the yellow, when obviously the same tubers were significantly smaller in size
149 than the yellow [11].

150 **Extract yield**

151 The highest percentage yields of 89.5% and 92% from the yellow and brown tubers,
152 respectively, were obtained from the 24-hour extracts (Figure 3); and these values were
153 significant.
154



155

156

157

158

159

160

KEY:

VSF-Brown variety

BBF- Yellow variety


Figure 3: Percentage yield of extracts from tiger nut tubers

161 Afterwards, there was a gradual decline in the extract yields obtained as fermentation time
 162 increased. The period of the peak extract yield corresponded with when the tubers were at
 163 their most rapid intake in moisture absorption abilities, as earlier discussed.


164 Generally, the extract yields from the brown tubers were significantly higher than those from
 165 the yellow. This observation differed from the reports of [12] that the yellow tiger nut variety
 166 yield more extracts than other varieties in the Nigerian market.

167 **Sensory evaluation of extracts**

168 The results of the 43-panel sensory evaluation of extracts from the brown and yellow tiger
 169 nut tubers are shown in Figure 4 and Figure 5, respectively.




170 **Figure 4:** Mean sensory evaluation of extracts from brown tiger nut tubers
 171



172 **Figure 5:** Mean sensory evaluation of extracts from yellow tiger nut tubers
 173
 174

175 In Figure 4, the 96-hour extract had mean scores less than 5 in all the parameters of sensory
 176 evaluation measured. The mean sensory evaluation score of less than 5 obtained for the 96-
 177 hour extract in all the parameters assessed, may be due to the prolonged natural or
 178 spontaneous fermentation and its subsequent production of unacceptable aroma, taste,
 179 colour and general acceptability. Tapped palm saps in Malaysia subjected to natural
 180 fermentation, have been reported by [13] to be highly fermentable in nature, with the results
 181 that products having fermentation lengths longer than 48 hours tend to possess
 182 unacceptable taste and aroma because of vinegar production. As regards aroma, even
 183 though the 24-hour extract had a higher mean score (7.12) than the 48-hour (6.63), the
 184 difference was not significant; however, the aroma from these extracts were significantly
 185 preferred to those from the 0-hour and 72- hour extracts. Furthermore, there was a
 186 significant preference in colour of the 0-hour extract obtained from the unsteeped brown
 187 tubers over those of other extracts from the same steeped tubers. This may be attributed to

188 Maillard reaction. [14] have explained that Maillard reaction occurs when reducing sugars
 189 and amino acids or proteins in foods interact, resulting in brown coloured products
 190 responsible for the difference in colour between fresh and processed foods. It is believed
 191 that the tuber-steeping process (a form of food processing) could have created the
 192 opportunity for the interaction of these compounds. Lastly, the 24-hour extract was
 193 significantly preferred over the other extracts, in regard to general acceptability.
 194 In Figure 5, the 0-hour and 24-hour extracts did not differ significantly in colour and taste
 195 from each other, but were highly significantly preferred to the other extracts from the same
 196 yellow tubers. Furthermore, the 24-hour extract was significantly preferred in aroma and
 197 general acceptability to the other extracts.
 198 The observation that the aroma from the 24-hour and 48-hour extracts of the brown tubers
 199 was significantly preferred over those in 0-hour, 72-hour and 96-hour extracts from the same
 200 tubers, as well as that, the 24-hour extract from the yellow tubers was significantly preferred
 201 to those in other extracts from the same tuber, may be explained. These observations
 202 maybe attributed to the development of diversities of microorganisms in the course of the
 203 natural fermentation process. [15] have proffered that natural fermentation has the
 204 advantage of developing complex ('rich') flavours and aroma from the diversities of
 205 microorganisms occurring during fermentation process.
 206 Figure 6, shows the mean sensory panel preference scores for the beverage extracts from
 207 the different varieties of tubers.



208 **Figure 6:** Mean sensory evaluation scores for beverage extracts from the different varieties
 209 of tubers
 210

211
 212 Extracts from the brown tubers were preferred in taste and colour to those from the yellow.
 213 In addition, extracts from the different tubers did not differ significantly in aroma and general
 214 acceptability. In all, the extracts from the brown tubers were preferred in taste and colour to
 215 those from the yellow variety. This observation is contrary to the report of [12] that the yellow
 216 variety is preferred to the brown and black varieties because of its inherent properties.

217 **Conclusion**

218
 219 This study has shown that when soaked in water at room temperature, the brown and yellow
 220 tubers had the highest rate of moisture absorption ability of 52.22% and 35.56%,
 221 respectively, occurring after 24 hours of the soaking process. Furthermore, at same period,
 222 the resultant beverage extracts obtained from the soaked tubers were also at a significant
 223 peak yields of 92% and 89.5%, respectively. In sensory evaluation, the 24-hour extract from
 224 each of the studied tuber variety was significantly preferred in general acceptability over
 225 other extracts within the same tuber type. This information is essential at determining
 226 beverage extraction and quality in the quest to move tiger nut beverage production from a
 227 local process into a conventional technological form.
 228
 229

230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Aliyu A. W., Leye J. B., Adedara A. O., Basiru D. I., Characterization of oil extracted from two varieties of tiger nut (*Cyperus esculentus* L.) tubers. *American Journal of Heterocyclic Chemistry*. 2017; 3 (3): 28-36,
2. Jansen, L. L. Morphology and photoperiodic responses of yellow nutsedge (*Cyperus esculentus*). *Weed Science*. 1971; 19:210-219,
3. Oladele, K.A. Osundahunsi, F.O. Adebowale, A.Y. Influence of processing techniques on the nutrients and anti-nutrients of tiger nut (*C. esculentus*), *World Journal of Dairy and Food Science*. 2009; 2: 88-93.
4. Adama K. K., Afolayan M. O., Oberafo A. A., Thomas S. Isolation and physicochemical characterization of tigernut (*Cyperus esculentus*) starch as a potential industrial biomaterial. *International Journal of Material Science and Applications*. 2014; 3 (2): 37-41.
5. Temple V., Ojobe T., Kapu M., Chemical analysis of tigernut (*Cyperus esculentus*) *Journal of the Science of Food and Agriculture* 1990;50 (2): 261-263.
6. Turhan M., Sayar S., Gunasekaran S., Application of peleg model to study water absorption in chickpea during soaking. *Journal of Food Engineering* 2002;53: Pp 153-159.
7. Geshinde, A.F., Oyawoye, O.M. and Adebisi, A., Comparative studies on the quality and quantity of soymilk from different varieties of soybean *Pakistan Journal of Nutrition* 2008;7 (1): 157-160.
8. Shurtleff, W. and Aoyagi, A., *Tofu and Soymilk Production: a Craft and Technical Manual*. Soyfoods Center, Lafayette, California, USA. 2000; 116-117.
9. Sanchez-Zapata E., Fernandez-Lopez J. Perez-Alvarez J.A.. Tiger nut (*Cyperus esculentus*) commercialization: health aspects, composition, properties and food applications. *Comprehensive Reviews in Food Science and Food Safety* 2012;11:366-376.
10. Coskuner Y, Ercan R, Karababa E, Nazlican AN. Physical and chemical properties of chufa (*Cyperus esculentus* L) tubers grown in the Cukurova region of Turkey. *J Sci Food Agri* 2002;82:625–31.
11. Elutade O.O., Oyawoye, O.M. and Agbo E.B., *Cyperus esculentus*: physical properties of tubers and microbiological changes during natural fermentation in Nigeria, *Food and Environment Safety* 2018;17 (2): 131-140.
12. Okafor, J.N.C., Mordi, J.I., Ozumba, A.U., Solomon, H.M. and Olatunji, O. Preliminary studies on the characterization of contaminants in tiger nuts (yellow variety). In: Proceedings of 27th Annual Nigerian Institute of Food Science and Technology (NIFST) conference. 13-17 October, 2003. Pp 210-211
13. Law, S.V., Abubakar, F., Mathashim, D. and Abdulhamid, A. Minireview: Popular fermented foods and beverages in SouthEast Asia *International Food Research Journal* 2011;18: 475-484.

278
279
280
281

282
283
284
285
286
287
288
289
290
291
292
293
294
295

14. Morris A., Barnett A. and Burrows O. Food spoilage, packaging and storage *Caarticlesj* 2004;37 (3): Pp. 165-168.
15. Nout M., Rombouts F., Fermentative preservation of plant foods *Journal of Applied Bacteriology Symposium Supplement* 1992;73: 136S-147S.

UNDER PEER REVIEW