

## Gaussian Generalized Tetranacci Numbers

**Abstract.** In this paper, we define Gaussian generalized Tetranacci numbers and as special cases, we investigate Gaussian Tetranacci and Gaussian Tetranacci-Lucas numbers with their properties.

**2010 Mathematics Subject Classification.** 11B37, 11B39, 11B83.

**Keywords.** Tetranacci numbers, Gaussian generalized Tetranacci numbers, Gaussian Tetranacci numbers, Gaussian Tetranacci-Lucas numbers.

### 1. Introduction and Preliminaries

In this work, we define Gaussian generalized Tetranacci numbers and give properties of Gaussian Tetranacci and Gaussian Tetranacci-Lucas numbers as special cases. First, we present some background about generalized Tetranacci numbers and Gaussian numbers before defining Gaussian generalized Tetranacci numbers.

There have been so many studies of the sequences of numbers in the literature which are defined recursively. Two of these type of sequences are the sequences of Tetranacci and Tetranacci-Lucas which are special case of generalized Tetranacci numbers. A generalized Tetranacci sequence  $\{V_n\}_{n \geq 0} = \{V_n(V_0, V_1, V_2, V_3)\}_{n \geq 0}$  is defined by the fourth-order recurrence relations

$$(1.1) \quad V_n = V_{n-1} + V_{n-2} + V_{n-3} + V_{n-4},$$

with the initial values  $V_0 = c_0, V_1 = c_1, V_2 = c_2, V_3 = c_3$  not all being zero.

This sequence has been studied by many authors and more detail can be found in the extensive literature dedicated to these sequences, see for example [12], [16], [17], [21], [28], [29].

The sequence  $\{V_n\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$V_{-n} = -V_{-(n-1)} - V_{-(n-2)} - V_{-(n-3)} + V_{-(n-4)}$$

for  $n = 1, 2, 3, \dots$ . Therefore, recurrence (1.1) holds for all integer  $n$ .

The first few generalized Tetranacci numbers with positive subscript and negative subscript are given in the following Table 1:

Table 1. A few Tetranacci numbers

$n$	0	1	2	3	4	5	...
$V_n$	$c_0$	$c_1$	$c_2$	$c_3$	$c_0 + c_1 + c_2 + c_3$	$c_0 + 2c_1 + 2c_2 + 2c_3$	...
$V_{-n}$	$c_0$	$c_3 - c_2 - c_1 - c_0$	$2c_2 - c_3$	$2c_1 - c_2$	$2c_0 - c_1$	$2c_3 - 2c_2 - 2c_1 - 3c_0$	...

We consider two special cases of  $V_n : V_n(0, 1, 1, 2) = M_n$  is the sequence of Tetranacci numbers (sequence A000078 in [22]) and  $V_n(4, 1, 3, 7) = R_n$  is the sequence of Tetranacci-Lucas numbers (A073817 in [22]). In other words, Tetranacci sequence  $\{M_n\}_{n \geq 0}$  and Tetranacci-Lucas sequence  $\{R_n\}_{n \geq 0}$  are defined by the fourth-order recurrence relations

$$(1.2) \quad M_n = M_{n-1} + M_{n-2} + M_{n-3} + M_{n-4}, \quad M_0 = 0, M_1 = 1, M_2 = 1, M_3 = 2$$

and

$$(1.3) \quad R_n = R_{n-1} + R_{n-2} + R_{n-3} + R_{n-4}, \quad R_0 = 4, R_1 = 1, R_2 = 3, R_3 = 7$$

respectively.

Next, we present the first few values of the Tetranacci and Tetranacci-Lucas numbers with positive and negative subscripts in the following Table 2:

Table 2. A few Tetranacci and Tetranacci-Lucas Numbers

$n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	...
$M_n$	0	1	1	2	4	8	15	29	56	108	208	401	773	1490	...
$M_{-n}$	0	0	0	1	-1	0	0	2	-3	1	0	4	-8	5	...
$R_n$	4	1	3	7	15	26	51	99	191	367	708	1365	2631	5071	...
$R_{-n}$	4	-1	-1	-1	7	-6	-1	-1	15	-19	4	-1	31	-53	...

It is well known that for all integers  $n$ , usual Tetranacci and Tetranacci-Lucas numbers can be expressed using Binet's formulas

$$M_n = \frac{\alpha^{n+2}}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)} + \frac{\beta^{n+2}}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)} + \frac{\gamma^{n+2}}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)} + \frac{\delta^{n+2}}{(\delta - \alpha)(\delta - \beta)(\delta - \gamma)}$$

(see for example [12] or [30])

or

$$(1.4) \quad M_n = \frac{\alpha - 1}{5\alpha - 8} \alpha^{n-1} + \frac{\beta - 1}{5\beta - 8} \beta^{n-1} + \frac{\gamma - 1}{5\gamma - 8} \gamma^{n-1} + \frac{\delta - 1}{5\delta - 8} \delta^{n-1}$$

(see for example [5])

and

$$R_n = \alpha^n + \beta^n + \gamma^n + \delta^n$$

respectively, where  $\alpha, \beta, \gamma$  and  $\delta$  are the roots of the equation  $x^4 - x^3 - x^2 - x - 1 = 0$ . Moreover,

$$\begin{aligned} \alpha &= \frac{1}{4} + \frac{1}{2}\omega + \frac{1}{2}\sqrt{\frac{11}{4} - \omega^2 + \frac{13}{4}\omega^{-1}}, \\ \beta &= \frac{1}{4} + \frac{1}{2}\omega - \frac{1}{2}\sqrt{\frac{11}{4} - \omega^2 + \frac{13}{4}\omega^{-1}}, \\ \gamma &= \frac{1}{4} - \frac{1}{2}\omega + \frac{1}{2}\sqrt{\frac{11}{4} - \omega^2 - \frac{13}{4}\omega^{-1}}, \\ \delta &= \frac{1}{4} - \frac{1}{2}\omega - \frac{1}{2}\sqrt{\frac{11}{4} - \omega^2 - \frac{13}{4}\omega^{-1}}, \end{aligned}$$

where

$$\omega = \sqrt{\frac{11}{12} + \left(\frac{-65}{54} + \sqrt{\frac{563}{108}}\right)^{1/3} + \left(\frac{-65}{54} - \sqrt{\frac{563}{108}}\right)^{1/3}}.$$

Note that we have the following identities:

$$\begin{aligned} \alpha + \beta + \gamma + \delta &= 1, \\ \alpha\beta + \alpha\gamma + \alpha\delta + \beta\gamma + \beta\delta + \gamma\delta &= -1, \\ \alpha\beta\gamma + \alpha\beta\delta + \alpha\gamma\delta + \beta\gamma\delta &= 1, \\ \alpha\beta\gamma\delta &= -1. \end{aligned}$$

We present an identity related with generalized Tetranacci numbers and Tetranacci numbers.

**THEOREM 1.1.** *For  $n \geq 0$  and  $m \geq 0$  the following identity holds:*

$$(1.5) \quad V_{m+n} = M_{m-2}V_{n+3} + (M_{m-3} + M_{m-4} + M_{m-5})V_{n+2} + (M_{m-3} + M_{m-4})V_{n+1} + M_{m-3}V_n$$

*Proof.* We prove the identity by induction on  $m$ . If  $m = 0$  then

$$V_n = M_{-2}V_{n+3} + (M_{-3} + M_{-4} + M_{-5})V_{n+2} + (M_{-3} + M_{-4})V_{n+1} + M_{-3}V_n$$

which is true because  $M_{-2} = 0, M_{-3} = 1, M_{-4} = -1, M_{-5} = 0$ . Assume that the equality holds for all  $m \leq k$ . For  $m = k + 1$ , we have

$$\begin{aligned}
 V_{(k+1)+n} &= V_{n+k} + V_{n+k-1} + V_{n+k-2} + V_{n+k-3} \\
 &= (M_{k-2}V_{n+3} + (M_{k-3} + M_{k-4} + M_{k-5})V_{n+2} + (M_{k-3} + M_{k-4})V_{n+1} + M_{k-3}V_n) \\
 &\quad + (M_{k-3}V_{n+3} + (M_{k-4} + M_{k-5} + M_{k-6})V_{n+2} + (M_{k-4} + M_{k-5})V_{n+1} + M_{k-4}V_n) \\
 &\quad + (M_{k-4}V_{n+3} + (M_{k-5} + M_{k-6} + M_{k-7})V_{n+2} + (M_{k-5} + M_{k-6})V_{n+1} + M_{k-5}V_n) \\
 &\quad + (M_{k-5}V_{n+3} + (M_{k-6} + M_{k-7} + M_{k-8})V_{n+2} + (M_{k-6} + M_{k-7})V_{n+1} + M_{k-6}V_n) \\
 &= (M_{k-2} + M_{k-3} + M_{k-4} + M_{k-5})V_{n+3} \\
 &\quad + ((M_{k-3} + M_{k-4} + M_{k-5} + M_{k-6}) + (M_{k-4} + M_{k-5} + M_{k-6} + M_{k-7}) \\
 &\quad \quad + (M_{k-5} + M_{k-6} + M_{k-7} + M_{k-8}))V_{n+2} \\
 &\quad + ((M_{k-3} + M_{k-4} + M_{k-5} + M_{k-6}) + (M_{k-4} + M_{k-5} + M_{k-6} + M_{k-7}))V_{n+1} \\
 &\quad + (M_{k-3} + M_{k-4} + M_{k-5} + M_{k-6})V_n \\
 &= M_{k-1}V_{n+3} + (M_{k-2} + M_{k-3} + M_{k-4})V_{n+2} + (M_{k-2} + M_{k-3})V_{n+1} + M_{k-2}V_n \\
 &= M_{(k+1)-2}V_{n+3} + (M_{(k+1)-3} + M_{(k+1)-4} + M_{(k+1)-5})V_{n+2} \\
 &\quad + (M_{(k+1)-3} + M_{(k+1)-4})V_{n+1} + M_{(k+1)-3}V_n.
 \end{aligned}$$

By induction on  $m$ , this proves (3.8).

The previous Theorem gives the following results as particular examples: For  $n \geq 0$  and  $m \geq 0$ , we have (taking  $V_n = M_n$ )

$$M_{m+n} = M_{m-2}M_{n+3} + (M_{m-3} + M_{m-4} + M_{m-5})M_{n+2} + (M_{m-3} + M_{m-4})M_{n+1} + M_{m-3}M_n$$

and (taking  $V_n = R_n$ )

$$R_{m+n} = M_{m-2}R_{n+3} + (M_{m-3} + M_{m-4} + M_{m-5})R_{n+2} + (M_{m-3} + M_{m-4})R_{n+1} + M_{m-3}R_n.$$

Next we present the Binet's formula of the generalized Tetranacci sequence.

LEMMA 1.2. *The Binet's formula of the generalized Tetranacci sequence  $\{V_n\}$  is given as*

$$V_n = M_{n-3}V_0 + (M_{n-3} + M_{n-4})V_1 + (M_{n-3} + M_{n-4} + M_{n-5})V_2 + M_{n-2}V_3.$$

*Proof.* Take  $n = 0$  and then replace  $n$  with  $m$  in Theorem 1.1.

For another proof of the Lemma 1.2, see [21]. This Lemma is also a special case of a work on the  $n$ th  $k$ -generalized Fibonacci number (which is also called  $k$ -step Fibonacci number) in [2, Theorem 2.2.].

COROLLARY 1.3. *The Binet's formula of the generalized Tetranacci sequence  $\{V_n\}$  is given as*

$$V_n = A\alpha^{n-6} + B\beta^{n-6} + C\gamma^{n-6} + D\delta^{n-6}$$

where

$$\begin{aligned}
 A &= \frac{\alpha - 1}{5\alpha - 8}(V_3\alpha^3 + (V_0 + V_1 + V_2)\alpha^2 + (V_1 + V_2)\alpha + V_2) \\
 B &= \frac{\beta - 1}{5\beta - 8}(V_3\beta^3 + (V_0 + V_1 + V_2)\beta^2 + (V_1 + V_2)\beta + V_2) \\
 C &= \frac{\gamma - 1}{5\gamma - 8}(V_3\gamma^3 + (V_0 + V_1 + V_2)\gamma^2 + (V_1 + V_2)\gamma + V_2) \\
 D &= \frac{\delta - 1}{5\delta - 8}(V_3\delta^3 + (V_0 + V_1 + V_2)\delta^2 + (V_1 + V_2)\delta + V_2)
 \end{aligned}$$

*Proof.* The proof follows from Lemma 1.2 and (1.4).

In fact, Corollary 1.3 is a special case of a result in [2, Remark 2.3].

Next, we give the ordinary generating function  $\sum_{n=0}^{\infty} a_n x^n$  of the sequence  $V_n$ .

LEMMA 1.4. *Suppose that  $f_{V_n}(x) = \sum_{n=0}^{\infty} a_n x^n$  is the ordinary generating function of the generalized Tetranacci sequence  $\{V_n\}_{n \geq 0}$ . Then  $f_{V_n}(x)$  is given by*

$$(1.6) \quad f_{V_n}(x) = \frac{V_0 + (V_1 - V_0)x + (V_2 - V_1 - V_0)x^2 + (V_3 - V_2 - V_1 - V_0)x^3}{1 - x - x^2 - x^3 - x^4}.$$

*Proof.* Using (1.1) and some calculation, we obtain

$$f_{V_n}(x) - x f_{V_n}(x) - x^2 f_{V_n}(x) - x^3 f_{V_n}(x) = V_0 + (V_1 - V_0)x + (V_2 - V_1 - V_0)x^2$$

which gives (1.6).

The previous Lemma gives the following results as particular examples: generating function of the Tetranacci sequence  $M_n$  is

$$f_{M_n}(x) = \sum_{n=0}^{\infty} M_n x^n = \frac{x}{1 - x - x^2 - x^3 - x^4}$$

and generating function of the Tetranacci-Lucas sequence  $R_n$  is

$$f_{R_n}(x) = \sum_{n=0}^{\infty} R_n x^n = \frac{4 - 3x - 2x^2 - x^3}{1 - x - x^2 - x^3 - x^4}.$$

In literature, there have been so many studies of the sequences of Gaussian numbers. A Gaussian integer  $z$  is a complex number whose real and imaginary parts are both integers, i.e.,  $z = a + ib$ ,  $a, b \in \mathbb{Z}$ . These numbers are denoted by  $\mathbb{Z}[i]$ . The norm of a Gaussian integer  $a + ib$ ,  $a, b \in \mathbb{Z}$  is its Euclidean norm, that is,  $N(a + ib) = \sqrt{a^2 + b^2} = \sqrt{(a + ib)(a - ib)}$ . For more information about this kind of integers, see the work of Fraleigh [6].

If we use together sequences of integers defined recursively and Gaussian type integers, we obtain a new sequences of complex numbers such as Gaussian Fibonacci, Gaussian Lucas, Gaussian Pell, Gaussian Pell-Lucas and Gaussian Jacobsthal numbers; Gaussian Padovan and Gaussian Pell-Padovan numbers; Gaussian Tribonacci numbers.

In 1963, Horadam [13] introduced the concept of complex Fibonacci number called as the Gaussian Fibonacci number. Pethe [19] defined the complex Tribonacci numbers at Gaussian integers, see also [8].

There are other several studies dedicated to these sequences of Gaussian numbers such as the works in [1], [3], [4], [8], [9], [10], [11], [13], [14], [15], [18], [23], [24], [25], [26], among others.

### 2. Gaussian Generalized Tetranacci Numbers

Gaussian generalized Tetranacci numbers  $\{GV_n\}_{n \geq 0} = \{GV_n(GV_0, GV_1, GV_2, GV_3)\}_{n \geq 0}$  are defined by

$$(2.1) \quad GV_n = GV_{n-1} + GV_{n-2} + GV_{n-3} + GV_{n-4},$$

with the initial conditions

$$GV_0 = c_0 + i(c_3 - c_2 - c_1 - c_0), GV_1 = c_1 + ic_0, GV_2 = c_2 + ic_1, GV_3 = c_3 + ic_2,$$

not all being zero. The sequences  $\{GV_n\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$GV_{-n} = -GV_{-(n-1)} - GV_{-(n-2)} - GV_{-(n-3)} + GV_{-(n-4)}$$

for  $n = 1, 2, 3, \dots$ . Therefore, recurrence (2.1) hold for all integer  $n$ . Note that for  $n \geq 0$

$$(2.2) \quad GV_n = V_n + iV_{n-1}.$$

and

$$GV_{-n} = V_{-n} + iV_{-n-1}$$

The first few generalized Gaussian Tetranacci numbers with positive subscript and negative subscript are given in the following Table 3:

Table 3. A few generalized Gaussian Tetranacci numbers

$n$	$GV_n$	$GV_{-n}$
0	$c_0 + i(c_3 - c_2 - c_1 - c_0)$	$c_0 + i(c_3 - c_2 - c_1 - c_0)$
1	$c_1 + ic_0$	$(c_3 - c_2 - c_1 - c_0) + i(2c_2 - c_3)$
2	$c_2 + ic_1$	$2c_2 - c_3 + i(2c_1 - c_2)$
3	$c_3 + ic_2$	$2c_1 - c_2 + i(2c_0 - c_1)$
4	$c_0 + c_1 + c_2 + c_3 + ic_3$	$2c_0 - c_1 + i(2c_3 - 3c_0 - 2c_1 - 2c_2)$

We consider two special cases of  $GV_n$  :  $GV_n(0, 1, 1 + i, 2 + i) = GM_n$  is the sequence of Gaussian Tetranacci numbers and  $GV_n(4 - i, 1 + 4i, 3 + i, 7 + 3i) = GR_n$  is the sequence of Gaussian Tetranacci-Lucas numbers. We formally define them as follows:

Gaussian Tetranacci numbers are defined by

$$(2.3) \quad GM_n = GM_{n-1} + GM_{n-2} + GM_{n-3} + GM_{n-4},$$

with the initial conditions

$$GM_0 = 0, GM_1 = 1, GM_2 = 1 + i, GM_3 = 2 + i$$

and Gaussian Tetranacci-Lucas numbers are defined by

$$(2.4) \quad GR_n = GR_{n-1} + GR_{n-2} + GR_{n-3} + GR_{n-4}$$

with the initial conditions

$$GR_0 = 4 - i, GR_1 = 1 + 4i, GR_2 = 3 + i, GR_3 = 7 + 3i.$$

Note that for  $n \geq 0$

$$GM_n = M_n + iM_{n-1}, GR_n = R_n + iR_{n-1}$$

and

$$GM_{-n} = M_{-n} + iM_{-n-1}, GR_{-n} = R_{-n} + iR_{-n-1}.$$

Next, we present the first few values of the Gaussian Tetranacci and Tetranacci-Lucas numbers with positive and negative subscripts in the following Table 4:

Table 4. A few Gaussian Tetranacci and Tetranacci-Lucas Numbers

$n$	0	1	2	3	4	5	6	7	8
$GM_n$	0	1	$1 + i$	$2 + i$	$4 + 2i$	$8 + 4i$	$15 + 8i$	$29 + 15i$	$56 + 29i$
$GM_{-n}$	0	0	$i$	$1 - i$	-1	0	$2i$	$2 - 3i$	$-3 + i$
$GR_n$	$4 - i$	$1 + 4i$	$3 + i$	$7 + 3i$	$15 + 7i$	$26 + 15i$	$51 + 26i$	$99 + 51i$	$191 + 99i$
$GR_{-n}$	$4 - i$	$-1 - i$	$-1 - i$	$-1 + 7i$	$7 - 6i$	$-6 - i$	$-1 - i$	$-1 + 15i$	$15 - 19i$

The following Theorem presents the generating function of Gaussian generalized Tetranacci numbers.

THEOREM 2.1. *The generating function of Gaussian generalized Tetranacci numbers is given as*

$$(2.5) \quad f_{GV_n}(x) = \frac{GV_0 + (GV_1 - GV_0)x + (GV_2 - GV_1 - GV_0)x^2 + (GV_3 - GV_2 - GV_1 - GV_0)x^3}{1 - x - x^2 - x^3 - x^4}.$$

*Proof.* Let

$$f_{GV_n}(x) = \sum_{n=0}^{\infty} GV_n x^n$$

be generating function of Gaussian generalized Tetranacci numbers. Then using the definition of generalized Gaussian Tetranacci numbers, and subtracting  $xf(x)$ ,  $x^2f(x)$ ,  $x^3f(x)$  and  $x^4f(x)$  from  $f(x)$  we obtain (note

the shift in the index  $n$  in the third line)

$$\begin{aligned}
 & (1 - x - x^2 - x^3 - x^4)f_{GV_n}(x) \\
 = & \sum_{n=0}^{\infty} GV_n x^n - x \sum_{n=0}^{\infty} GV_n x^n - x^2 \sum_{n=0}^{\infty} GV_n x^n - x^3 \sum_{n=0}^{\infty} GV_n x^n - x^4 \sum_{n=0}^{\infty} GV_n x^n \\
 = & \sum_{n=0}^{\infty} GV_n x^n - \sum_{n=0}^{\infty} GV_n x^{n+1} - \sum_{n=0}^{\infty} GV_n x^{n+2} - \sum_{n=0}^{\infty} GV_n x^{n+3} - \sum_{n=0}^{\infty} GV_n x^{n+4} \\
 = & \sum_{n=0}^{\infty} GV_n x^n - \sum_{n=1}^{\infty} GV_{n-1} x^n - \sum_{n=2}^{\infty} GV_{n-2} x^n - \sum_{n=3}^{\infty} GV_{n-3} x^n - \sum_{n=4}^{\infty} GV_{n-4} x^n \\
 = & (GV_0 + GV_1 x + GV_2 x^2 + GV_3 x^3) - (GV_0 x + GV_1 x^2 + GV_2 x^3) - (GV_0 x^2 + GV_1 x^3) - GV_0 x^3 \\
 & + \sum_{n=4}^{\infty} (GV_n - GV_{n-1} - GV_{n-2} - GV_{n-3} - GV_{n-4}) x^n \\
 = & GV_0 + (GV_1 - GV_0)x + (GV_2 - GV_1 - GV_0)x^2 + (GV_3 - GV_2 - GV_1 - GV_0)x^3
 \end{aligned}$$

Rearranging above equation, we get

$$f_{GV_n}(x) = \frac{GV_0 + (GV_1 - GV_0)x + (GV_2 - GV_1 - GV_0)x^2 + (GV_3 - GV_2 - GV_1 - GV_0)x^3}{1 - x - x^2 - x^3 - x^4}.$$

The previous Theorem gives the following results as particular examples: the generating function of Gaussian Tetranacci numbers is

$$(2.6) \quad f_{GM_n}(x) = \frac{x + ix^2}{1 - x - x^2 - x^3 - x^4}$$

and the generating function of Gaussian Tetranacci-Lucas numbers is

$$(2.7) \quad f_{GR_n}(x) = \frac{-(1 + i)x^3 - (2 + 2i)x^2 - (3 - 5i)x + 4 - i}{1 - x - x^2 - x^3 - x^4}.$$

The result (2.6) is already known, (see [24]).

We now present the Binet formula for the Gaussian generalized Tetranacci numbers.

**THEOREM 2.2.** *The Binet formula for the Gaussian generalized Tetranacci numbers is*

$$GV_n = (A\alpha^{n-6} + B\beta^{n-6} + C\gamma^{n-6} + D\delta^{n-6}) + i(A\alpha^{n-6} + B\beta^{n-6} + C\gamma^{n-6} + D\delta^{n-6})$$

where  $A, B, C$  and  $D$  are as in Corollary (1.3).

*Proof.* The proof follows from Corollary (1.3) and  $GV_n = V_n + iV_{n-1}$ .

The previous Theorem gives the following results as particular examples: the Binet formula for the Gaussian Tetranacci numbers is

$$GM_n = \left( \frac{\alpha^{n+2}}{(\alpha-\beta)(\alpha-\gamma)(\alpha-\delta)} + \frac{\beta^{n+2}}{(\beta-\alpha)(\beta-\gamma)(\beta-\delta)} \right) + i \left( \frac{\alpha^{n+2}}{(\alpha-\beta)(\alpha-\gamma)(\alpha-\delta)} + \frac{\beta^{n+2}}{(\beta-\alpha)(\beta-\gamma)(\beta-\delta)} \right)$$

and the Binet formula for the Gaussian Tetranacci-Lucas numbers is

$$GR_n = (\alpha^n + \beta^n + \gamma^n + \delta^n) + i(\alpha^{n-1} + \beta^{n-1} + \gamma^{n-1} + \delta^n).$$



The following Theorem present some formulas of Gaussian generalized Tetranacci numbers.

**THEOREM 2.3.** *For  $n \geq 1$  we have the following formulas:*

**(a):** *(Sum of the Gaussian generalized Tetranacci numbers)*

$$\sum_{k=1}^n GV_k = \frac{1}{3}(GV_{n+2} + 2GV_n + GV_{n-1} - GV_0 + GV_1 - GV_3)$$

**(b):**  $\sum_{k=1}^n GV_{2k+1} = \frac{1}{3}(2GV_{2n+2} + GV_{2n} - GV_{2n-1} - 2GV_0 - GV_1 - 3GV_2 + GV_3)$

**(c):**  $\sum_{k=1}^n GV_{2k} = \frac{1}{3}(2GV_{2n+1} + GV_{2n-1} - GV_{2n-2} + GV_0 - GV_1 + 3GV_2 - 2GV_3).$

*Proof.*

**(a):** Using the recurrence relation

$$GV_n = GV_{n-1} + GV_{n-2} + GV_{n-3} + GV_{n-4}$$

i.e.

$$GV_{n-4} = GV_n - GV_{n-1} - GV_{n-2} - GV_{n-3}$$

we obtain

$$GV_0 = GV_4 - GV_3 - GV_2 - GV_1$$

$$GV_1 = GV_5 - GV_4 - GV_3 - GV_2$$

$$GV_2 = GV_6 - GV_5 - GV_4 - GV_3$$

$$GV_3 = GV_7 - GV_6 - GV_5 - GV_4$$

$$GV_4 = GV_8 - GV_7 - GV_6 - GV_5$$

$\vdots$

$$GV_{n-4} = GV_n - GV_{n-1} - GV_{n-2} - GV_{n-3}$$

$$GV_{n-3} = GV_{n+1} - GV_n - GV_{n-1} - GV_{n-2}$$

$$GV_{n-2} = GV_{n+2} - GV_{n+1} - GV_n - GV_{n-1}$$

$$GV_{n-1} = GV_{n+3} - GV_{n+2} - GV_{n+1} - GV_n$$

$$GV_n = GV_{n+4} - GV_{n+3} - GV_{n+2} - GV_{n+1}.$$

If we add the equations by side by, we get

$$\begin{aligned} \sum_{k=1}^n GV_k &= \frac{1}{3}(GV_{n+4} - GV_{n+2} - 2GV_{n+1} - GV_0 + GV_1 - GV_3) \\ &= \frac{1}{3}(GV_{n+2} + 2GV_n + GV_{n-1} - GV_0 + GV_1 - GV_3). \end{aligned}$$

(b): When we use (2.1), we obtain the following equalities:

$$\begin{aligned}
 GV_k &= GV_{k-1} + GV_{k-2} + GV_{k-3} + GV_{k-4} \\
 GV_4 &= GV_3 + GV_2 + GV_1 + GV_0 \\
 GV_6 &= GV_5 + GV_4 + GV_3 + GV_2 \\
 GV_8 &= GV_7 + GV_6 + GV_5 + GV_4 \\
 GV_{10} &= GV_9 + GV_8 + GV_7 + GV_6 \\
 &\vdots \\
 GV_{2n+2} &= GV_{2n+1} + GV_{2n} + GV_{2n-1} + GV_{2n-2}.
 \end{aligned}$$

If we rearrange the above equalities, we obtain

$$\begin{aligned}
 GV_3 &= GV_4 - GV_2 - GV_1 - GV_0 \\
 GV_5 &= GV_6 - GV_4 - GV_3 - GV_2 \\
 GV_7 &= GV_8 - GV_6 - GV_5 - GV_4 \\
 GV_9 &= GV_{10} - GV_8 - GV_7 - GV_6 \\
 &\vdots \\
 GV_{2n-1} &= GV_{2n} - GV_{2n-2} - GV_{2n-3} - GV_{2n-4} \\
 GV_{2n+1} &= GV_{2n+2} - GV_{2n} - GV_{2n-1} - GV_{2n-2}.
 \end{aligned}$$

Now, if we add the above equations by side by, we get

$$\begin{aligned}
 \sum_{k=1}^n GV_{2k+1} &= GV_{2n+2} - GV_2 - \sum_{k=1}^{2n-1} GV_k - GV_0 \\
 &= GV_{2n+2} - GV_2 - \frac{1}{3}(GV_{(2n-1)+4} - GV_{(2n-1)+2} - 2GV_{(2n-1)+1} - GV_0 + GV_1 - GV_3) - GV_0 \\
 &= GV_{2n+2} - GV_2 - \frac{1}{3}(GV_{2n+3} - GV_{2n+1} - 2GV_{2n} - GV_0 + GV_1 - GV_3) - GV_0 \\
 &= -\frac{1}{3}(-3GV_{2n+2} + GV_{2n+3} - GV_{2n+1} - 2GV_{2n} + 2GV_0 + GV_1 + 3GV_2 - GV_3) \\
 &= \frac{1}{3}(3GV_{2n+2} - GV_{2n+3} + GV_{2n+1} + 2GV_{2n} - 2GV_0 - GV_1 - 3GV_2 + GV_3)
 \end{aligned}$$

and

$$\begin{aligned}
 3GV_{2n+2} - GV_{2n+3} + GV_{2n+1} + 2GV_{2n} &= 2GV_{2n+2} + (GV_{2n+2} + GV_{2n+1} + GV_{2n} - GV_{2n+3}) + GV_{2n} \\
 &= 2GV_{2n+2} + GV_{2n} - GV_{2n-1}
 \end{aligned}$$

So

$$\sum_{k=1}^n GV_{2k+1} = \frac{1}{3}(2GV_{2n+2} + GV_{2n} - GV_{2n-1} - 2GV_0 - GV_1 - 3GV_2 + GV_3).$$

(c): Since

$$\sum_{k=1}^n GV_{2k+1} + \sum_{k=1}^n GV_{2k} = \sum_{k=1}^{2n+1} GV_k - GV_1$$

we have

$$\begin{aligned} \sum_{k=1}^n GV_k &= \frac{1}{3}(GV_{n+4} - GV_{n+2} - 2GV_{n+1} - GV_0 + GV_1 - GV_3) \\ &= \frac{1}{3}(GV_{n+2} + 2GV_n + GV_{n-1} - GV_0 + GV_1 - GV_3), \\ \sum_{k=1}^n GV_{2k+1} &= \frac{1}{3}(2GV_{2n+2} + GV_{2n} - GV_{2n-1} - 2GV_0 - GV_1 - 3GV_2 + GV_3) \end{aligned}$$

$$\begin{aligned} \sum_{k=1}^n GV_{2k} &= \sum_{k=1}^{2n+1} GV_k - \sum_{k=1}^n GV_{2k+1} - GV_1 \\ &= \frac{1}{3}(GV_{(2n+1)+4} - GV_{(2n+1)+2} - 2GV_{(2n+1)+1} - GV_0 + GV_1 - GV_3) \\ &\quad - \frac{1}{3}(2GV_{2n+2} + GV_{2n} - GV_{2n-1} - 2GV_0 - GV_1 - 3GV_2 + GV_3) - GV_1 \\ &= \frac{1}{3}(GV_{2n+5} - GV_{2n+3} - 2GV_{2n+2} - GV_0 + GV_1 - GV_3) + \frac{1}{3}(-2GV_{2n+2} - GV_{2n} \\ &\quad + GV_{2n-1} + 2GV_0 + GV_1 + 3GV_2 - GV_3 - 3GV_1) \\ &= \frac{1}{3}(GV_{2n+5} - GV_{2n+3} - 2GV_{2n+2} - GV_0 + GV_1 - GV_3 - 2GV_{2n+2} - GV_{2n} + GV_{2n-1} \\ &\quad + 2GV_0 + GV_1 + 3GV_2 - GV_3 - 3GV_1) \\ &= \frac{1}{3}(GV_{2n+5} - GV_{2n+3} - 4GV_{2n+2} - GV_{2n} + GV_{2n-1} + GV_0 - GV_1 + 3GV_2 - 2GV_3) \\ &= \frac{1}{3}(2GV_{2n+1} + GV_{2n-1} - GV_{2n-2} + GV_0 - GV_1 + 3GV_2 - 2GV_3) \end{aligned}$$

This completes the proof.

As special cases of above Theorem, we have the following two Corollaries. First one present some formulas of Gaussian Tetranacci numbers.

COROLLARY 2.4. For  $n \geq 1$  we have the following formulas:

(a): (Sum of the Gaussian Tetranacci numbers)

$$\sum_{k=1}^n GM_k = \frac{1}{3}(GM_{n+2} + 2GM_n + GM_{n-1} - (1 + i))$$

(b):  $\sum_{k=1}^n GM_{2k+1} = \frac{1}{3}(2GM_{2n+2} + GM_{2n} - GM_{2n-1} - 2 - 2i)$

(c):  $\sum_{k=1}^n GM_{2k} = \frac{1}{3}(2GM_{2n+1} + GM_{2n-1} - GM_{2n-2} - 2 + i).$

Second Corollary gives some formulas of Gaussian Tetranacci-Lucas numbers.

COROLLARY 2.5. For  $n \geq 1$  we have the following formulas:

(a): (Sum of the Gaussian Tetranacci-Lucas numbers)

$$\sum_{k=1}^n GR_k = \frac{1}{3}(GR_{n+2} + 2GR_n + GR_{n-1} - 10 + 2i)$$

(b):  $\sum_{k=1}^n GR_{2k+1} = \frac{1}{3}(2GR_{2n+2} + GR_{2n} - GR_{2n-1} - 11 - 2i)$

(c):  $\sum_{k=1}^n GR_{2k} = \frac{1}{3}(2GR_{2n+1} + GR_{2n-1} - GR_{2n-2} - 2 - 8i)$ .

In fact, using the method of the proof of Theorem 2.3, we can prove the following formulas of generalized Tetranacci numbers.

THEOREM 2.6. For  $n \geq 1$  we have the following formulas:

(a): (Sum of the generalized Tetranacci numbers)

$$\sum_{k=1}^n V_k = \frac{1}{3}(V_{n+2} + 2V_n + V_{n-1} - V_0 + V_1 - V_3)$$

(b):  $\sum_{k=1}^n V_{2k+1} = \frac{1}{3}(2V_{2n+2} + V_{2n} - V_{2n-1} - 2V_0 - V_1 - 3V_2 + V_3)$

(c):  $\sum_{k=1}^n V_{2k} = \frac{1}{3}(2V_{2n+1} + V_{2n-1} - V_{2n-2} + V_0 - V_1 + 3V_2 - 2V_3)$ .

As special cases of above Theorem, we have the following two Corollaries. First one present some formulas of Tetranacci numbers.

COROLLARY 2.7. For  $n \geq 1$  we have the following formulas:

(a): (Sum of the Tetranacci numbers)

$$\sum_{k=1}^n M_k = \frac{1}{3}(M_{n+2} + 2M_n + M_{n-1} - 1)$$

(b):  $\sum_{k=1}^n M_{2k+1} = \frac{1}{3}(2M_{2n+2} + M_{2n} - M_{2n-1} - 2)$

(c):  $\sum_{k=1}^n M_{2k} = \frac{1}{3}(2M_{2n+1} + M_{2n-1} - M_{2n-2} - 2)$ .

Second Corollary gives some formulas of Tetranacci-Lucas numbers.

COROLLARY 2.8. For  $n \geq 1$  we have the following formulas:

(a): (Sum of the Tetranacci-Lucas numbers)

$$\sum_{k=1}^n R_k = \frac{1}{3}(R_{n+2} + 2R_n + R_{n-1} - 10)$$

(b):  $\sum_{k=1}^n R_{2k+1} = \frac{1}{3}(2R_{2n+2} + R_{2n} - R_{2n-1} - 11)$

(c):  $\sum_{k=1}^n R_{2k} = \frac{1}{3}(2R_{2n+1} + R_{2n-1} - R_{2n-2} - 2)$ .

Note that if the sum starts with the zero then the constant in the formula may only change, for example

$$\sum_{k=0}^n R_k = R_0 + \sum_{k=1}^n R_k = 4 + \frac{1}{3}(R_{n+2} + 2R_n + R_{n-1} - 10) = \frac{1}{3}(R_{n+2} + 2R_n + R_{n-1} + 2)$$

but

$$\sum_{k=0}^n M_k = M_0 + \sum_{k=1}^n M_k = \sum_{k=1}^n M_k = \frac{1}{3}(M_{n+2} + 2M_n + M_{n-1} - 1).$$

### 3. Some Identities Connecting Gaussian Tetranacci and Gaussian Tetranacci-Lucas Numbers

In this section, we obtain some identities of Gaussian Tetranacci numbers and Gaussian Tetranacci-Lucas numbers.

First, we can give a few basic relations between  $\{GM_n\}$  and  $\{GR_n\}$  as

$$(3.1) \quad GR_n = -GM_{n+3} + 6GM_{n+1} - GM_n$$

$$(3.2) \quad GR_n = -GM_{n+2} + 5GM_{n+1} - 2GM_n - GM_{n-1}$$

and also

$$(3.3) \quad GR_n = 4GM_{n+1} - 3GM_n - 2GM_{n-1} - GM_{n-2}.$$

Note that the last three identities hold for all integers  $n$ . For example, to show (3.1), writing

$$GR_n = aGM_{n+3} + bGM_{n+2} + cGM_{n+1} + dGM_n$$

and solving the system of equations

$$GR_0 = aGM_3 + bGM_2 + cGM_1 + dGM_0$$

$$GR_1 = aGM_4 + bGM_3 + cGM_2 + dGM_1$$

$$GR_2 = aGM_5 + bGM_4 + cGM_3 + dGM_2$$

$$GR_3 = aGM_6 + bGM_5 + cGM_4 + dGM_3$$

we find that  $a = -1, b = 0, c = 6, d = -1$ . Or using the relations  $GM_n = M_n + iM_{n-1}$ ,  $GR_n = R_n + iR_{n-1}$  and identity  $R_n = -M_{n+3} + 6M_{n+1} - M_n$  we obtain the identity (3.1). The others can be found similarly.

We will present some other identities between Gaussian Tetranacci and Gaussian Tetranacci-Lucas numbers with the help of generating functions.

The following lemma will help us to derive the generating functions of even and odd-indexed Gaussian Tetranacci and Gaussian Tetranacci-Lucas sequences.

LEMMA 3.1 ([7]). *Suppose that  $f(x) = \sum_{n=0}^{\infty} a_n x^n$  is the generating function of the sequence  $\{a_n\}_{n \geq 0}$ . Then the generating functions of the sequences  $\{a_{2n}\}_{n \geq 0}$  and  $\{a_{2n+1}\}_{n \geq 0}$  are given as*

$$f_{a_{2n}}(x) = \sum_{n=0}^{\infty} a_{2n} x^n = \frac{f(\sqrt{x}) + f(-\sqrt{x})}{2}$$

and

$$f_{a_{2n+1}}(x) = \sum_{n=0}^{\infty} a_{2n+1} x^n = \frac{f(\sqrt{x}) - f(-\sqrt{x})}{2\sqrt{x}}$$

respectively.

The next Theorem presents the generating functions of even and odd-indexed generalized Tetranacci sequences.

**THEOREM 3.2.** *The generating functions of the sequences  $V_{2n}$  and  $V_{2n+1}$  are given by*

$$f_{V_{2n}}(x) = \frac{V_0 + (-3V_0 + V_2)x + (-2V_0 + V_1 - 2V_2 + V_3)x^2 + (-2V_2 + V_3)x^3}{x^4 + x^3 - 3x^2 - 3x + 1}$$

and

$$f_{V_{2n+1}}(x) = \frac{V_1 + (-3V_1 + V_3)x + (V_0 - V_1 + 2V_2 - V_3)x^2 + (V_0 + V_1 + V_2 - V_3)x^3}{x^4 + x^3 - 3x^2 - 3x + 1}$$

respectively.

*Proof.* Both statements are consequences of Lemma 3.1 applied to (1.6) and some lengthy work.

From the previous Theorem we get the following results as particular examples: the generating functions of the sequences  $M_{2n}$  and  $M_{2n+1}$  are given by

$$f_{M_{2n}}(x) = \frac{x^2 + x}{x^4 + x^3 - 3x^2 - 3x + 1}, \quad f_{M_{2n+1}}(x) = \frac{-x^2 - x + 1}{x^4 + x^3 - 3x^2 - 3x + 1}$$

and the generating functions of the sequences  $R_{2n}$  and  $R_{2n+1}$  are given by

$$f_{R_{2n}}(x) = \frac{x^3 - 6x^2 - 9x + 4}{x^4 + x^3 - 3x^2 - 3x + 1}, \quad f_{R_{2n+1}}(x) = \frac{x^3 + 2x^2 + 4x + 1}{x^4 + x^3 - 3x^2 - 3x + 1}.$$

The next Theorem presents the generating functions of even and odd-indexed Gaussian generalized Tetranacci sequences.

**THEOREM 3.3.** *The generating functions of the sequences  $GV_{2n}$  and  $GV_{2n+1}$  are given by*

$$f_{GV_{2n}} = \frac{GV_0 + (-3GV_0 + GV_2)x + (-2GV_0 + GV_1 - 2GV_2 + GV_3)x^2 + (-2GV_2 + GV_3)x^3}{x^4 + x^3 - 3x^2 - 3x + 1}$$

and

$$f_{GV_{2n+1}} = \frac{GV_1 + (-3GV_1 + GV_3)x + (GV_0 - GV_1 + 2GV_2 - GV_3)x^2 + (GV_0 + GV_1 + GV_2 - GV_3)x^3}{x^4 + x^3 - 3x^2 - 3x + 1}$$

respectively.

*Proof.* Both statements are consequences of Lemma 3.1 applied to (2.5) and some lengthy algebraic calculations.

The previous theorem gives the following two corollaries as particular examples. Firstly, the next one presents the generating functions of even and odd-indexed Gaussian Tetranacci sequences.

**COROLLARY 3.4.** *The generating functions of the sequences  $GM_{2n}$  and  $GM_{2n+1}$  are given by*

$$(3.4) \quad f_{GM_{2n}} = \frac{(1+i)x + (1-i)x^2 - ix^3}{x^4 + x^3 - 3x^2 - 3x + 1}$$

and

$$(3.5) \quad f_{GM_{2n+1}} = \frac{1 - (1-i)x - (1-i)x^2}{x^4 + x^3 - 3x^2 - 3x + 1}$$

respectively.

The following Corollary gives the generating functions of even and odd-indexed Gaussian Tetranacci-Lucas sequences.

COROLLARY 3.5. *The generating functions of the sequences  $GR_{2n}$  and  $GR_{2n+1}$  are given by*

$$(3.6) \quad f_{GR_{2n}}(x) = \frac{(4-i) - (9-4i)x - (6-7i)x^2 + (1+i)x^3}{x^4 + x^3 - 3x^2 - 3x + 1}$$

and

$$(3.7) \quad f_{GR_{2n+1}}(x) = \frac{(1+4i) + (4-9i)x + (2-6i)x^2 + (1+i)x^3}{x^4 + x^3 - 3x^2 - 3x + 1}$$

respectively.

The next Corollary present identities between Gaussian Tetranacci and Gaussian Tetranacci-Lucas sequences.

COROLLARY 3.6. *We have the following identities:*

$$\begin{aligned} & (4-i)GM_{2n} - (9-4i)GM_{2n-2} - (6-7i)GM_{2n-4} + (1+i)GM_{2n-6} \\ = & (1+i)GR_{2n-2} + (1-i)GR_{2n-4} - iGR_{2n-6}, \\ & (1+4i)GM_{2n} + (4-9i)GM_{2n-2} + (2-6i)GM_{2n-4} + (1+i)GM_{2n-6} \\ = & (1+i)GR_{2n-1} + (1-i)GR_{2n-3} - iGR_{2n-5}, \\ & (4-i)GM_{2n+1} - (9-4i)GM_{2n-1} - (6-7i)GM_{2n-3} + (1+i)GM_{2n-5} \\ = & GR_{2n} - (1-i)GR_{2n-2} - (1-i)GR_{2n-4} \\ & (1+4i)GM_{2n+1} + (4-9i)GM_{2n-1} + (2-6i)GM_{2n-3} + (1+i)GM_{2n-5} \\ = & GR_{2n+1} - (1-i)GR_{2n-1} - (1-i)GR_{2n-3} \end{aligned}$$

*Proof.* From (3.4) and (3.6) we obtain

$$((4-i) - (9-4i)x - (6-7i)x^2 + (1+i)x^3)f_{GM_{2n}} = ((1+i)x + (1-i)x^2 - ix^3)f_{GR_{2n}}.$$

The LHS (left hand side) is equal to

$$\begin{aligned} LHS &= ((4-i) - (9-4i)x - (6-7i)x^2 + (1+i)x^3) \sum_{n=0}^{\infty} GM_{2n}x^n \\ &= (5-i)x^2 + (5+3i)x + \sum_{n=3}^{\infty} ((4-i)GM_{2n} - (9-4i)GM_{2n-2} \\ &\quad - (6-7i)GM_{2n-4} + (1+i)GM_{2n-6})x^n \end{aligned}$$

whereas the RHS is

$$\begin{aligned} RHS &= ((1+i)x + (1-i)x^2 - ix^3) \sum_{n=0}^{\infty} GR_{2n}x^n \\ &= (5-i)x^2 + (5+3i)x + \sum_{n=3}^{\infty} ((1+i)GR_{2n-2} + (1-i)GR_{2n-4} - iGR_{2n-6})x^n. \end{aligned}$$

Compare the coefficients and the proof of the first identity is done. The other identities can be proved similarly by using (3.4)-(3.7).

We present an identity related with Gaussian general Tetranacci numbers and Tetranacci numbers.

**THEOREM 3.7.** *For  $n \geq 0$  and  $m \geq 0$  the following identity holds:*

$$(3.8) \quad GV_{m+n} = M_{m-2}GV_{n+3} + (M_{m-3} + M_{m-4} + M_{m-5})GV_{n+2} + (M_{m-3} + M_{m-4})GV_{n+1} + M_{m-3}GV_n$$

*Proof.* We prove the identity by strong induction on  $m$ . If  $m = 0$  then

$$GV_n = M_{-2}GV_{n+3} + (M_{-3} + M_{-4} + M_{-5})GV_{n+2} + (M_{-3} + M_{-4})GV_{n+1} + M_{-3}GV_n$$

which is true because  $M_{-2} = 0, M_{-3} = 1, M_{-4} = -1, M_{-5} = 0,$ . Assume that the equality holds for  $m \leq k$ .

For  $m = k + 1$ , we have

$$\begin{aligned} GV_{(k+1)+n} &= GV_{n+k} + GV_{n+k-1} + GV_{n+k-2} + GV_{n+k-3} \\ &= (M_{k-2}GV_{n+3} + (M_{k-3} + M_{k-4} + M_{k-5})GV_{n+2} + (M_{k-3} + M_{k-4})GV_{n+1} + M_{k-3}GV_n) \\ &\quad + (M_{k-3}GV_{n+3} + (M_{k-4} + M_{k-5} + M_{k-6})GV_{n+2} + (M_{k-4} + M_{k-5})GV_{n+1} + M_{k-4}GV_n) \\ &\quad + (M_{k-4}GV_{n+3} + (M_{k-5} + M_{k-6} + M_{k-7})GV_{n+2} + (M_{k-5} + M_{k-6})GV_{n+1} + M_{k-5}GV_n) \\ &\quad + (M_{k-5}GV_{n+3} + (M_{k-6} + M_{k-7} + M_{k-8})GV_{n+2} + (M_{k-6} + M_{k-7})GV_{n+1} + M_{k-6}GV_n) \\ &= (M_{k-2} + M_{k-3} + M_{k-4} + M_{k-5})GV_{n+3} \\ &\quad + ((M_{k-3} + M_{k-4} + M_{k-5} + M_{k-6}) + (M_{k-4} + M_{k-5} + M_{k-6} + M_{k-7})) \\ &\quad + (M_{k-5} + M_{k-6} + M_{k-7} + M_{k-8})GV_{n+2} \\ &\quad + ((M_{k-3} + M_{k-4} + M_{k-5} + M_{k-6}) + (M_{k-4} + M_{k-5} + M_{k-6} + M_{k-7}))GV_{n+1} \\ &\quad + (M_{k-3} + M_{k-4} + M_{k-5} + M_{k-6})GV_n \\ &= M_{k-1}GV_{n+3} + (M_{k-2} + M_{k-3} + M_{k-4})GV_{n+2} + (M_{k-2} + M_{k-3})GV_{n+1} + M_{k-2}GV_n \\ &= M_{(k+1)-2}GV_{n+3} + (M_{(k+1)-3} + M_{(k+1)-4} + M_{(k+1)-5})GV_{n+2} \\ &\quad + (M_{(k+1)-3} + M_{(k+1)-4})GV_{n+1} + M_{(k+1)-3}GV_n \end{aligned}$$

By strong induction on  $m$ , this proves (3.8).



The previous Theorem gives the following results as particular examples: For  $n \geq 0$  and  $m \geq 0$ , we have (taking  $GV_n = GM_n$ )

$$GM_{m+n} = M_{m-2}GM_{n+3} + (M_{m-3} + M_{m-4} + M_{m-5})GM_{n+2} + (M_{m-3} + M_{m-4})GM_{n+1} + M_{m-3}GM_n$$

and (taking  $GV_n = GR_n$ )

$$GR_{m+n} = M_{m-2}GR_{n+3} + (M_{m-3} + M_{m-4} + M_{m-5})GR_{n+2} + (M_{m-3} + M_{m-4})GR_{n+1} + M_{m-3}GR_n.$$

#### 4. Matrix Formulation of $GV_n$

Now, consider the sequence  $\{U_n\}$  which is defined by the fourth-order recurrence relation

$$U_n = U_{n-1} + U_{n-2} + U_{n-3} + U_{n-4}, \quad U_0 = U_1 = 0, U_2 = U_3 = 1.$$

Next, we present the first few values of numbers  $U_n$  with positive and negative subscripts in the following Table 5:

Table 5. A few values of the numbers  $U_n$

$n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	...
$U_n$	0	0	1	1	2	4	8	15	29	56	108	208	401	773	1490	...
$U_{-n}$	0	0	1	-1	0	0	2	-3	1	0	4	-8	5	-1	8	...

Note that some authors call  $\{U_n\}$  as a Tetranacci sequence instead of  $\{M_n\}$ . The numbers  $U_n$  can be expressed using Binet's formula

$$U_n = \frac{\alpha^n}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)} + \frac{\beta^n}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)} + \frac{\gamma^n}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)} + \frac{\delta^n}{(\delta - \alpha)(\delta - \beta)(\delta - \gamma)}.$$

The matrix method is very useful method in order to obtain some identities for special sequences. We define the square matrix  $A$  of order 4 as:

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

such that  $\det M = -1$ . Induction proof may be used to establish

$$(4.1) \quad A^n = \begin{pmatrix} M_{n+1} & M_n + M_{n-1} + M_{n-2} & M_n + M_{n-1} & M_n \\ M_n & M_{n-1} + M_{n-2} + M_{n-3} & M_{n-1} + M_{n-2} & M_{n-1} \\ M_{n-1} & M_{n-2} + M_{n-3} + M_{n-4} & M_{n-2} + M_{n-3} & M_{n-2} \\ M_{n-2} & M_{n-3} + M_{n-4} + M_{n-5} & M_{n-3} + M_{n-4} & M_{n-3} \end{pmatrix}$$

$$(4.2) \quad = \begin{pmatrix} U_{n+2} & U_{n+1} + U_n + U_{n-1} & U_{n+1} + U_n & U_{n+1} \\ U_{n+1} & U_n + U_{n-1} + U_{n-2} & U_n + U_{n-1} & U_n \\ U_n & U_{n-1} + U_{n-2} + U_{n-3} & U_{n-1} + U_{n-2} & U_{n-1} \\ U_{n-1} & U_{n-2} + U_{n-3} + U_{n-4} & U_{n-2} + U_{n-3} & U_{n-2} \end{pmatrix}$$

$$(4.3) \quad = \begin{pmatrix} U_{n+2} & U_{n+2} - U_{n-2} & U_{n+1} + U_n & U_{n+1} \\ U_{n+1} & U_{n+1} - U_{n-3} & U_n + U_{n-1} & U_n \\ U_n & U_n - U_{n-4} & U_{n-1} + U_{n-2} & U_{n-1} \\ U_{n-1} & U_{n-1} - U_{n-5} & U_{n-2} + U_{n-3} & U_{n-2} \end{pmatrix}.$$

Matrix formulation of  $M_n$  and  $R_n$  can be given as

$$(4.4) \quad \begin{pmatrix} M_{n+3} \\ M_{n+2} \\ M_{n+1} \\ M_n \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^n \begin{pmatrix} M_3 \\ M_2 \\ M_1 \\ M_0 \end{pmatrix}$$

and

$$(4.5) \quad \begin{pmatrix} R_{n+3} \\ R_{n+2} \\ R_{n+1} \\ R_n \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^n \begin{pmatrix} R_3 \\ R_2 \\ R_1 \\ R_0 \end{pmatrix}.$$

Induction proofs may be used to establish the matrix formulations  $M_n$  and  $R_n$ . Note that

$$GM_n = iU_n + U_{n+1}.$$

Consider the matrices  $N_M, E_M$  defined by as follows:

$$N_M = \begin{pmatrix} 2+i & 1+i & 1 & 0 \\ 1+i & 1 & 0 & 0 \\ 1 & 0 & 0 & i \\ 0 & 0 & i & 1-i \end{pmatrix}$$

$$E_M = \begin{pmatrix} GM_{n+3} & GM_{n+2} & GM_{n+1} & GM_n \\ GM_{n+2} & GM_{n+1} & GM_n & GM_{n-1} \\ GM_{n+1} & GM_n & GM_{n-1} & GM_{n-2} \\ GM_n & GM_{n-1} & GM_{n-2} & GM_{n-3} \end{pmatrix}$$

Next Theorem presents the relations between  $A^n, N_M$  and  $E_M$ .

**THEOREM 4.1.** *For  $n \geq 3$ , we have*

$$A^n N_M = E_M.$$

*Proof.* Using the relation

$$GM_n = iU_n + U_{n+1},$$

and the calculations

$$\begin{aligned} a &= (2+i)U_n + (1+i)U_{n-1} + (2+i)U_{n+1} + (2+i)U_{n+2} \\ &= 2U_n + iU_n + U_{n-1} + iU_{n-1} + 2U_{n+1} + iU_{n+1} + 2U_{n+2} + iU_{n+2} \\ &= i(U_{n+2} + U_{n+1} + U_n + U_{n-1}) + (2U_{n+2} + 2U_{n+1} + 2U_n + U_{n-1}) \\ &= iU_{n+3} + (U_{n+2} + U_{n+1} + U_n + (U_{n+2} + U_{n+1} + U_n + U_{n-1})) \\ &= iU_{n+3} + (U_{n+2} + U_{n+1} + U_n + U_{n+3}) = iU_{n+3} + U_{n+4} = GM_{n+3} \end{aligned}$$

and

$$\begin{aligned} U_n + U_{n-1} + U_{n+1} + (1+i)U_{n+2} &= U_n + U_{n-1} + U_{n+1} + U_{n+2} + iU_{n+2} \\ &= iU_{n+2} + U_{n+3} = GM_{n+2}, \end{aligned}$$

we get

$$\begin{aligned}
 A^n N_M &= \begin{pmatrix} U_{n+2} & U_{n+1} + U_n + U_{n-1} & U_{n+1} + U_n & U_{n+1} \\ U_{n+1} & U_n + U_{n-1} + U_{n-2} & U_n + U_{n-1} & U_n \\ U_n & U_{n-1} + U_{n-2} + U_{n-3} & U_{n-1} + U_{n-2} & U_{n-1} \\ U_{n-1} & U_{n-2} + U_{n-3} + U_{n-4} & U_{n-2} + U_{n-3} & U_{n-2} \end{pmatrix} \begin{pmatrix} 2+i & 1+i & 1 & 0 \\ 1+i & 1 & 0 & 0 \\ 1 & 0 & 0 & i \\ 0 & 0 & i & 1-i \end{pmatrix} \\
 &= \begin{pmatrix} GM_{n+3} & GM_{n+2} & GM_{n+1} & GM_n \\ GM_{n+2} & GM_{n+1} & GM_n & GM_{n-1} \\ GM_{n+1} & GM_n & GM_{n-1} & GM_{n-2} \\ GM_n & GM_{n-1} & GM_{n-2} & GM_{n-3} \end{pmatrix}.
 \end{aligned}$$

Above Theorem can be proved by mathematical induction as well.

Consider the matrices  $N_R, E_R$  defined by as follows:

$$\begin{aligned}
 N_R &= \begin{pmatrix} 7+3i & 3+i & 1+4i & 4-i \\ 3+i & 1+4i & 4-i & -1-i \\ 1+4i & 4-i & -1-i & -1-i \\ 4-i & -1-i & -1-i & -1+7i \end{pmatrix} \\
 E_R &= \begin{pmatrix} GR_{n+3} & GR_{n+2} & GR_{n+1} & GR_n \\ GR_{n+2} & GR_{n+1} & GR_n & GR_{n-1} \\ GR_{n+1} & GR_n & GR_{n-1} & GR_{n-2} \\ GR_n & GR_{n-1} & GR_{n-2} & GR_{n-3} \end{pmatrix}.
 \end{aligned}$$

The following Theorem presents the relations between  $A^n, N_R$  and  $E_R$ .

**THEOREM 4.2.** *We have*

$$A^n N_R = E_R.$$

*Proof.* The proof requires some lengthy calculation, so we omit it.

The previous Theorem, also, can be proved by mathematical induction.

Similarly, matrix formulation of  $V_n$  can be given as

$$\begin{pmatrix} V_{n+3} \\ V_{n+2} \\ V_{n+1} \\ V_n \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^n \begin{pmatrix} V_3 \\ V_2 \\ V_1 \\ V_0 \end{pmatrix}$$

Consider the matrices  $N_V, E_V$  defined by as follows:

$$N_V = \begin{pmatrix} ic_2+c_3 & ic_1+c_2 & ic_0+c_1 & a_1 \\ ic_1+c_2 & ic_0+c_1 & (1-i)c_0-ic_1-ic_2+ic_3 & a_2 \\ ic_0+c_1 & (1-i)c_0-ic_1-ic_2+ic_3 & (1-i)c_3-c_1-(1-2i)c_2-c_0 & a_3 \\ (1-i)c_0-ic_1-ic_2+ic_3 & (1-i)c_3-c_1-(1-2i)c_2-c_0 & 2ic_1+(2-i)c_2-c_3 & a_4 \end{pmatrix},$$

$$E_V = \begin{pmatrix} GV_{n+3} & GV_{n+2} & GV_{n+1} & GV_n \\ GV_{n+2} & GV_{n+1} & GV_n & GV_{n-1} \\ GV_{n+1} & GV_n & GV_{n-1} & GV_{n-2} \\ GV_n & GV_{n-1} & GV_{n-2} & GV_{n-3} \end{pmatrix}.$$

where

$$\begin{aligned} a_1 &= (1-i)c_0 - ic_1 - ic_2 + ic_3 \\ a_2 &= (1-i)c_3 - c_1 - (1-2i)c_2 - c_0 \\ a_3 &= 2ic_1 + (2-i)c_2 - c_3 \\ a_4 &= 2ic_0 + (2-i)c_1 - c_2. \end{aligned}$$

We now present our final Theorem.

**THEOREM 4.3.** *We have*

$$A^n N_V = E_V.$$

*Proof.* The proof requires some lengthy work, so we omit it.

## 5. Conclusions

- In the section (1), we present some background about generalized Tetranacci numbers and Gaussian numbers.
- In the section (2), we define Gaussian generalized Tetranacci numbers and as special cases, we investigate Gaussian Tetranacci and Gaussian Tetranacci-Lucas numbers with their properties such as the generating functions, Binet's formulas and sums formulas of these Gaussian numbers.
- In the section (3), we obtain some identities of Gaussian Tetranacci numbers and Gaussian Tetranacci-Lucas numbers.
- In the section (4), we give matrix formulation of Gaussian generalized Tetranacci numbers.

### Competing Interests

Author have declared that no competing interests exist.

## References

- [1] Asci M., and Gurel E., Gaussian Jacobsthal and Gaussian Jacobsthal Polynomials, Notes on Number Theory and Discrete Mathematics, Vol.19, pp.25-36, 2013.
- [2] Bacani, J. B., Rabago, J. F. T., On Generalized Fibonacci Numbers, Applied Mathematical Sciences, 9 (25), 3611-3622, 2015.
- [3] Berzsenyi G., Gaussian Fibonacci Numbers, Fibonacci Quart., Vol.15(3), pp.233-236, 1977.
- [4] Catarino, P., and Campos, H., A note on Gaussian Modified Pell numbers, Journal of Information & Optimization Sciences, Vol. 39, No. 6, pp. 1363–1371, 2018.
- [5] G. P. Dresden G. P., Du, Z., A Simplified Binet Formula for k-Generalized Fibonacci Numbers, J. Integer Seq. 17, art. 14.4.7, 9 pp., 2014.
- [6] Fraleigh J.B., A First Course In Abstract Algebra, (2nd ed.), Addison-Wesley, Reading, ISBN 0-201-01984-1, 1976.
- [7] Frontczak, R., Convolutions for Generalized Tribonacci Numbers and Related Results, International Journal of Mathematical Analysis, Vol. 12, 2018, no. 7, 307 - 324.
- [8] E. Gurel, k-Order Gaussian Fibonacci and k-Order Gaussian Lucas Recurrence Relations, Ph.D Thesis, Pamukkale University Institute of Science Mathematics, Denizli, Turkey (2015).
- [9] Halici S., Öz S., On some Gaussian Pell and Pell-Lucas numbers, Ordu University Science and Technology Journal, Vol.6(1), pp.8-18, 2016.
- [10] Halici S., Öz S., On Gaussian Pell Polynomials and Their Some Properties, Palestine Journal of Mathematics, Vol 7(1), 251-256, 2018.
- [11] Harman C. J., Complex Fibonacci Numbers, Fibonacci Quart., Vol.19(1), pp. 82-86, 1981.
- [12] Hathiwala, G. S., Shah, D. V., Binet–Type Formula For The Sequence of Tetranacci Numbers by Alternate Methods, Mathematical Journal of Interdisciplinary Sciences 6 (1), 37–48, 2017.
- [13] Horadam, A. F., Complex Fibonacci Numbers and Fibonacci quaternions, Amer. Math. Monthly 70, 289–291, 1963.
- [14] Jordan J.H., Gaussian Fibonacci and Lucas Numbers, Fibonacci Quart., Vol.3, pp. 315-318, 1965.
- [15] Pethe S., Horadam A.F., Generalised Gaussian Fibonacci numbers, Bull. Austral. Math. Soc., Vol.33, pp.37-48, 1986.
- [16] Melham, R. S., Some Analogs of the Identity  $F_n^2 + F_{n+1}^2 = F_{2n+1}^2$ , Fibonacci Quarterly, 305-311, 1999.
- [17] Natividad, L. R., On Solving Fibonacci-Like Sequences of Fourth, Fifth and Sixth Order, International Journal of Mathematics and Computing, 3 (2), 2013.
- [18] Pethe S., Horadam A.F., Generalised Gaussian Lucas Primordial numbers, Fibonacci Quart., pp. 20-30, 1988.
- [19] Pethe, S., Some Identities for Tribonacci Sequences, The Fibonacci Quarterly, 26, 144-151, 1988.
- [20] Simson. R., An Explanation of an Obscure Passage in Albrecht Girard’s Commentary upon Simon Stevin’s Works, Philosophical Transactions of the Royal Society 48.1, 368-377, 1753.
- [21] Singh, B., Bhadouria, P., Sikhwal, O., Sisodiya, K., A Formula for Tetranacci-Like Sequence, Gen. Math. Notes, 20 (2), 136-141, 2014.
- [22] Sloane, N.J.A., The on-line encyclopedia of integer sequences, <http://oeis.org/>
- [23] Soykan, Y., Taşdemir, E., Okumuş, İ., Göcen, M., Gaussian Generalized Tribonacci Numbers, Journal of Progressive Research in Mathematics(JPRM), 14 (2), 2373-2387, 2018.
- [24] Taşcı, D., and Acar, H., Gaussian Tetranacci Numbers, Communications in Mathematics and Applications, 8 (3), 379–386, 2017.
- [25] Taşcı, D., and Acar, H., Gaussian Padovan and Gaussian Pell-Padovan Numbers, Commun. Fac. Sci. Ank. Ser. A1 Math. Stat., Volume 67, Number 2, pp. 82-88, 2018.

- [26] Yagmur, T., and Karaaslan, N., Gaussian Modified Pell Sequence and Gaussian Modified Pell Polynomial, Aksaray University Journal of Science and Engineering, Volume 2, Issue 1, pp. 63-72, 2018.
- [27] Yalavigi, C.C., A Note on 'Another Generalized Fibonacci Sequence', The Mathematics Student. 39, 407-408, 1971.
- [28] Waddill, Another Generalized Fibonacci Sequence, M. E., Fibonacci Quarterly., 5 (3), 209-227, 1967.
- [29] Waddill, M. E., The Tetranacci Sequence and Generalizations, The Fibonacci Quarterly, 9-20, 1992.
- [30] Zaveri, M. N., Patel, J. K., Binet's Formula for the Tetranacci Sequence, International Journal of Science and Research (IJSR), 5 (12), 1911-1914, 2016.