

PREVALENCE OF GBS SEROTYPES ASSOCIATED WITH ANOGENITAL
COLONIZATION AMONG PREGNANT WOMEN IN JOS

ABSTRACT

Aim

The aim of this study was to determine the prevalence of GBS serotype distribution in Jos University Teaching Hospital.

Materials and Methodology

This was a hospital based descriptive cross-sectional study of 300 women receiving health care at the Jos University Teaching Hospital between July 2017 and November 2017. Systematic sampling technique was employed in recruiting consenting subjects for this study. High vaginal and anorectal swabs were collected from each subject after obtaining their consent by signing a structured consent form. The identified GBS isolates were serotyped using immuLex strep-B antisera from SSI Diagnostica, 2 Herredysvejen, DK-3400 Hillerod Demnark to identify the different serotypes. The results obtained were computed using SPSS version 21.

Results

A total of 300 women obtaining health care in Jos University Teaching Hospital (JUTH) were enrolled in this study between the months of July, 2017 and November, 2017. In all, vaginal and anorectal swabs were taken from 200 pregnant women and 100 non-pregnant women. The age range of the study population was between 16 years to 48 years with a mean age of 31.9 year (SD \pm 6.6).

The prevalence rate among the study population was 6.3%. The colonization rate among pregnant and non-pregnant women was 6.5% and 6.0% respectively with no significant statistical difference.

28 Serotype Ia was the commonest isolate responsible for 42.1% of the GBS isolates. Serotype III
29 accounted for 31.6% of the isolates, followed by serotype V (15.8%). Serotype II was less common,
30 responsible for only 10.5%. Serotypes Ib, IV, VI, VII, VIII and IX were not isolated.

31 Conclusion

32 This study showed that GBS colonization rate among the study population was 6.3%. Approximately,
33 6.5% and 6.0% prevalence rate were found among pregnant and non-pregnant women respectively. Of
34 all the GBS isolates, serotypes Ia, II, III, and serotype V were isolated with serotype Ia being the most
35 prevalent serotype. Serotype Ib, IV, VI, VIII and IX were not isolated in this study. This knowledge of
36 serotype distribution will help in instituting serotype specific GBS vaccines for the prevention of GBS
37 diseases in Jos Plateau state, Nigeria.

38 Key words: *Streptococcus agalactiae*, Pregnant women, Serotypes

39 INTRODUCTION

40 *Streptococcus agalactiae* (Group B *Streptococcus*, GBS) is a normal flora of the genitourinary tract and
41 the rectum of humans [1]. *Streptococcus agalactiae* colonizes the human gastrointestinal and genital
42 tracts of 20-30% of healthy humans [2]. The colonization of the vagina by GBS can be transient,
43 chronic or intermittent and serves as potential sources of infection to the new born [3,4]. From this site,
44 the bacterium can colonize 50% - 75% of infants of colonized mothers via the amniotic fluid or the
45 mucous membranes during birth and produce an early onset disease (EOD) 24-48 hours after birth
46 [5,6]. About 1-2% of the new born of GBS colonized mothers may develop EOD with the maternal
47 genital area serving as source of infection [7,8].

48 Several maternal risk factors can significantly increase the possibility of development of GBS disease in
49 neonates. This is an important fact, since the presence of an isolated risk factor increases the
50 probability that a pregnant woman will have a child with early neonatal GBS disease by 6.5 times [9].
51 The identification of a high-risk population to be screened for GBS colonization is a considerable

52 challenge, since colonization is also observed in mothers who do not have the classically known risk
53 factors, and it represents 25% to 30% of neonates that develop early neonatal GBS [10].

54 GBS colonization rate has been found to be associated with prolonged labour, premature rupture of
55 membrane and preterm delivery [11]. Other factors equally associated with GBS colonization are low
56 levels of circulating maternal antibodies against GBS, maternal age of less than 20 years, and maternal
57 diabetes mellitus [10,12].

58 The frequency of GBS colonization is highest among black women, followed by Hispanic women and
59 white women [10]. However, in some countries, in view of the high degree of racial miscegenation, it is
60 difficult to establish a precise classification of a patient sample in terms of race or ethnic group making
61 it difficult to differentiate among races. Similar study in Saudi Arabia has reported no difference in GBS
62 prevalence among blacks and white Saudi women [13]. One important virulence factor of this GBS is
63 the capsular polysaccharide (CPS). This CPS antigen has been studied extensively and is important
64 antigenic marker. There are ten antigenically distinct serotypes identified, namely Ia, Ib, II, III, IV, V, VI,
65 VII, VIII, and IX [14]. Capsular type distribution varies with geographical region and ethnic origin of the
66 population [15]. In the United States and Europe, the most prevalent serotypes in human infections are
67 Ia, II, III, and V, whereas serotypes VI and VIII are the most predominant in Japan [16,17].

68 A study conducted in Gabon revealed that serotypes III and V are the most prevalent serotypes in
69 Gabon similar to what was found in South Africa [18,19]. This knowledge of serotype distribution is
70 necessary for the selection and development of serotype-based vaccine for the prevention of invasive
71 disease in a given country [20,21,22].

72 2.0 MATERIALS AND METHODS

73 2.1 Study Area

74 The study was carried out in Jos University Teaching Hospital (JUTH). Jos University Teaching Hospital
75 is located in Jos the Plateau State capital. The hospital is a tertiary health institution with a 600 beds

76 capacity serving Plateau State and majority of the states in the North-central and part of North-east
77 geopolitical zones of Nigeria. Jos University Teaching Hospital is also a centre for AIDS Prevention
78 Initiative in Nigeria (APIN) that cater for most people living with HIV (PLHIV) from within and the
79 bordering states. The main occupation of the people is farming with majority of them in the city being
80 civil servants and businessmen and women.

81 2.2 Study Population

82 The study population included 200 pregnant and 100 non-pregnant women attending antenatal clinic at
83 the Jos University Teaching Hospital between July 2017 and November 2017.

84

85

86 2.3 Study Design

87 The study was a hospital based descriptive, cross-sectional study that recruited 300 consenting
88 pregnant and non-pregnant women attending antenatal and gynaecology clinics at the Jos University
89 teaching Hospital

90 2.4 Sample Collection

91 Anorectal and vaginal swabs were carefully collected from 200 pregnant women and 100 non-pregnant
92 women using sterile swab sticks by the attending physicians after given them appropriate instructions
93 on how the sample should be collected [23].

94 2.5 Specimen Transport

95 The collected specimens were immediately inoculated into a selective enrichment broth, Todd-Hewitt
96 broth (Oxoid LTD) supplemented with gentamycin (8ug/ml), nalidixic acid (15ug/ml) and 5% sheep

97 blood to increase the recovery rate of GBS [16,23]. These were transported to the laboratory within
98 three hours of inoculation.

99 2.6 Culture and Incubation

100 The tubes of inoculated Todd-Hewitt broth were incubated aerobically at 37°C for 18 to 24 hours. After
101 an overnight incubation, the broths were subcultured onto 10% sheep blood agar and chromatic Strepto
102 B agar (Liofilchem, Italy), a selective medium for GBS.

103 The inoculated 10% sheep blood agar plates were incubated aerobically in 5-10% CO₂ (candle
104 extinction jar) at 37°C for 18 to 24 hours, while the inoculated chromatic Strepto B agar plates were
105 incubated aerobically at 37°C for 18 to 24 hours [24]. The *Streptococcus agalactiae* control strain was
106 also inoculated onto 10% sheep blood agar and chromatic Strepto B agar and incubated as stated
107 above respectively.

108

109 2.7 Identification of GBS Isolates

110 GBS isolates were identified by their beta haemolytic pattern on 5% sheep blood agar and blue-green
111 colour on chromatic Strepto B agar. The isolates were further subjected to Gram staining, catalase test,
112 and Serogrouping using streptococcal grouping kit (DR0585A OXOID) from Oxoid.

113 2.8 Serotyping of Isolates

114 The group B B-haemolytic (GBS) isolates were serotyped using immuLex™ strep-B antisera from SS1
115 Diagnostica, 2 Herredysvejen, DK-3400 Hillerød Denmark to identify the different serotypes according
116 to manufacturers guidelines.

117 2.9 Data Analysis

The data obtained from the study were analysed using Statistical Package for Social Sciences (SPSS) version 21 (IBM SPSS Inc, USA). Proportions were compared using Chi-square with confidence limit (p-value) of < 0.05 considered significant.

3.0 RESULTS

A total of 300 women obtaining health care in Jos University Teaching Hospital (JUTH) were enrolled in this study between the months of July, 2017 and November, 2017. In all, vaginal and anorectal swabs were taken from 200 pregnant women and 100 non-pregnant women. The age range of the study population was between 16 years to 48 years with a mean age of 31.9 years (SD 16.6).

The prevalence rate among the study population was 19(6.3%). The pregnant women were colonized in 6.5% while the colonization rate was 6.0% among the non-pregnant women. The difference in colonization between pregnant and non-pregnant was not statistically significant (Table 1) ($X^2 = 0.028$, $P = .87$).

Out of the 19 isolates of GBS isolated, 42.1% belonged to serotype Ia, which was the most prevalent serotype among the study population. Serotype III accounted for 31.6% of the isolates, followed by serotype V (15.8%). Serotype II was less common responsible for only 10.5% of the 19 isolates. Serotypes Ib, IV, VI, VII, VIII and IX were not isolated.

Among the pregnant women, serotypes Ia and III were isolated in 53.8% and 30.8% respectively while serotype II and V accounted for 7.7% each. In non-pregnant women, serotype III and V were the most common serotypes accounting for 33.3% each while serotype Ia and II had equal distribution, responsible for 16.7% each. The serotypes distribution among the pregnant and non-pregnant was not statistically significant (Table 2) ($X^2 = 3.380$, $P = .34$).

Table I

Group B Streptococcal carriage rates among pregnant and non-pregnant women in Jos University Teaching Hospital.

Category of women	No. tested	No. Positive	% Positive
Pregnant	200	13	6.5
Non-Pregnant	100	6	6.0
Total	300	19	6.3

$$\chi^2 = 0.028 \quad P = .87 \quad df = 1$$

Table 2

Serotypes distribution of Streptococcus agalactiae isolates from pregnant and non-pregnant Women in Jos University Teaching Hospital

Serotype	Pregnant (%)	Non-pregnant (%)	Total
Ia	7(53.8)	1(16.7)	8(42.1)
II	1(7.7)	1(16.7)	2(10.5)
III	4(30.8)	2(33.3)	6(31.6)
V	1(7.7)	2(33.3)	3(15.8)
Total	13(6.5)	6(6.0)	19(6.3)

$$\chi^2 = 3.380 \quad P = .34 \quad df = 3$$

4.0 DISCUSSION

The study revealed GBS colonization rate of 19(6.3%) among the population tested. The carriage rate was 6.5% in pregnant women and 6.0% in non-pregnant women. This result is lower than the 7.0% previously reported by Nsagha et al (1997) in Jos [25]. The slight decrease in the colonization rate may be attributed to improve health awareness among the general public and, improvement in culturing technique as Todd-Hewitt broth and chromatic Strepto B agar which are selective media for *Streptococcus agalactiae* were used in this study making identification and differentiation easy rather than just blood agar.

In other parts of Nigeria, 19.0% was reported in Ibadan by Onile (1990) [26] and Uhiara (1993) reported a carriage rate of 9.0% in Calaber [27]. Onipede and his colleagues in 2012 reported a higher prevalence rate of 11.3% in Ile-Ife [28] while 9.8% was reported by Okon et al 2013 in North-eastern Nigeria [29].

The group B streptococcal colonization rate observed in this study was also lower compare to colonization rates reported from several African countries. About 31.6% was reported in Zimbabwe by Moyo et al (2000) [30], 25.3% in Egypt by Shabayek et al (2014) [31] and 23% in Tanzania [6].

When compared with studies conducted in developed countries, it was observed that the result of this study was lower to what was obtained in United States of America [32]. Tor-Udom et al (2006) reported a carriage rate of 16.0% in Thailand [33] while 24.0% was reported in Belgium in 2009 [34]. A similar study conducted in Poland by Brzychczy-Wloch et al (2013) reported 29.5% carriage rate [35]. These results are higher to 5.7% obtained in Israel [36]. The variations between countries could possibly be due, at least in part, to differences in sampling and culturing techniques, types of media used as Well as the population studied [37]. For instance, in this study, samples were collected from HIV positive and HIV negative pregnant women regardless of gestational age.

Serotyping of the GBS isolates revealed that the isolates belong to serotypes Ia, II, III, and V whereas serotypes Ib, IV, VI, VII, VIII and IX were not isolated. As indicated in table above, serotype Ia (42.1%) was the predominant serotype followed by serotype III (31.6%). Serotypes II and V were isolated in 10.5% and 15.8% respectively. The isolated serotypes have been reported as the predominant causes of human infection worldwide [38] and serotype III as the most prevalent and invasive of all the serotypes [39]. Capsular type distribution varies between countries, geographical region and ethnic origin of a population [15]. A study conducted in Gabon revealed that serotype V and III are the most common, which is in consistent with findings in South Africa [18,19]. Serotypes Ia, Ib and II have been reported in Brazil [40] as the predominant serotypes causing most of the infections there. In United States of America and Europe, serotypes Ia, II, III, and V have been isolated in 80-90% of clinical infections while serotypes VI and VIII are the most predominant in Japan.

This knowledge of serotype distribution is necessary for the selection and development of serotype-based vaccine for the prevention of invasive disease in a given country [20].

5.0 CONCLUSION

This study showed that GBS colonization rate among the study population was 6.3%. Approximately, 6.5% and 6.0% prevalence rate were found among pregnant and non-pregnant women respectively. Of all the GBS isolates, serotypes Ia, II, III, and serotype V were isolated with serotype Ia being the most prevalent serotype. Serotype Ib, IV, VI, VIII and IX were not isolated in this study. This knowledge of serotype distribution will help in instituting serotype specific GBS vaccines for the prevention of GBS diseases in Jos Plateau state, Nigeria.

204 **ETHICAL APPROVAL**

205 This study was approved by the research ethical committee of Jos University Teaching Hospital with
206 reference number JUTH/DCS/ADM/127/XIX/6583. Written informed consents were also signed by all
207 subjects before enrolment in the study.

208 **CONSENT**

209 As per international standard or university standard, patients' written consent has been collected and
210 preserved by the authors.

211 **COMPETING INTERESTS**

212 Authors have declared that no competing interests exist.

213 **REFERENCE**

- 214 1. Keefe, G.P. Streptococcus agalactiae mastitis: a review. The Can Vet J. 1997;38(7):429-437.
- 215 2. Kathryn AP, Philip AW, Berenice R, John DH, John RT, Kelly SD. Streptococcus salivarius K12
216 limits group B Streptococcus vaginal colonization. Infect Immun. 2015;83(9):3438-3444.
- 217 3. Chakshu G, Laurence EG. Comparison of two culture media and three sampling techniques for
218 sensitive and rapid screening of vaginal colonization by group B streptococcus in pregnant
219 women. J Clinical Microbiol. 2004;42(9):3975-3977.
- 220 4. Verani JR, McGee L, Schrag SJ. Prevention of perinatal group B streptococcal disease:
221 Revised guidelines from CDC. MMWR. 2010;59(R10):1-36.
- 222 5. Fatemi F, Chamani L, Pakzad P, Zeraati H, Rabbani H, Asgari S. Colonization rate of group B
223 Streptococcus (GBS) in pregnant women using GBS agar medium. Acta Med Iran. 2009;47(1):
224 25-30.

6. Joachim A, Matee MI, Massawe FA, Lyamuya EF. Maternal and neonatal colonisation of group B streptococcus at Muhimbili National Hospital in Dar es Salaam, Tanzania: prevalence, risk factors and antimicrobial resistance. BMC Public Health, 2009;9:437.
7. Chung MY, K0 DJ, Chen CC, Huang CB, Chung CH, Chen FS, Hwang KP. Neonatal group B streptococcal infection: a 7-year experience. Chang Gung Med J. 2004;27(7):501-508.
8. Melin P. Neonatal group B streptococcal disease: from pathogenesis to prevention strategies. Clin Microbiol Infect. 2011;17(9):1294-1303.
9. Boyer KM, Gotoff SP. Prevention of early-onset neonatal group B streptococcal disease with selective intrapartum chemoprophylaxis. N Engl J Med. 1986;314:1665-1669.
10. Centres for Disease Control and Prevention (CDC). (2002). Prevention of Perinatal Group B Streptococcal Disease: Revised Guidelines from CDC. MMWR, 2002;51:1-22.
11. Aali B, Abdollahi H, Nakhaee N, Davazdahemami Z, Mehdizadeh A. The association of preterm labor with vaginal colonization of group B streptococci. IJRM. 2007; 5(4):191-194.
12. Lin FY, Weisman LE, Azimi PH, Joseph BP, Penny CJ, Regan GG, et al. Level of maternal IgG anti-group B streptococcus type III antibody correlated with protection of neonates against early-onset disease caused by this pathogen. J Infect. Dis.2004;190(5): 928-934.
13. Milyani R, Rokbah AR. Factors affecting vaginal and rectal carriage rate of Streptococcus agalactiae among pregnant and nonpregnant Saudi women. Adv Sch Med. 2011;1(3): 26-32.
14. Slotved HC, Kong F, Lambertsen L, Sauer S, Gilbert GL. Latex assay for serotyping of group B Streptococcus isolates. J Clin Microb. 2007; 45(9):2929-2936.
15. Manning SD, Tallman P, Baker CJ, Gillespie B, Marrs CF, Foxman B. Determinants of colonization with group B Sytreptococcus among heterosexual college couples. Epidemiol. 2002; 13:533-537.
16. Mario MS, Valenzuela I, Vasquez AE, Illane SE. Prevention of early-onset neonatal group B streptococcal disease. Rev Obstet Gyne. 2013; 6(2):63-68.

17. Martina OC, Rooyen TM, Charles MM, John YB, Sogolo LL, Motiatji RM, Maphoshane N, Sylvester RM. *Germes*, 5(4), 125-133.
18. Madzivhandila M, Adrian PV, Cutland CL, Kuwanda L, Schrag SJ, Madhi SA. Serotype Distribution and Invasive Potential of Group B Streptococcus Isolates Causing Disease in Infants and Colonizing Maternal-Newborn Dyads. *PLOS*.2011; 6(3):17861.
19. Sabine B, Nicole T, Meskure C, Ghyslain M, Rella Z, Mirjam G, et al. Streptococcus agalactiae serotype distribution and antimicrobial susceptibility in pregnant women in Gabon, Central Africa. *Sci Rep*. 2015; 5:17281.
20. Fluegge K, Supper S, Siedler A, Berner R. Serotype distribution of invasive group B streptococcal isolates in infants: Results from a nationwide active laboratory surveillance study over 2 years in Germany. *Clin Infect Dis*. 2005; 40(5):760-763.
21. Johri AK, Paoletti LC, Glaser P, Dua M, Shanna PK, Grandi G, et al. Group B Streptococcus: global incidence and vaccine development. *Nat Rev Microbiol*. 2006; 4(12):932-942.
22. Baker CJ, Carey VJ, Rench MA, Edwards MS, Hillier SH, Kasper DL, et al. Maternal Antibody at Delivery Protects Neonates from Early Onset Group B Streptococcal Disease Maternal Antibody at Delivery Protects Neonates from Early Onset group B Streptococcal Disease. *J Infect Dis*. 2014; 209:1781-788.
23. CDC. Prevention of perinatal group B streptococcal disease-revised guideline from centres for disease control (CDC). *MMWR*. 2010;59(10):1-36.
24. Dunne WM. Comparison of selective broth medium plus neomycin-nalidixic acid agar and selective broth medium plus Columbia colistin-nalidixic acid agar for detection of group B streptococcal colonization in women. *J Clin Microbiol*. 1999;37(11):3705-3706.
25. Nsagha DS, Bello CSS, Kandakai-Olukemi VT. Maternal carriage in pregnancy of Group B streptococcus in Jos: Relation of Endocervical and Anorectal colonization. *NigQt J Hosp Med*. 1997; 7:53-56.

26. Onile BA. Group B Streptococcus carriage in Nigeria. *Trans R Soc Trop Med Hyg.* 1990; 74:367-370.
27. Uhiara, J.E. (1993). Group B streptococcal carriage among parturient and their neonates in Zaria, Nigeria. *Afr J Med.*1993; 22(3):78-83.
28. Onipede A, Adefusi O, Adeyemia, Adejuyigbe E, Oyelese A, Ogunniyi T. Group B streptococcus carriage during late pregnancy in Ile-Ife, Nigeria. *Afr J Clin Exper Microbiol.* 2012;13(3):135-143.
29. Okon KO, Usman H, Umar Z, Balogun ST. Prevalence of group B streptococcus colonization among pregnant women attending antenatal clinic of a tertiary hospital in Northeastern, Nigeria. *Am J Res Commun.* 2013; 1(6):54-66.
30. Moyo SR, Mudzori J, Tswana SA, Maeland JA. Prevalence, capsular type distribution, anthropometric and obstetric factors of group B streptococcus (*Streptococcus agalactiae*) colonization in pregnancy. *Cent Afr J Med.* 2000; 46(5):115-120.
31. Shabayek S, Abdalla S, Abouzeid AM. Serotype and surface protein gene distribution of colonizing group B streptococcus in women in Egypt. *Epidemiol Infect.* 2014; 142(1):208-210.
32. McKenna, D.S., Matson, S., & Northern, I. (2003). Maternal group B streptococcal (GBS) genital tract colonization at term in women who have asymptomatic GBS bacteriuria. *Infect Dis J Obstet Gyne.* 2003; 11(4):203-207.
33. Tor-Udom S, Tor-Udom P, Hiriote W. The prevalence of streptococcus agalactiae (group B) colonization in pregnant women at Thammasat Hospital. *J Med Asso Thai.* 2006; 89(4): 411-414.
34. El Aila NA, Tency I, Claeys G, Saerens B, De Backer E, Temmerman M, et al. Genotyping of *Streptococcus agalactiae* (group B streptococci) isolated from vaginal and rectal swabs of women at 35-37 weeks of pregnancy. *BMC infectious diseases.* 2009; 9:153.

- 299 35. Brzychczy-Wloch M, Wojkowska-Mach J, Helwich E, Heczko PB. Incidence of maternal GBS
300 colonization and neonatal GBS disease among very low birth weight Polish neonates. Medical
301 science monitor: Intern Med J Exper Clin Res. 2013; 19:34-39.
- 302 36. Makhoul IR, Sprecher H, Sawaid R, Jakobi P, Smolkin T, Sujov P, et al. Early-onset group B
303 Streptococcus sepsis in high risk neonates born after prolonged rupture of membranes. Isr Med
304 Asso J. 2009; 11(1):34-38.
- 305 37. Schuchat A, Wenger JD. Epidemiology of group B streptococcal disease. Epidemiol Rev. 1994;
306 16(2):374-402.
- 307 38. Gherardi G, Imperi M, Baldassarri L, Pataracchia M, Alfarone G, Recchia S, et al. Molecular
308 epidemiology and distribution of serotypes, surface proteins, and antibiotic resistance among
309 group B streptococci in Italy. J Clin Microbiol. 2007; 45:2909-2916.
- 310 39. Rojo P, Araya P, Martinez TMA, Hormazabal JC, Maldonado A, and Fernandez J. Molecular
311 characterization of Chilean isolates of *Streptococcus agalactiae*. Rev Med Chil.2008; 136:606-612.
- 312 40. Simões JÁ, Neder AV, Fracalanza SE, Soares de Camargo R, Mathias L, Pires Milanez M.
313 Phenotypical characteristics of group B *Streptococcus* in parturients. Braz J Infect Dis. 2007;
314 11:261-266.