

Performance of Corn Cultivars in Different Row Spacing and Types of Fertilizations

ABSTRACT

Aims: This study aimed to evaluate the productivity and production components of two cultivars of corn for different types of fertilization and row spacings on irrigated farming system.

Study design: It was adopted a randomized block design at 3 x 2 x 2 factorial experiment with four replications, the treatments consisted of three fertilization (OF - Organic Fertilization; OMF – Organomineral Fertilization and MF - Mineral Fertilization), two cultivars of corn (Bras 3010 and Potiguar) and two row spacings (80 cm and 50 cm).

Place and Duration of Study: The experiment was carried out from June to October 2013, at Fazenda Experimental Rafael Fernandes, in the community of Alagoinha, belonging to the Federal Rural University of Semi-Arido (UFERSA), lying 20 km from the city of Mossoró, Rio Grande do Norte, Brazil.

Methodology: The organic fertilization (OF) was performed as minimum recommendation corresponding to 10 t ha⁻¹ of bovine manure. The organomineral fertilization (OMF) was made by applying 50% of the recommended amount of manure recommended in organic fertilization (OF), 5 t ha⁻¹, and 50% of the recommendation of mineral fertilizer (MF). The mineral fertilization (MF) was performed based on the parameters observed in the soil analysis and recommendation for the corn crop in the region due to an expected maximum productivity.

Results: The study showed that the mineral fertilization (MF) provided an increase in the productivity components of both evaluated, the hybrid variety Bras 3010 and Potiguar. The OMF did not show significant differences regarding the MF in production components of corn cultivars evaluated.

Conclusion: Components of production and productivity of corn cultivars were incremented when using the spacing of 80 cm between rows. The variety Potiguar corn had higher grain productivity compared with the Bras 3010 hybrid when used the organic fertilization.

12
13
14
15
16
17
18
19
20
21
22
23
24

Keywords: Zea mays L.; organic fertilizer; spatial arrangement; production.

1. INTRODUCTION

Aspects such as nutritional quality, ease of adaptation and applicability, do corn crop, a worldwide important cereal, especially in the United States, China and Brazil, which are among the largest producers of this cereal with 371.0, 259.1 and 82.0 million tons in the 2017/18 harvest, respectively, representing 66.16% of world production [1]. World demand for food, feed and fuel is increasing continuously. The corn is frequently used to meet these three uses. This multipurpose characteristic grain led to a dramatic increase in cereal production demand during the last decade [2].

25 Among the crops grown in Brazil, the main cereal is corn (*Zea mays* L.) highlighting the
26 volume of production and the socio-economic importance. It is estimated that in the 2017/18
27 harvest were grown 16.63 million hectares of this culture, in which they were produced 81.35
28 million tons of grain in the country [3]. Corn also is configured as one of the important crops
29 in the state of Rio Grande do Norte, being cultivated in all 167 counties of this state, culture
30 is mainly explored by small farmers, but also by large companies of fruit growing, exploring
31 the cereal, especially, at the melon crop (*Cucumis melo* L.) off season.
32

33 The corn productivity is dependent on the technological level and the interaction between
34 genetic factors, environmental and crop management [4]. It is considered that soil fertility is
35 one of the main factors responsible for the low productivity of areas intended to produce corn
36 grains. This is due not only to low levels of nutrients present in the soil, but also to the
37 inappropriate use of liming and fertilization [5]. Several studies have shown the positive
38 effect of mineral fertilizer in the production components for the corn crop, as the number of
39 grains per ear, mass of ear with and without husk and grain yield by crop [6,7,8].
40

41 In order to increase the efficiency in the use of fertilizers in agricultural systems while
42 maintaining the ecological balance, interest in the use of biofertilizers or organic fertilizers
43 has recently increased, associated with the use of mineral fertilizers. [9] found that continued
44 use of organomineral fertilizer in the corn crop, for several year, caused significant increases
45 in grain production, which was more efficient fertilization that single application of organic or
46 mineral fertilizer. A practice commonly adopted to increase production is the use of manure
47 as an organic fertilizer for the supply of Nitrogen and Phosphorus in soils [10]. Information
48 available in the literature regarding the combined use of organic fertilization with mineral
49 fertilizers are scarce. Working in an Ultisol [11] found that the association of organic compost
50 and mineral fertilizer did not influence significantly the productivity of corn crop. The
51 productive potential of corn crop can be exploited by the judicious implementation of
52 technical aspects such as the choice of cultivar best suited to growing conditions, spacing
53 and proper management.
54

55 One of the practical ways for the family farmer adopts a new technology is the introduction of
56 new cultivars. Thus, research works have been developed trying to evaluate and validate
57 new corn cultivars under different soil and climatic conditions in different regions [12,13]. [14]
58 in studies in the Northeast Nebraska, reported that reducing the spacing between the crop
59 rows can favor an increase in grain production, because smaller spacing in the row result in
60 a distribution more equidistant between plants, increasing the area leaf, trapping solar
61 radiation and shading the soil, leading to a reduction in water loss by evaporation, resulting
62 in a higher photosynthetic capacity and consequently in a higher grain yield.
63

64 In Guarapuava, Brazil, [15] did not obtain difference in the ear lengths when compared
65 spacings of 40 and 80 cm in a population of 50,000 plants ha⁻¹. [16] evaluating the
66 performance of three corn hybrids, AS 1570, AS 1565 and AS 1575, at three locations in
67 western Paraná, Brazil, did not observe significant changing in the number of grains per ear,
68 in three different spacing between rows (45, 68 and 90 cm). [17] found that row spacing did
69 not affect the number of grain lines per ear, evaluating five corn hybrids early and super
70 early maturing, when the row spacing was reduced from 90 to 45 cm.
71

72 With the growing need for an increase of the agricultural productivity in Brazil, it becomes
73 paramount the scientific advancement in the supply studies of nutritional deficiencies and
74 management deficiencies in corn, acting increasingly as relevant factor for science and
75 agriculture. In this scenario, the aim of this work was to evaluate the production components
76 and the productivity of two corn cultivars in function of different types of fertilization and row
77 spacing on irrigated farming system in the western of Rio Grande do Norte state.

78

79 2. MATERIAL AND METHODS

80

81 The experiment was carried out from June to October 2013, at Fazenda Experimental Rafael
82 Fernandes, in the community of Alagoinha, belonging to the Universidade Federal Rural do
83 Semi-Árido (UFERSA). Located at Latitude 5°03'37" S e Longitude 37°23'50" W, with an
84 average altitude of 72 meters and slope between 0 and 2%, lying 20 km from the city of
85 Mossoró, Brazil. The city of Mossoró is in the Rio Grande do Norte state Northwest region.
86 According to W. Koeppen climate classification, the climate is BSw^h type, dry climate, very
87 hot and rainy season in the summer lingering for fall, with an average annual temperature of
88 27,4°C, annual rainfall very irregular, averaging 673.9 mm and relative humidity 68.9% [18].

89

90 The soil of the experimental area was labeled according to the brazilian soil classification as
91 ARGISSOLO VERMELHO-AMARELO eutrófico latossólico de textura franco-arenosa [19],
92 belonging predominantly to the Ultisol Order by U.S. Soil Taxonomy [20]. The chemical
93 analysis of the soil, carried out before trial installation at 0-20 cm depth, showed the
94 following results: pH 4.8; 0.14 g kg⁻¹ N; 4.19 g kg⁻¹ of organic matter; 8.1 mg dm⁻³ P; 40.1 mg
95 dm⁻³ K; 7.6 mg dm⁻³ Na; 0.52 cmolc dm⁻³ Ca; 0.44 cmolc dm⁻³ Mg and 0.15 cmolc dm⁻³ Al.

96 The area had small shrub native vegetation by the year 2010, which was subsequently
97 removed, in 2011 was barred, scarified and cultivated with bean in conventional farming
98 system. In 2012 the area was set aside. In soil preparation was done plowing, followed by
99 leveling grading. Being held liming the soil of the experimental area, raising the pH to a
100 suitable range to the nutritional requirements of corn: 5.5 to 6.5. Was distributed 2.5 t ha⁻¹ of
101 lime, with 12% MgO, broadcasted 60 days before sowing, being distributed at a depth of 0-
102 10 cm, with the aid of a grading. Irrigation was twice a week made for 60 days period to
103 complete the product reaction with the soil mineral particles.

104

105 The experimental design was randomized blocks in factorial 3 x 2 x 2, composed of three
106 types of fertilizers, (OF - Organic Fertilization; OMF – Organomineral Fertilization and MF -
107 Mineral Fertilization), two row spacings, E1 (80 cm) and E2 (50 cm), and two cultivars of
108 corn, hybrid Bras 3010 and the cultivar Potiguar, with four repetitions, totaling 48
109 experimental units of 4 x 30 m each. The organic fertilization (OF) was performed as
110 minimum recommendation of [21], corresponding to 10 t ha⁻¹ of bovine manure. The material
111 was collected in the cattle sector from the Federal Rural University of Semi-Árido, which
112 material was chemically analyzed and obtained the following characteristics: pH 7.7; 10.22 g
113 kg⁻¹ N; 34.68 g kg⁻¹ of organic matter; 806.7 mg dm⁻³ P; 5178.5 mg dm⁻³ K; 1887.4 mg dm⁻³
114 Na; 9.6 cmolc dm⁻³ Ca; 8.3 cmolc dm⁻³ Mg and 0.44 cmolc dm⁻³ Al. The organomineral
115 fertilization (OMF) was made by applying 50% of the recommended amount of manure
116 recommended in organic fertilization (OF), 5 t ha⁻¹, and 50% of the recommendation of
117 mineral fertilizer (MF). The mineral fertilization (MF) was performed based on the parameters
118 observed in the soil analysis and recommendation for the corn crop in the region due to an
119 expected maximum productivity, 15 kg ha⁻¹ of Nitrogen, being applied in the foundation, and
120 60 kg ha⁻¹ in coverage fertilization. It was applied to 80 kg of phosphorus and 50 kg ha⁻¹ of
121 potassium in the foundation. Before sowing operation, the seeds were treated with
122 imidacloprid insecticide active principle and thiodicarb at a dose of 0.35 L ha⁻¹.

123

124 With emerging percentage values and purity of each batch of seeds used in the experiment,
125 the seeder was set to distribute 4.18 and 3.46 seeds per meter spacing for 80 and 50 cm,
126 respectively. The expected values were 69,200 and 52,250 seeds per hectare for a desired
127 population of 50,000 plants per hectare. In the experiment was used a precision seeder,
128 Marchesan brand T2SI model chassis 2,800 mm, weight 656 kg and required power of 60
129 HP operating at an average speed of about 5 km h⁻¹, adjusted to 80 and 50 cm between
130 rows, respectively. The irrigation water available at the Experimental Farm came from a well

131 at Sandstone aquifer, characterized by presenting approximate depth of 1000 m, with good
 132 quality electrical conductivity (ECw) of 0.58 dS m⁻¹ and pH 7.5. The irrigation system used
 133 was by spraying, powered by a three-phase hydraulic pump Thebe brand, with capacity of
 134 7.5 hp and maximum flow of 38 m³ h⁻¹, consisting of 9 sidelines spaced 12 m, with 8
 135 sprinklers brand agropolo NY 25, each row also spaced 12 m. The spray had 25 mca
 136 working pressure of 12 m range, flow rate of 528 L h⁻¹ and height 2.5 m jet. With the
 137 meteorological station installed near the experiment was determined and applied to the
 138 amount of water necessary for each stage of culture. Irrigation was always done at night
 139 because the best application efficiency, lower drift caused by wind and consequently a better
 140 water use by the crop.

141
 142 To evaluate the production components were collected 10 ears, at random, from two central
 143 lines of the plot, targeting the following determinations: mass of ear with and without husk,
 144 ear length, ear diameter and mass of 1000 grains. The grain productivity was obtained by
 145 weighing the grain harvested in the area of the experimental plot, threshed mechanically,
 146 also summing up the grain mass of the collected ears, correcting the moisture to 13%, being
 147 held adjustment for kg ha⁻¹. The data were submitted to analysis of variance by F test at 5%
 148 probability. Then the averages were compared by Tukey test at 5% probability. In the
 149 statistical analysis was used the software SISVAR 5.0 [22].

151 3. RESULTS AND DISCUSSION

152
 153 Data from the mass of ear with husk (MEWH), mass of ear without husk (MENP), ear length
 154 (EL) ear diameter (ED), mass of 1000 grains (M1000G) and productivity (P) obtained in the
 155 experiment, as well as F values, are presented in Table 1. Analyzing the data presented, it
 156 was found that the variables: mass of ear with and without husk and ear diameter showed
 157 significant interaction for factors, cultivar and row spacing (F x C x RS), respectively.

158
 159 **Table 1. Mean and the F values of productivity and production components that**
 160 **resulted from the analysis of variance**

161


Sources of variation	MEWH (g)	MENH (g)	EL (cm)	ED (mm)	M1000G (g)	P (kg ha ⁻¹)
Fertilizations (F)						
Organic (OF)	125.28 b	107.78 b	15.00 b	39.73 b	127.01 a	1255.79 a
Organomineral (OMF)	137.23 a	118.33 a	15.32 ab	40.31 ab	128.04 a	1435.58 a
Mineral (MF)	141.32 a	122.55 a	15.69 a	40.75 a	127.00 a	1415.83 a
Cultivars (C)						
Bras 3010	119.69 b	105.04 b	15.56 a	38.06 b	124.63 b	1245.99 b
Potiguar	149.53 a	127.41 a	15.10 b	42.47 a	130.07 a	1492.14 a
Row spacing (RS)						
E1 (80 cm)	147.42 a	127.30 a	15.51 a	41.03 a	128.54 a	1379.31 a
E2 (50 cm)	121.80 b	105.13 b	15.16 b	39.49 b	126.15 a	1358.82 a
Values of F						
Fertilizations (F)	7.66 **	7.75 **	5.18 **	3.40 *	0.07 ^{ns}	1.31 ^{ns}
Cultivars (C)	73.68 **	50.23 **	6.67 *	187.55 **	4.17 *	6.10 *
Row Spacing (RS)	54.28 **	49.33 **	3.95 *	22.86 **	0.80 ^{ns}	0.04 ^{ns}

F x C	12.29 **	10.91 **	4.73 **	5.92 **	1.13 ^{ns}	4.02 *
F x RS	5.96 **	4.94 **	1.28 ^{ns}	6.31 **	0.08 ^{ns}	0.01 ^{ns}
C x RS	0.04 ^{ns}	0.00 ^{ns}	0.91 ^{ns}	0.05 ^{ns}	0.04 ^{ns}	0.34 ^{ns}
A x C x RS	5.15 **	5.41 **	1.20 ^{ns}	3.88 *	1.34 ^{ns}	0.45 ^{ns}
Average	134.60	116.22	15.33	40.26	127.35	1369.06
CV (%)	28.29	29.75	12.66	8.77	7.24	25.21

162 *Mass of ear with husk (MEWH), mass of ear without husk (MENP), ear length (EL) ear diameter (ED),*
 163 *mass of 1000 grains (M1000G) and productivity (P). Means followed by the same letter do not differ by*
 164 *Tukey test at 5% probability. *P < 0.05; **P < 0.01; ^{ns}Not significant; C.V.: Coefficient of variation.*
 165

166 Fertilization, cultivars and evaluated spacings had significant effect on the ear diameter
 167 (ED). The mineral fertilization increased the ear diameter, followed by organomineral
 168 fertilization, differing significantly from organic fertilization. The ear length and the
 169 productivity showed significant interaction only to the factors fertilization and cultivar (F x C).
 170 [23] state that the proportionate differences in the components of corn yield, including the
 171 thousand grain weight, is more due to population density than exclusively to spacing.
 172

173 The splitting of significant triple interaction between fertilization, cultivars and spacings for
 174 the mass of ear with husk are shown in Figure 1. It was found that the hybrid Bras 3010
 175 showed greater mass of ear with husk when used organomineral fertilization, use spacing of
 176 80 cm between rows, not differing significantly from the mineral fertilizer. The combination of
 177 organic fertilization with mineral fertilizer is presented as a viable alternative to the increased
 178 mass of ear with husk, it reduces at 50% of costs, as the acquisition of mineral fertilizers,
 179 providing soil improvements in physical, chemical and biological terms. Since the spacing of
 180 50 cm between rows promoted a decrease in the mass of ear with husk in the hybrid Bras
 181 3010 with the use of organic fertilizer, and this may have been due more competition for
 182 light, because there was a faster closing of corn plants in this spacing, as well as the time of
 183 decomposition of the organic matter in the soil may have been less, causing a slow nutrient
 184 availability.
 185



186
 187
 188
 189


Fig. 1. Mass of ear with husk of corn cultivars (Bras 3010 and Potiguar) submitted to organic, organomineral and mineral fertilizations in the row spacings of 80 and 50 cm

190 Means followed by the same letters, uppercase and lowercase to Bras 3010 in each type of fertilization
 191 in both row spacings, respectively, Greek letters (α) and (γ) followed by the same numbering for
 192 Potiguar in both row spacings, do not differ by Tukey test at 5% probability.

193
 194 [16] states that in tight spacing there is an increase in the production of components and
 195 consequently, increased grain yield, due to a more equidistant distribution of plants in the
 196 area, increasing the efficiency of utilization of sunlight, water and nutrients, improving weed
 197 control because of the faster closure of the spaces available, reducing erosion, improving
 198 the quality of seeding through the slower speed of rotation of the seed distribution systems,
 199 maximizing the use of seeders. The variety of Potiguar corn had a higher mass of ear with
 200 husk in relation to hybrid Bras 3010, especially when fertilized in mineral form in the spacing
 201 of 80 cm between rows (190.10 g). These results corroborate those of [24], who evaluated
 202 two spacings (90 and 45 cm) in corn, found greater results of mass of ear with husk in higher
 203 assessed spacing. The spacing of 50 cm between rows did not significantly alter the mass of
 204 ear with husk for Potiguar variety of corn in the evaluated fertilizations.

205
 206 The mass of ear without husk followed the tendency of the mass of ear with husk. The
 207 fertilization significantly alters the mass of ear without husk. The hybrid Bras 3010 showed
 208 higher mass of ear without husk when fertilized in mineral form, no significant interference of
 209 organomineral fertilization, differing only treatment with organic fertilization. The spacing of
 210 80 cm between rows positively influenced the mass of ear without husk in Bras 3010 corn,
 211 especially with mineral and organomineral fertilization, promoting higher values in the mass
 212 of ear without husk as shown in Figure 2. The spacing of 50 cm between rows significantly
 213 changed the mass of ear without husk in the hybrid tested, with higher values with
 214 organomineral and mineral fertilization, respectively.

215



216
 217

218 **Fig. 2. Mass of ear without husk of corn cultivars (Bras 3010 and Potiguar) submitted**
 219 **to organic, organomineral and mineral fertilizations in the row spacings of 80 and**
 220 **50 cm**


221 Means followed by the same letters, uppercase and lowercase to Bras 3010 in each type of fertilization
 222 in both row spacings, respectively, Greek letters (α) and (γ) followed by the same numbering for
 223 Potiguar in both row spacings, do not differ by Tukey test at 5% probability.

224

225 The splitting of significant triple interaction between fertilization, cultivars and spacing to the
 226 ear diameter is shown in Figure 3. The spacing between rows of 80 cm significantly interfere

227 in the ear diameter of hybrid evaluated, having noticed that with organomineral fertilization
 228 way, the hybrid Brasmilho 3010 presented a larger ear diameter. The results differ from
 229 those obtained by [25], which did not find significance in the ear diameters at different
 230 spacings. They could only observe a decreasing trend in the ear diameter with increased
 231 spacing, which claim to be associated with increased intra-specific population and
 232 competition. The spacing between rows of 50 cm did not cause significant changes in the
 233 ear diameter of Bras 3010 when subjected to the three types of evaluated fertilizations.
 234 According to [26] the ear diameter reflects the productive capacity of the plants, the largest
 235 diameter favors the formation of a larger amount of grains. The author found that the manure
 236 doses responsible for the best effect in diameter were 20, 40, 50 and 60 t ha⁻¹. Studying
 237 different corn hybrid under organic fertilization, [27] found the largest 44 mm ear diameter.
 238

239 The spacing of 80 cm between rows also interfered with significance in the ear diameter
 240 Potiguar corn, having noticed that when fertilized in mineral fertilization form highest values
 241 were found for the measured variable. No significant difference of the variable when fertilized
 242 with the organic and organomineral fertilization. The increase in the ear diameter with the
 243 use of mineral fertilization corroborate with [28], when says that nutrients present in mineral
 244 fertilizers play an important role in the growth and development of crops. The ear diameter of
 245 Potiguar variety was not significantly altered by fertilizations evaluated using spacing of 50
 246 cm between rows. [29] claim that the equidistant distribution of corn plants in the field
 247 improves the components production, including the ear diameter. Affirming also that the
 248 equidistant distribution between plants favors the closing of the leading, improving the
 249 interception of solar radiation and the rate of growth of corn plants in the early stages,
 250 exactly as happen with the stem diameter.
 251



252
 253

254 **Fig. 3. Ear diameter of corn cultivars (Bras 3010 and Potiguar) submitted to organic,**
 255 **organomineral and mineral fertilizations in the row spacings of 80 and 50 cm**

256 *Means followed by the same letters, uppercase and lowercase to Bras 3010 in each type of fertilization*
 257 *in both row spacings, respectively, Greek letters (α) and (γ) followed by the same numbering for*
 258 *Potiguar in both row spacings, do not differ by Tukey test at 5% probability.*
 259

260 The splitting of significant interaction between fertilization x cultivar (F x C) for the variables
 261 ear length and productivity are shown in Table 2. For variable ear length it is observed that
 262 the mineral fertilization provided higher average values, followed by organomineral, 15.69
 263 cm and 15.32 cm, respectively, differing significantly only from organic fertilization results.

264 This variable was changed significantly by the evaluated spacings where the spacing of 80
 265 cm between rows promoted higher values.

266

267 **Table 2. Splitting of interaction between the factors fertilizations and cultivars**

268

Fertilizations	EL (cm)		P (kg ha ⁻¹)	
	Bras 3010	Potiguar	Bras 3010	Potiguar
Organic (OF)	14.94 Ba	15.05 ABa	933.04 Bb	1578.55 Aa
Organomineral (OMF)	15.92 Aa	14.72 Bb	1414.81 Aa	1456.34 Aa
Mineral (MF)	15.84 Aa	15.54 Aa	1390.13 Aa	1441.53 Aa

269 *Ear length (EL) and productivity (P). Means followed by the same letter, uppercase in column and*
 270 *lowercase in row, do not differ by Tukey test at 5% probability.*

271

272 These results corroborate the [30], who found significant change in the ear length subjected
 273 to organic fertilization, starting at a dose of 20 t ha⁻¹ of manure, compared to mineral
 274 fertilization. For spacings between lines, the results differ from those obtained by [31], that
 275 when assessing three varieties of corn (Arapuim, BR 106 and AL Bandeirante) and four row
 276 spacings (40, 60, 80 and 100 cm), found that there were no significant differences in the ear
 277 lengths, just checking out differences between varieties. [15] also did not obtain differences
 278 in ear length when compared to the spacing 40 to 80 cm in a population of 50,000 plants h⁻¹.
 279 Corroborating with those obtained by [32], which found that the spacing of 90 cm between
 280 rows showed greater ear length, compared with plants spaced 45 cm, claiming that a largest
 281 spacing between rows can provide higher incidence of light on the canopy and the
 282 competition of plants by incident solar radiation, for nutrients and water, determines the
 283 formation of the ear, especially in a dense crop, which may result in a deficit of carbon and
 284 nitrogen supply to the plants.

285

286 It was found that the fertilization significantly influenced the ear length of the hybrid corn
 287 Bras 3010. The organomineral and mineral fertilizations did not differ significantly from each
 288 other, differing only from organic fertilization results. The organomineral fertilization provided
 289 higher ear length (15.92 cm), followed by mineral fertilization (15.84 cm). Since in the ear
 290 length of the variety Potiguar was not found a significant difference between mineral and
 291 organic fertilizations, differing only from organomineral fertilization. The mineral fertilization
 292 provided an increase in the ear length (15.54 cm), followed by organic fertilization (15.05
 293 cm). [33] had higher average values of ear length ranging between 18.61 and 19.51 cm,
 294 according to the doses of Nitrogen. Corroborating the results [34], determined that higher
 295 values of ear length ranging from 16.57 to 18.22 cm, depending on the row spacing and
 296 population densities. [26] obtained higher values (15.96 cm) of ear length with 20 t ha⁻¹ of
 297 manure applied and lower average values when used organic fertilization (0, 10, 30, 40, 50
 298 and 60 t ha⁻¹) and mineral ranging between 14.03 and 15.12 cm for ear length, according to
 299 fertilization.

300

301 Fertilization exerted significant interference in the productivity of Bras 3010 hybrid.
 302 The organomineral fertilization increased grain productivity of assessed hybrid (1414.81 kg
 303 ha⁻¹), followed by mineral fertilization (1390.13 kg ha⁻¹), differing significantly from results of
 304 organic fertilization (933.04 kg ha⁻¹). The productivity of the Potiguar variety was not affected
 305 significantly by any types of evaluated fertilization. Observing increased productivity when
 306 fertilized only organically (1578.5 kg ha⁻¹). Although these results are lower than those of
 307 [21], they are positive, with a view to reducing costs with mineral fertilizers, proving positive
 308 response from the Potiguar variety of corn with organic fertilization, becoming a strategy for
 309 improving the quality of soil and, consequently, increase productivity. There was a significant
 310 change between the hybrid Bras 3010 and the Potiguar variety of corn in relation to
 productivity. There was a significant decrease in productivity of the hybrid Bras 3010 (933.04

311 cm) when fertilized organically with no significant reductions between this hybrid and the
312 Potiguar variety with the use of organomineral and mineral fertilizations.

313
314

315 **4. CONCLUSION**

316

317 The mineral fertilization provided an increase in the components of production of the hybrid
318 Bras 3010 and variety Potiguar. The organomineral fertilization showed no significant
319 differences from the mineral fertilization in the production components of evaluated corn.
320 The components of production and productivity of corn cultivars were incremented when
321 using the spacing of 80 cm between rows. The variety Potiguar corn had higher grain
322 productivity compared with the Bras 3010 hybrid when used the organic fertilization.

323
324

325 **COMPETING INTERESTS**

326

327 Authors have declared that no competing interests exist.

328

329

330 **REFERENCES**

331

332 1. United States Department of Agriculture. 2018. Accessed 11 December 2018. Available:
333 <https://apps.fas.usda.gov/psdonline/app/index.html#/app/downloads>.

334 2. Bremner J, Frost A, Haub C, Mather M, Ringheim K, Zuehlke E. World population
335 highlights: key finding from prb's 2010 world population data sheet. *Population Reference*
336 *Bulletin* 2010; 65: 1-14.

337 3. National Supply Company. Follow up of the Brazilian grain harvest - Safra 2017/18. 2018;
338 5 (12): 1-148.

339 4. Raun WR, Solie JB, Stone ML. Independence of yield potential and crop nitrogen
340 response. *Precision Agric.* 2010; 12: 508-518.

341 5. Rabbit AM, France GE, Pitta GVE, Alves VMC, Hernani LC. Production systems.
342 *Cultivation of corn. Embrapa CNPMS.* 2007.

343 6. Aratani RG, Fernandes FM, Mello LMM. (2006). Coverage nitrogen fertilization in irrigated
344 maize crop, under no-tillage system. *Electronic Journal of Agronomy.* 2006; 9: 1-10. English.

345 7. Gonçalves Júnior AC, Trautmann RR, Marengoni NG, Ribeiro OL, Santos AL. Corn
346 productivity in response to fertilization with NPK and Zn in Eutrophic Yellow Red Argisol and
347 Eutroferic Red Latosol. *Science and Agrotechnology.* 2007; 31: 1231-1236. English.

348 8. Deparis GA, Lana MC, Frandoloso JF. Nitrogen and potassium fertilization and spreading
349 in corn crop. *Acta Scientiarum Agronomy.* 2007; 29: 517-525.

350 9. Silva AP, Silveira JPA, Santos D, Fraga VS, Silva E, Souza JM, Lima LPF, Nascimento
351 JAM. Edaphic Breathing After the Application of Biofertilizers in Organic Corn Crops.
352 *Brazilian Journal of Agroecology.* 2007; 2: 1251-1254.

353 10. Menezes RSC, Silva TO. Changes in the fertility of a Neolithic Regolith after six years of
354 organic fertilization. *Brazilian Journal of Agricultural and Environmental Engineering.* 2008;
355 12: 251-257.

356 11. Gomes JA, Scapim CA, Braccini AL, Filho PSV, Sagrilo E, Mora F. Organic and mineral
357 fertilization, maize yield and physical and chemical characteristics of a yellow red Argisol.
358 *Acta Scientiarum Agronomy.* 2005; 27: 521-529.

359 12. Grigulo ASM, Azevedo VH.; Krause, W.; Azevedo, PH. Evaluation of performance of
360 corn genotypes for in natura consumption in Tangará da Serra, MT, Brazil. *Bioscience*
361 *Journal.* 2011; 27: 603-608.

- 362 13. Santos JF, Graniteiro JIT, Brito LMP. Behavior of maize cultivars under the
363 edaphoclimatic conditions of the Paraíba swamp. *Environmental engineering*. 2011; 8: 81-
364 90.
- 365 14. Shapiro CA, Wortmann CS. Corn Response to Nitrogen rate, Row Spacing, and Plant
366 Density in Eastern Nebraska. *Agronomy Journal*. 2006; 98: 529-535.
- 367 15. Turkish GMS. Production and physical composition of corn plant for silage, cultivated at
368 two levels of fertilization, two spacings between rows and two planting densities. *UECO /*
369 *PR*. 2011; 65.
- 370 16. Gonçalves ML. Agronomic performance of maize hybrids as a function of population
371 spacing and densities at three sites. *UEOP. Marechal Cândido Rondon*. 2008; 68.
- 372 17. Kappes C, Andrade JAC, Arf O, Oliveira AC, Arf MV, Ferreira JP. Arrangement of plants
373 for different corn hybrids. *Tropical Agriculture Research*. 2011; 41: 348-359. English.
- 374 18. Pereira VC, Espinola Sobrinho J, Oliveira AD, Melo TK, Vieira RYM. Influence of El Nino
375 and La Nina events on rainfall in Mossoró-RN. *Encyclopedia Biosphere*. 2011; 7: 1-13.
376 English.
- 377 19. Brazilian Agricultural Research Corporation. Brazilian system of soil classification. 3rd
378 ed. Rio de Janeiro: Embrapa Solos. 2013; 353.
- 379 20. United States Department of Agriculture. Soil Taxonomy. *Agricultural Handbook No.436*.
380 2nd ed. 1999; 863.
- 381 21. RN's Agricultural Research Company. Technical recommendations for corn crop.
382 Christmas. 2010; 21.
- 383 22. Ferreira DF. SISVAR: A program for analysis and teaching of statistics. *Symposium*
384 *Magazine*. 2008; 6: 36-41.
- 385 23. Modolo AJ, Carneletto R, Kolling EM, Trogello E.; Sgarbossa M. Performance of maize
386 hybrids in the Southwest of Paraná region under different spacings between rows.
387 *Agronomic Science Journal*. 2010; 41: 435-441.
- 388 24. Calonego JC, Poleto LC, Domingue SFN, Tiritan CS. Corn productivity and growth in
389 different plant arrangements. *Agrarian Magazine*. 2011; 4: 84-90.
- 390 25. Carvalho HWL, Cardoso MJ, Leal MLS, Santos MX, Tabosa JN, Souza, MS. Adaptability
391 and stability of maize cultivars in Northeast Brazil. *Pesquisa Agropecuária Brasileira*. 2005;
392 40: 471-477. English.
- 393 26. Mata JF, Silva JC, Ribeiro JF, Afféri FS, Vieira LM. Production of hybrid corn under
394 doses of bovine manure. *Applied Research and Agrotechnology*. 2010; 3 (3): 125-133.
395 Portuguese.
- 396 27. Santos IC, Miranda GV, Melo AV, Mattos RN, Oliveira LR, Lima JS, Galvão JAC.
397 Behavior of organically grown corn cultivars and correlations between traits of harvested
398 ears in the green stage. *Brazilian Journal of Corn and Sorghum*. 2005; 4: 45-53. English.
- 399 28. NHR cyanide. Grain yield, dry matter and nutrient accumulation in crops submitted to
400 organic and mineral fertilization. *UFMS / RS*. 2010; 86.
- 401 29. Nummer Filho I, Hentschke CW. Reduction of line spacing in maize crop. *Direct Plant*
402 *Review*. 2006; 92. English.
- 403 30. Reina E, Afféri FS, Carvalho EV, Dotto MA, Peluzio JM. Effect of bovine manure doses
404 on the sowing line on corn yield. *Green Magazine on Agroecology and Sustainable*
405 *Development*. 2010; 5: 158-164. English.
- 406 31. Porto APF, Vasconcelos RC, Viana AES, Almeida MRS. Varieties of corn at different
407 spacings in the Plateau of Vitória da Conquista-BA. *Brazilian Journal of Agricultural*
408 *Sciences*. 2011; 6: 208-214.
- 409 32. Gilo EG, Silva Junior CA, Torres FE, Nascimento ES, Lourenção AS. Behavior of maize
410 hybrids in the cerrado south-mato-grossense, under different spacings between lines.
411 *Bioscience Journal*. 2011; 27: 908-914.
- 412 33. Carmo MS, Cruz SCS, Souza EJ, Campos LFC, Machado CG. Nitrogen doses and
413 sources in the development and productivity of sweet corn (*Zea mays*convar, *saccharatavar*.
414 *Rugosa*). *Bioscience Journal*. 2012; 28: 223-231.

415 34. Stacciarini TCV, Castro PHC, Borges MA, Guerin HF, Moraes PAC, Gotardo M.
416 Evaluation of agronomic traits of maize crop by reducing line spacing and increasing
417 population density. Ceres Journal. 2010; 57 (4): 516-519. Portuguese.
418
419
420
421

422

UNDER PEER REVIEW