

Coverage and use of the soil of the hydrographic sub-bowl of Rio Espinharas, PB / RN / PE with emphasis in areas of permanent preservation (APP) and areas of restricted use (AUR)

ABSTRACT

Introduction: The changes made in the natural dynamics cause risks that influence the equilibrium, terrestrial and atmospheric, which often ends in the disappearance of species, either by agricultural activities or livestock, in addition to disproportionate human occupation.

Aims: Objective of this work was to characterize the coverage and use of the Sub-Basin Hydrographic Basin (SBH) of the Espinharas river, with emphasis on Areas of Permanent Preservation (APP) and Areas of Restricted Use (AUR).

Methodology: The SBH rio Espinharas is part of the Northern Country Depression, it has one of the most typical landscapes of the northeastern semi-arid region. For the analyzes, multispectral images of the Landsat 8 OLI satellites were used, from the orbits and points 215/65, 216/64, bands 3, 4 and 5. The delineation of the sub-basin began with obtaining the hydrological attributes in the Software QGIS. For the identification of the areas of land use conflicts in APP and AUR, the map algebra was used to perform an overlay of the land cover and use map with the Map of the APP and AUR, using SIG Idrisi Software.

Results and discussion: The classes of land use and land cover in the SBH of the Espinharas River has the predominance of the Open Arboreal Arboreal Caatinga (CAAA) typologies with 2,239.37 km² (68.13%), Caatinga Arbustiva Arborea Closed (CAAF) with 203.17 km² (6.18%) of the total SBH area. It was also verified that 752.67 km² (22.90%) of the total area corresponds to anthropism. The satellite images allowed to have a clear, comprehensive and current view of the use and land cover of SBH of the river Espinharas.

Conclusion: Discrimination, mapping and quantification of land use and land cover areas through the Geographical Information System (IDRISI, QGIS GRASS) classification allowed us to obtain results with greater agility regarding the integration and manipulation of the areas. The data obtained will help recovery plans and planning of the area, since a part of SBH is not complying with the current environmental legislation.

Keywords: *Anthropism, Semi-arid, Riparian forest, environmental degradation.*

1. INTRODUCTION

The changes made in the natural dynamics cause risks that influence the equilibrium, terrestrial and atmospheric, which often ends in the disappearance of species, either by agricultural activities or livestock, in addition to disproportionate human occupation. These facts bring to light the increasingly rapid need for studies aimed at changes and landscape composition, with greater emphasis on land use and coverage in watersheds [1].

From the last decade of the last century, the understanding that it is necessary to combine sustainability with development requires new positions in relation to such farms, currently represented by Federal Law No. 9,433 of January 8, 1997, [2] better known as - "Water Law" establishing the National Water Resources Policy and Creates the National System of Water Resources Management (SINGERH) and Law 12.651 of May 25, 2012, which provides for the preservation of native vegetation and determines the presence of Areas of Permanent Preservation (APP) and Restricted Use (AUR).

Another factor that corroborates is that the preservation of the vegetal cover is a fundamental condition for the conservation of the water resources. The removal of the same causes a series of modifications in the physical environment and in the water cycle, because the dynamics and behavior of the vegetation directly affect the water regime, both in a beneficial way, by its maintenance and circulation, and by making it unavailable on the planet [3].

Silva et al. [4] reinforce that their removal discharacterizes the original environments as well as, interferes in the water balance of the BH, compromising the water supply and the sustainability of the most varied life forms, notably in the northeastern semi-arid region.

The integrated planning of BH is one of the main management techniques of a given territorial unit with regard to the socioeconomic-environmental aspect. For this, indicators should be used to systematically reduce socio-environmental conflicts, to perform actions of recovery, preservation, conservation and management of natural ecosystems, considering as essential point the quality of life of society [5].

However, when studying BH, it becomes increasingly necessary to analyze and characterize the Permanent Preservation Areas (APP) and AUR. For Boin [6], the quantity and quality of water resources is influenced by the conflicts between use and occupation of these areas, in which the importance of compliance with legislation is highlighted.

For such management, tools in the area of geotechnology allow an integrated analysis of the environment in order to understand how issues related to environmental changes behave in space. This is one of the strengths, allowing the environment to be studied in parts and understood as a whole [7].

Remote sensing, together with geographic information systems (GIS), are highly efficient tools for surveying, mapping and monitoring natural resources. Through satellite imagery it is possible to have a broad view of the study area, to have frequent monitoring of the changes that have occurred in the region over time, in an economically viable way [8].

Therefore, the objective of this work was to characterize the soil cover and use of SBH of the Espinharas river, with emphasis on APP and AUR.

2. MATERIAL AND METHODS

2.1 Characterization of the study area

The SBH rio Espinharas is inserted in the Northern Sertanea Depression, where it has one of the most typical landscapes of the northeastern semi-arid that are, the extensive plain, predominantly soft-wavy relief, residual elevations (inselbergs) [9]


It is inserted more specifically, in the intermediate regions of Patos and Campina Grande (PB), Caicó (RN), Serra Talhada (PE) [10], (Figure 01).

Figure 01 - SBH location map of the Espinharas river, semi-arid northeast, Brazil.

According to Alvares et al. [11] in the SBH area studied, climates such as Bsh and Aw 'are characterized. The Bsh type is defined as a hot and dry climate, with summer rains and with annual rainfall around 500 mm and an annual average temperature of 26 ° C; the Aw 'type is present in the western center portion of the SBH, presenting warm and semi-humid conditions with summer-fall rains, with a rainfall average of around 500 mm and an average annual temperature of 27 ° C, and extends through the southeast portion of the sub- basin [12,13].

Soils are generally shallow, stony, of crystalline origin and very vulnerable to erosion, with predominance of the following types: Luvisol chrome and Litho Neosol [14].

The vegetation present in the study area is composed of small woody species, endowed with spines and usually deciduous leaves that lose their leaves in the dry period, with a marked



presence of cactáceas and bromeliáceas [4].

According to SUDEMA [15], the Open Arboreal Arboreal Caatinga (CAAA) is present in most of the studied area, characterized by sparse vegetation with some arboreal individuals with a mean height of 3m, with herbaceous and cactaceous vegetation, being high degree of degradation in the flat relief areas.

The vegetation is classified as Closed Arboreal Shrub Caatinga (CAAF) and is found on the slopes of hills and mountains [16]. This vegetation has as characteristics the predominance of arboreal individuals.

2.2 Materials Used

- Planialtimetric Letters from SUDENE, edited in 1985 and scanned in 1996; (SB.24 - Z - A - VI), Serra Negra do Norte - RN (SB.24 - Z - B - IV), Piancó - PB (SB.24 - Z - C - III) and Ducks-PB (SB.24-Z-D-I).
- SRTM MDE covering scenes 07_w038_1arc_v3.tif.aux; s08_w038_1arc_v3.tif.aux
- QGIS softwares plus add-ons and GRASSGIS and IDRISI.
- Landsat 8 satellite imagery, OLI sensor, resolution 30m, bands 3,4 and 5.
- VANT (Drone Phantom 3, brand DJI).

For the development of the proposed work, it was important a bibliographic review to deepen and contextualize some concepts such as BH, bioma caatinga, remote sensing, geoprocessing, land use map.

Table 01 - Identification of orbital images with coverage for the study area.

Satellite	Sensor	Orbit / Point	Date
LANDSAT 8	OLI	216/064	06/08/2017
LANDSAT 8	OLI	215/065	15/08/2017

The second step consisted in selecting the software to be used in the data processing, a limitation is the cost involved in the acquisition of the data, however an alternative form is the use of open source programs, such as the QGIS and GRASS GIS, indicated for data processing, analysis and visualization, and used in this work, making this type of operation, which is more expensive, a lot more economical. The model was used to extract the morphometric characteristics in a GIS environment with the aid of the QGIS and GRASS tools.




Figure 02. Flowchart of the methodological steps.

2.3 Methods applied

2.3.1 MDE SRTM Processing

The SRTM MDE was acquired in GeoTIFF format from the USGS Earth Explore site containing the data corresponding to scenes 07_w038_1arc_v3.tif.aux; s08_w038_1arc_v3.tif.aux with a resolution of 1 arc of a second, which corresponds to approximately 30 meters, referenced in the WGS84 Datum.

The model was used to extract the APP from the water courses and from the top of the hill with the help of the QGIS tools and complements, and the processing of the data contained in the MDE comprised the following steps:

- STRM mosaic composition (Raster> miscellaneous> mosaic);
- Reprojection of the mosaic for flat coordinates, referenced to the Datum Sirgas2000, Zone 24 S (Raster> Projections> Redesign);
- Clipping involving the study area (Vector> Geoprocessing tool> crop);
- Filling of regions without data in the SRTM MDE using the "r.fillnulls" module, which operates with the Spline Adjustment Algorithm [17], implementing in SIG GRASS.

2.3.2 Processing orbital images

Multispectral images of the LANDSAT 8 OLI satellites, orbit and points 215/65, 216/64 were used. The dates of the satellite images were selected corresponding to the dry period, and presented lower cloud interference in order to provide a better evaluation of the soil use and cover, through the contrast between vegetation and soil [18].

According to Silva [18], to perform the georeferencing of orbital images, the SIRGAS 2000 horizontal Datum and the UTM Projection System are used, using control points in the field (PC), based on intersections between roads, roads and paths, confluences of rivers and other reliable and recognizable mooring points, both in orbital images and images of Google Earth 6.0.2.2014. In this way the images will be corrected geometrically applying the resampling by the method of the nearest neighbor.

2.3.3 Delimitation of the study area.

The SBH delimitation was started with the hydrological attributes obtained in the QGIS Software, in which they were executed by the GRASS complement "r.watershed" [19]. This module derives maps of flow accumulation, drainage direction, drainage location and SBH boundary.

2.3.4 Characterization of the cover and land use of SBH of the Espinharas river

After the pre-processing of the images, visual and supervised classifications were performed. For the visual method, the first step was the image processing, where it consisted of the mosaic> reprojection for the Datum SIRGAS 2000, Zone 24Sul> RGB color composition in the bands 3, 4 and 5.

183 The second stage consisted of the vector representation of each identified theme, rasterizing
184 on a mask previously generated with definition of the polygon of the basin.
185

186 The subjects chosen for use map were based on field sampling, and three samples were
187 selected previously for the thematic class. For automatic classification, the likelihood method
188 (Maxlike) was used. Ten samples of each class were verified, considering the training based
189 on the labeling formulated in the visual interpretation of the image and related knowledge of
190 the study area. From the overlapping of the two classifications, a hybrid image was
191 generated with which the land use and cover map was created with the following typologies:
192 Open Arboreal Shrub Caatinga (CAAA) - with predominance of grasses, tree and sparse
193 trees and Caatinga Shrub Arborea Closed (CAAF) with the presence of shrubs and trees
194 with height varying from 6 to 8 m [15], Area antropizada Urban Area, Corpos D'água and
195 Rocky Outcrop. Then, the area values of each category of land use were calculated. After
196 the classified image, the Kappa statistic was used to evaluate the agreement between the
197 results observed and those classified in a contingency table (error matrix). According to
198 Landis; Koch [20], Kappa values are equivalent to classification quality (Table 01). During
199 the field work, the "terrestrial truth" was verified, in which areas with possible classification
200 errors were analyzed.
201

202 **Table 1 - Quality of the classification associated with Kappa index values.**
203

Kappa Index	Quality
≤0,00	Poor
0,01 a 0,20	Bad
0,21 a 0,40	Reasonable
0,41 a 0,60	Good
0,61 a 0,80	Very Good
0,81 a 1,00	Great

204 Source – Landis; Koch (1977).
205

206 **2.3.5 Delimitation of the APP and AUR of SBH of the Espinharas river** 207

208 For the generation of APP at the drain banks, perennial and intermittent drains were
209 considered, according to current legislation, with well defined channels [21], from the fifth
210 and fourth orders in the drain hierarchy. In sequence, the marginal bands were generated in
211 the QGIS buffer module according to the average width measurement of the regular river
212 bed.
213

214 For the areas located at the top of elevations (top of hills, mounts mountains or saws) with
215 slope higher than 25 ° of elevation or (area equal or greater than 100 meters), are
216 considered (AUR), and the MDE was used SRTM, following the following steps:
217

- 218 • Analysis of the MDE (Raster> Analysis> MDE Land Models), with this was obtained
219 the slope of the terrain in degrees;
- 220 • For area delimitation (Raster Menu> Convert> Raster to vector> Polygonize).
221

222 **2.3.6 Verification of the conflict between land use and land cover classes x APP / AUR** 223

224 In order to identify the areas of land use conflicts in APP and AUR, map algebra was used to
225 perform an overlapping of the land cover and use map with the Map of APP and AUR. The
226 procedures were performed in the IDRISI SIG Software, through the "CROSSTAB" module.

After the overlapping of these maps, the areas were duly quantified and characterized as to the use and coverage of the soil taking into account the current environmental legislation, executing the area calculation functions, by the tool "Database Query" menu area, belonging to the module IDRISI Analysis.

3. RESULTS AND DISCUSSION

The validation of the land use classification (Maxlike) presented a Kappa Index of 0.8852 (88.52%), considered in the range of excellent quality according to the classification used.

As can be observed, the classes of land use and land cover in the SBH of the Espinharas River shows the predominance of the Open Arboreal Shrub Caatinga (CAAA) typologies with 2,239.37 km², representing (68.13%) of the total area, characterized by a sparse vegetation with some arboreal individuals with a mean height of 3 m, with cactaceous and herbaceous vegetation being found, in most cases, with a high degree of degradation, located in the most flat areas and also in areas with strong slopes (Table 02).

Table 02. Land use and cover and their respective areas and percentage in relation to SBH area of Espinharas, Paraíba, Brazil.

Usage and Coverage	Area (km ²)	Area (%)
CAAA*	2239,37	68,13
Anthropized area	752,67	22,90
CAAF*	203,17	6,18
Urban area	44,63	1,36
Bodies of Water	25,05	0,76
Rock outcrop	22,06	0,67
Total	3286,95	100

* CAAA – Open Arboreal Shrub Caatinga

* CAAF – Closed Arboreal Shrub Caatinga

Another typology with the presence of vegetation is the Closed Arboreal Arboreal Caatinga (CAAF) with 203.67 km² (6.18%) of the total SBH area. In this typology, trees and shrubs occur more frequently in areas of higher slope, such as slopes of hills and mountains, where dense vegetation is present, with less herbaceous and cacti (Figure 03).




Figure 03. Use and land cover map of SBH of the Espinharas river.

It is known that the vegetation cover is important in the control of erosion, floods and in the recharge of the water table. The use of these areas, for the removal of wood and for extensive livestock activity, was verified "in loco", contributing to a greater degradation of the same ones.

In other works carried out in the northeastern semi-arid region, the predominance of vegetation was found occupying most of the BH's area, such as Andrade; Oliveira [22], Mendonça et al [23], Silva [24], Souza (25), Assis et al [26], Marcelino [27], Silva et al [4] and Assis [28].

It was also verified that 752.67 km² (22.90%) of the total area correspond to the anthropized area being the second largest typology found. This topology was classified between pasture (native and planted), subsistence agriculture where maize and beans predominate and mineral extraction. It is located to a large extent on the banks of the rivers and extends to the headwaters of the water courses that feed the SBH where they can contribute to the increase of the degradation of the area, mainly by the erosion, silting and pollution of the surface waters.

Different results were found by Mendonça et al., [23] when studying SBH of Rio Jatobá, Patos-PB, inserted in the study area, in which it was identified that, in this section, 41.4% of SBH area were occupied by anthropic area; 29.7% for CAAA, 23.2% for CAAF and 5.7% for Water bodies. This different result for anthropic area occurred due to the SBH of the Jatobá dam is located in an area closer to the urban area of the municipality of Patos-PB, resulting in greater anthropic action and pressure by natural resources.

In the semi-arid, most soils are fertile, but shallow [4]. This peculiarity requires the man of the field to use soil conservation practices, eliminating or reducing the risk of erosion and

transporting the thin layer of arable land, which normally does not occur. To that end, it is important to provide technical assistance, which must be prepared to face the low level of schooling of the rural producer and the few financial resources to invest in a sustainable and economically viable agriculture Mendonça et al. [23].

The typology of water bodies that presented an area of 25.5 km² (0.76%) of the total area of the SBH where it is represented by dams, dams and barriers.

In the sub-basin studied, there are few reservoirs of greater representativity, the best known being the Jatobá Dam with a capacity of 17,516,000 m³, Flour Açude with a capacity of 25,738,500 m³, both located in the municipality of Patos-PB and Capoeira Water is the most representative with 53,450,000 m³ located in the municipality of Santa Terezinha-PB [12] The rest of the water bodies are smaller, categorized as barriers and small dams, not constituting reservoirs that can store water for periods of drought, a fact that leaves the population depending on cars kites or even cacimbas and wells built emergentially.

The other typologies were Urban Area with 44.63 km² representing (1.36%) and finally Rocky outcrops with 22.06 km² representing (0.67%) of the total area of the SBH of the river Espinharas. The SBH drainage area of the Espinharas River extends through thirty-one (31) municipalities, twenty (25) in the State of Paraíba, three (03) of the State of Rio Grande do Norte and three (03) of the State of Pernambuco. The most representative municipalities in the São José de Espinharas and Patos-PB study area. The immediate geographic region of Patos is composed of nine municipalities, and presents the highest population index of the region, becoming one of the factors, together with its privileged geographic position, relevant to the strengthening of its centrality [29].

The outcrops are part of the most typical landscapes of the northeastern semi-arid region, being inserted in the Northern Sertaneja depression, with an extensive pediplanada plain, with altitudes varying from 250 m to 700 m. Some of these rocks are granitic in nature and are quite exploited for use in construction. Satellite imagery has provided a clear, comprehensive and current view of land use. Discrimination, mapping and quantification of land use areas through classification by the Geographic Information System (IDRISI, QGIS GRASS) allowed results to be obtained with greater agility regarding the integration and manipulation of the áreas.

3.2 Mapping of APP and AUR

From the current forest legislation and the aid of the geoprocessing, as described in the methodology, the map of APP and AUR was obtained (Figure 04).

It is observed that the AUR occupy a larger area with 105.64 km² (3.21%) followed by Drainage APPs with 82.36 km² (2.51%) APP of Water bodies, 56.45 km² (1, 72%) (Table 03). APPs are considered as protected areas, covered by native or exotic vegetation, with the environmental function of preserving water resources, landscape, geological stability and biodiversity, as well as facilitating the gene flow of fauna and flora, protecting the soil and ensuring the well-being of human populations [30].




Figure 04. Map of APP and AUR of SBH of the Espinharas river.

Table 03 APP and AUR and their respective areas and percentage in relation to the SBH area of the Espinharas river, Paraíba, Brazil.

APP's	Area (km ²)	% in relation to the sub-basin area
Areas of Restricted Use	105,64	3,21
Drain APP	82,36	2,51
Water Body APP	56,45	1,72
Total	244,45	7,44

In the SBH of the river Espinharas, anthropic activities have caused impacts to the natural environment, mainly by the removal of vegetation, such as silting, erosion of drains due to lack of riparian forest.

In a similar study, Melo et al. [31] observed that plant formation was suppressed due to anthropic activities such as farming and agriculture, where the lack of riparian forests caused silting of the surrounding rivers.

Another issue that deserves attention is the obligation of the APPs to be covered by native or exotic vegetation, since, according to Garcia et al. [32], these areas have the function of reducing the transportation of material to the watercourses, silting up its banks, minimizing erosion processes and, finally, helping to maintain and preserve biodiversity.

Silva et al.[33] In a study in the Ribeirão Ubá-MG Hydrographic Basin, found a restricted use area (AUR) of 3.9%, while the Permanent Preservation area was 0.037%.

In the recreational catchment area of Alegre-ES, Silva et al. [34], the Permanent Protected area occupied 8.85% of the microbasin, and the restricted use area was 1.83%.

3.3 Conflict between land use and land cover x APP and AUR

Despite their importance and being reserved by law, APP and AUR have been the target of anthropic activities, leading to conflicting interests in land use and occupation.

Regarding compliance with environmental legislation, especially with regard to the protection of APP and AUR, it is verified that the SBH of the Espinharas river presents a reality different from what is foreseen in the legislation.

It is observed in (Table 04) that most of the permanent preservation areas present conflicting use in relation to what is established by the current environmental legislation [30], with 144.95 km² of APP and AUR areas (59, 85%) covered by Caatinga Arbustiva Aberta (CAAA) and 29.53 km² of APP and AUR areas (9.77%) are covered by Caatinga Arbustiva Arborea Closed (CAAF). On the other hand, 60.31 km² (26.73%) of APP and AUR areas are being used for anthropic activities.

Similar results were obtained by Silva et al. [35], in the Paraíba basin, where areas with aptitude for forest protection or afforestation due to marked declivity were also used in agriculture (7.62%) and pasture (6.34%).

Table 04. Conflict between Use Classes and Soil Coverage and APP and Restricted Use Area in SBH of the Espinharas river.

Classes		Area (km ²)						
APP \ Usage and Coverage	Water Body APP	Área (%)	Drain APP	Área (%)	Restricted Use Area	Área (%)	Total (km ²)	Total (%)
CAAA	38,17	67,76	40,04	48,63	66,74	63,17	144,95	59,85
Anthropized Area	16,27	28,88	35,87	43,57	8,17	7,73	60,31	26,73
CAAF	1,23	2,18	1,27	1,55	27,03	25,58	29,53	9,77
Urban area	0,38	0,68	0,96	1,17	0,37	0,35	1,72	0,73
Bodies of Water	0,25	0,45	4,08	4,95	0,10	0,10	4,44	1,83
Rock outcrop	0,03	0,05	0,11	0,14	3,23	3,06	3,37	1,08
Total	56,34	100,00	82,33	100,00	105,64	100,00	244,31	100,00

According to Medeiros [36], one of the main impacts found in the Espinharas river is the removal of the trees from the riparian forest. He also mentioned that in these places the presence of solid and liquid wastes is more and more visible, invasion and irregular occupation.

Drainage APPs that are located along the perennial and intermittent riverbanks and the Water Bodies APP that are located at the banks of dams and dams together represent an area of 138.67 km² and are 52.14 km² , 37%) of the area occupied by anthropic activities.

These are areas with soils with higher moisture content and higher natural fertility and are present in the vicinities of water reservoirs, which are widely used with little or no conservation by man from the field for subsistence farming or pasture [23].

There are also conflicts where APP, AUR are occupied with 1.72 km² of urban area (0.73%) of the total area of the SBH of the river Espinharas. It is observed that, even if the APPs have legal protection at the federal, state and municipal level to control their occupation and degradation, they are occupied by the classes of land use and land cover, occurring non-compliance with the legislation, probably due to lack and / or omission of the competent bodies [37].

The riverside population is one of the main causes of these impacts, and also the first to be harmed, since, corroborating Mendonça et al.[23], in order to achieve a reversal of these processes, it is necessary to change society's position, that of managers in a joint action of the public authorities, of the population and of the entrepreneurs, contributing to the reduction of the impacts that afflict the SBH of the river Espinharas, as well as its recovery.

Similar results were observed by Nardini et al [38], in a study carried out in the Ribeirão do Morro Grande, SP, Brazil, that 21.13% of APP and AUR areas are being used for anthropic activities such as pasture and agriculture.

In the sub-basin of the Córrego dos Bois in Minas Gerais, Silva et al. [39], verified that most pastures are pasture, corresponding to 22.52% and with perennial crops with 1.47% respectively.

In the work done by Santos et al [40], it was observed in the Piauitinga-PE basin, that the areas of APP present conflicts with urban area 4.3%, agricultural crop 19.11% and pasture 44.11%.

It is observed that most of the permanent preservation areas in watersheds present conflicts with other anthropic activities, this is related to the incorrect use of the soil and also the environmental legislations are not long. For this, more studies are needed in watershed areas to establish recovery and conservation measures and also to establish public policies that encourage communities to use sustainable natural resources.

4. CONCLUSION

The classes of soil cover and use in the SBH of the Espinharas River show the predominance of the Open Arboreal Shrub Caatinga (CAAA) typologies with 2,239.37 km² (68.13%) and anthropic area with 752.67 km² (22.90%) of the total SBH area of the Espinharas river. Another typology found is the Arboreal Closed Arboreal Caatinga (CAAF) with 203.17 km² (6.18%).

The typology of water bodies presented an area of 25.5 km² (0.76%) of the total SBH area represented by dams, dams and barriers. The other typologies found were urban area with 1.36 km² (44.63%) and rocky outcrops with 22.06 km² representing (0.67%) of the total area.

The adoption of measures and practices for soil conservation in these areas is fundamental to maintain the ecological quality of these resources in the long term. Failure to observe this

balance in the formulation of agricultural systems has been responsible for the breakdown of this balance and the continuous degradation of this resource, mainly due to the loss of soil via erosion in the growing areas.

Satellite imagery has provided a clear, comprehensive and current view of land use. Discrimination, mapping and quantification of land use areas through geographic information system classification (IDRISI, QGIS GRASS) allowed results to be obtained with greater agility regarding the integration and manipulation of the areas.

The data obtained will help in the future recovery and planning projects of the area, since a part of SBH has not been preserved and is failing to comply with the current environmental legislation.

REFERENCES

1. Andrade FM, Lourenço RW. Soil Use and Plant Cover in the Una - Ibiúna / SP River Basin. *Journal of the Department of Geography*. 2016; 32: 48-60. (in Portuguese)
2. National Policy on Water Resources. Law No. 9,433, of January 08, 1997. Available at: <http://www.planalto.gov.br/ccivil_03/leis/L9433.HTM>. Accessed on: 20 January 2019. (In Portuguese)
3. Almeida AQ, Ribeiro A, Paiva YG, Souza CAM. Relationship between forest cover and hydrological response of a river basin. In: XIV Brazilian Symposium on Remote Sensing, Natal. Anais ... Natal, 2009; 2057-2513. (in Portuguese)
4. Silva RMP, Lima JR, Mendonça IFC. Alteration of the vegetation cover in the sub-basin of the Espinharas River in the period 2000-2010. *Brazilian Journal of Agricultural and Environmental Engineering*. 2014; 18 (2): 202-209. (In Portuguese)
5. Soares LS, Lopes WGR, Castro ACL, Araujo GMC. Morphometric analysis and prioritization of river basins as an integrated environmental planning instrument. *Journal of the Department of Geography*. 2016; 31: 82-100. DOI: <http://dx.doi.org/10.11606/rdg.v31i0.107715>. (In Portuguese)
6. Boin MN. Areas of Permanent Preservation: A practical vision. *Practical Handbook of the Justice of the Environment*. 1 ed. São Paulo: Official Press Office of the State of São Paulo. 2005:2
7. Pires EVR, Silva RA, Izippato FJ, Mirandola PH. Geoprocessing Applied to Land Use and Land Use Analysis for the Purposes of Environmental Planning in the River Stream Hydrographic Basin - Três Lagoas (MS). *Revista Geonorte*. 2012; 2 (4): 1528-1538. (In Portuguese)
8. Leal JV, Todt V, Thum AB. The use of GIS for Monitoring Degraded Areas- – Case Study: PPA Arroio Gil, Triunfo / RS. *Revista Brasileira de Cartografia*. 2013;65(5):967-983. (in portuguese)
9. Velloso AL, Sampaio EVSB, Pareyn FGC. Proposed ecoregions for the catinga biome. Recife: Associação Plantas do Nordeste; Institute of Environmental Conservation The Nature Conservancy of Brazil, 2002. 76 p. (in Portuguese)

- 493 10. IBGE - Brazilian Institute of Geography and Statistics. Regional division of Brazil in
494 immediate geographic regions and intermediate geographical regions, Rio de
495 Janeiro, 2017. Available at:
496 <<https://biblioteca.ibge.gov.br/visualizacao/livros/liv100600.pdf>>. Access: 30 June
497 2018. (in portuguese)
498
499 11. Alvares CA, Stape JL, Sentelhas PC, Gonçalves JLM, Sparovek G. Köppen's
500 climate classification map for Brazil. *Meteorologische Zeitschrift*. 2014;22:711–728.
- 501 12. AESA. Executive Agency for Water Management of the State of Paraíba. State Plan
502 for Water Resources of the State of Paraíba. Final report. Hydrological year 2008-
503 2009. 2010. (In portuguese)
- 504 13. Paraíba. Secretary of State for Science and Technology and Environment -
505 SECTMA. PERH-PB: state water resources plan: executive summary & atlas / State
506 Government of Paraíba; State Secretariat for Science and Technology and the
507 Environment, SECTMA; Executive Agency for Water Management of the State of
508 Paraíba, AESA. - Brasília, DF: Consortium TC / BR - Concremat, 112p. : il. 2006. (In
509 Portuguese).
510
511 14. Santos HG, Jacomine PKT, LHC Anjos, Oliveira VA, Lumbrreras JF, Coelho MR,
512 Almeida JÁ, Cunha TJF, Oliveira JB. Brazilian Soil Classification System, 3 ed.
513 Brasília, DF: Embrapa, 353 p, 2013. (In portuguese)
- 514 15. SUDEMA - Superintendence of Administration of the Environment. Update of the
515 forest diagnosis of the State of Paraíba. João Pessoa: SUDEMA, 268p. 2004. (In
516 portuguese)
- 517 16. AESA - Executive Agency for Water Management of the State of Paraíba.
518 Interactive map. AESA, 2010a. (In portuguese)
- 519 17. Mitsova H, Mitsova L. Interpolation by regularized Spline with tension: I. Theory and
520 implementation. In: *Mathematical Geology*. 1993;25:641–655. (In portuguese)
521
- 522 18. Silva RMP. Alteration of the vegetation cover in the sub-basin of the Espinharas
523 River in the period 2000-2010: geoprocessing as a tool for environmental
524 management. 2011. Dissertation (Master's). Postgraduate Program in Forest
525 Sciences. CSTR / PPGF, Patos-PB, 143p. : il. 2011. (In portuguese)
- 526 19. Neteler M. Introduction to GRASS GIS software, Hannover, 2nd ed.1998.
- 527 20. Landis J, Koch GG. The measurements of agreement for categorical data.
528 *Biometrics*. 1977;33(3):159-179.
- 529 21. Lima WP. Forest hydrology applied to the management of hydrographic basins. 2.
530 ed. São Paulo: School of Agriculture Luiz de Queiroz, 245p, 2008. (In portugue)
- 531 22. Andrade JB, Oliveira TS. Spatial-temporal analysis of land use in part of the semi-
532 arid region of Ceará. *Brazilian Journal of Soil Science*. 2004; 28: 393-401. (In
533 portuguese)

- 534 23. Mendonça IFC, Silva JER, Souza ATA, Lopes IS, Neto PNM. Adequacy of the use
535 of the soil according to the environmental legislation in the hydrographic basin of the
536 Jatobá dam, Patos-PB. *Geography*. 2010; 19 (2). (In portuguese)
- 537 24. Silva L, Lima ERV, Almeida HÁ, Costa Filho JF. Geomorphometric characterization
538 and mapping of conflicts of use in the drainage basin of Aude Soledade. *Brazilian
539 journal of geography and physics*. 2010; 3 (2): 112-122. (In portuguese)
- 540 25. Souza ATA. Physical - conservationist diagnosis of the watershed 139 of the Jatobá
541 reservoir, Patos - PB. Patos - PB: CSTR, UFCG, Monografia (Forest Engineering) -
542 Federal University of Campina Grande / Rural Health and Technology Center. 23p.
543 2010. (In portuguese)
- 544 26. Assis FRV, Lima JR, Mendonça IFC, Silva JER, Santos HCM, Medeiros JX. Use of
545 geoprocessing in the study of soil cover in the Brazilian semi-arid region. *Scientia
546 Plena*. 2012; 8 (4): 1-6, 2012. (In portuguese)
- 547 27. Marcelino RL. Risks and vulnerabilities of the Santa Luzia basin - PB. 2012. 138 f.
548 Thesis (Doctorate in Agricultural Engineering) - Center for Science and Technology
549 602, Federal University of Campina Grande, Campina Grande, 2012. (In
550 portuguese)
- 551 28. Assis FRV. Identification of potential areas for environmental vulnerability in the
552 Talhado watershed, Santa Luzia-PB, 2015. 105 f. Dissertation (Master in Forest
553 Science) - Federal University of Campina Grande, Patos, 2015. (In portuguese)
- 554 29. Cavalcante VLU. THE CITY CENTER OF PATOS-PB: a study based on spatial
555 arrangements. Dissertation (masters in geography) - Federal University of Paraíba -
556 UFPB. João Pessoa-PB. 117p. 2008. (In portuguese)
- 557 30. BRAZIL. Current Forest Code. Law No. 12,651 of May 25, 2012a. Available at
558 <http://www.planalto.gov.br/ccivil_03/_ato2011-2014/2012/lei/l12651.htm>. Access:
559 28 January 2019
- 560 31. Melo IS, Silva DB, Santos ALA, Santana FS, SANTANA BLP. 2016. Anthropogenic
561 activities and environmental degradation in the sub-basin of the river Itamirim-SE,
562 REGNE. 2016; 2, Special Issue. (In portuguese)
- 563 32. Garcia YM, Campos S, Spadotto AJ, Campos M, Silveira GRP. Characterization of
564 land use conflicts in PPAs in the Barra Seca stream basin (Pederneiras / SP).
565 *Energy in Agriculture (UNESP, Botucatu, CD-Rom)*. 2015; 30: p. 68. (In portuguese)
- 566 33. Silva FP, Rocha HB, Marques Neto R. Land use conflicts in permanent preservation
567 areas at Ribeirão Ubá Hydrographic Basin-MG. *Journal of Geography*. 2015; 5 (2):
568 141-156. (In portuguese)
- 569 34. Silva KG, Ferrari JL, Belan LL. Environmental compliance of a microwatershed
570 based on brazilian forest code. *Revista de Ciências Agrárias*. 2016;59(1):87-92.
571 DOI: <http://dx.doi.org/10.4322/rca.2023>. (In portuguese)
- 572 35. Silva ER, Lima JR, Mendonça IFC, Assis FRV. Classification of the suitability of land
573 use in the Talhado microbasin, Santa Luzia municipality - PB, *Revista de Ciências
574 Agroambientais*. 2017; 15 (1): 96-105. (In portuguese)

- 575 36. Medeiros FS. Environmental impacts and delimitation of the permanent preservation
576 area in the Espinharas River in the urban stretch of Patos-PB. (Monograph) Federal
577 University of Campina Grande, UFCG / CSTR - Patos, 88f. : il. color. 2015. (In
578 portuguese)
- 579 37. Alves JB, Medeiros FS. Environmental Impacts and Delimitation of the Permanent
580 Preservation Area in the Espinharas River in the urban stretch of Patos-PB.
581 Networking Magazine. 2016; 21 (20): 207-130. (In portuguese)
- 582 38. Nardini RC, Campos S, Ribeiro FL, Gomes LN, Felipe AC, Campos M. Evaluation of
583 the use of conflict areas in app stream Morro Grande Watershed. Caminhos de
584 Geografia. 2015;16(55): 104-113. (In portuguese)
- 585 39. Silva MS, Bueno IT, Acerbi Júnior, FW, Borges LAC, Calegario N. Evaluation of the
586 land cover as water management indicator: a case study at Córrego dos Bois sub-
587 basin, Minas Gerais. Engenharia Sanitaria Ambiental. 2017; 22(3):445-452. DOI:
588 10.1590/S1413-41522017149673. (In portuguese)
- 589 40. Santos WA, Almeida AQ, Cruz JF, Mello AA, Santos RB, Loureiro DC. Land use
590 conflicts in permanent preservation areas of a Piauitinga river basin, Sergipe, Brazil.
591 Revista de Ciências Agrárias. 2017;60(1):19-24.
592 DOI:<http://dx.doi.org/10.4322/rca.2281>. (In portuguese)