

Evaluating Landuse Change along Thiruvananthapuram Coast, South West Coast of India Using Geo-spatial Techniques

10 **ABSTRACT**

Coastal zone across the world is typified with intensive multiple human activities. Historic maps and multi-temporal satellite images are helpful in detecting the change in landuse over the years. An attempt is made to evaluate the landuse/landcover change along the coastal zone of Thiruvananthapuram using Geographic Information System (GIS) and Remote Sensing techniques. Landuse scenarios for 1988, 2000 and 2010 have been generated using Survey of India Topographical Sheets, Landsat ETM+ Image and Quickbird Image respectively. ERDAS software is used for image processing and landuse change is analysed using ArcGis 10.2.1 Software. The reasons and implications of landuse change in the coast are also studied through field investigation and valid conclusions are drawn. Agricultural land, clustered settlement area, settlement with vegetation, sandy area and rocky area are the broad landuse categories delineated in the coastal zone. The analysis shows that agricultural land area has decreased from 9.34% to 3.89% while area under clustered settlement increased from 5.75 % to 25.65% during 1988-2010. This study can provide base information on landuse scenario of the coast while framing out ICZM plan.

12 *Keywords: Landuse change, coastal zone, GIS, Thiruvananthapuram coast*

13 **1. INTRODUCTION**

14
15
16
17 Coastal zone, the triple interface of land, water and air supports an earth's most dynamic fragile and productive ecosystems, requires pertinent sustainable strategies for the protection of coastal inhabitants and development of its resources. Landuse management plays a crucial part in the management of coastal zone, as it limits and delimits human activities and utilization of resources within the coast. Use of land for various human activities within the coast is highly conflicting due to high competition among its users and its limited availability for use. Inappropriate and uncontrolled use of land in the coastal zone lead to unscientific use of resources, congestion of human settlements, increase of pollution, and land degradation. The global climate change and rapidly rising of sea level exacerbate the already prevailing vulnerability of the coastal people and assets to hazards. Reliable knowledge on the coastal landuse, its evaluation on spatial pattern of distribution, and temporal variation are essential inputs in framing out ICZM plans.

26 Coastal zone is a common property resource. Although it covers only 6.3% of the earth's surface area, it provides about 43% of estimated value of world's ecosystem services (Costanza et al. 1997). About 23% of the global population lives within the coastal zone. The average population density in coastal zone is 112 persons/km² and is several times higher than the average global population density of 44 persons/km² (Small and Nicholls, 2003). The ever increasing anthropogenic pressure on the coasts over the last century has substantially reduced or depleted many valuable natural ecosystems in the coast. For example, area under mangrove vegetation of the world has declined from 19.8 million ha in 1980 to 14.7 million ha in 2000 (Howari et al. 2009). Area under natural coastal systems is now converting at a faster rate for human inhabitation. Urbanization and tourism development could cause an irreversible change in the coastal landuse in the future. The situation in tropical maritime countries like India is also not different. According to 2011 Census data, nearly 250 million people or 17% of Indian population live within a distance of 50kms from the coast spreading over 66 coastal districts in 9 coastal states. There are 2661 urban centres and 3827 rural villages in the coastal zone of India. The

average population density in the coastal zone of India was 455 persons/km² in 2001 which has increased to 558 persons/km² in 2011 (Senapati and Gupta, 2014). This warrants effective land use planning at various scales in the coastal zone in order to effectively manage the coastal resources.

Coastal land use change studies provided a base to coastal zone management even from a very earlier time. Historical maps, aerial photographs and satellite images and field survey are the sources of data traditionally used in land use mapping by scientific community across the globe. Land use change detection using Landsat MSS data in the Matagorda Bay estuarine system of Texas coast is one of the earliest attempts in this regard (Weismiller et al. 1977). A Land use classification scheme developed by USGS (1997) is commonly followed in western countries for their land use/land cover investigation purposes. Nayak (2001) developed a classification scheme for coastal land use mapping applicable to Indian coastal zone. A land use zonation scheme developed for Segara Anakan coast of Indonesia applying ICZM principles satisfied the needs of different resource users, preserved ecologically important areas and settled land use conflicts (Chua, 1992). Land use change can have a corresponding change in the ecosystem services provided by the coasts (Zhao et al. 2004). Climatic implications of land use change was analysed for Godavari Delta region using multi-temporal remote sensing data (Sharma et al. 2001).

Land use change predictions are now attempted in different parts of the world especially urban areas using various stochastic and cellular models (Lacono and Levinson, 2009) while few of them are in the coastal zone. Land use change in Abudhabi coastal zone was analysed using remote sensing data (Yagoub and Kolan, 2006). The coastal land use change along with its reasons was evaluated for North Western Egypt (Shalaby and Tateshi, 2007). Lo and Gunasiri (2014) analysed the impact of land use change on shoreline change dynamism in Taiwan, and established both positive and negative influence of land use change in causing erosion and accretion. Land use change mapping within the coastal regulation zone (up to 500m from High Tide Line) of Kerala was attempted for the year 1995 by Centre for Earth Science Studies (Baba et al. 1998) and for 2011 by SAC. However, a comprehensive study to detect change in land use within coastal zone of Kerala is yet to be attempted. The present study on Thiruvananthapuram coast is expected to fill this gap.

Thiruvananthapuram, the southernmost district of Kerala on the southwest coast of India supports 75km long, 5km wide coastal zone between North latitudes of 8° 17' 39.16" and 8° 47' 51.77" and East longitudes of 76° 40' 22.7" and 77° 07' 44.14" (Fig.1). The coastal zone covers a land area of about 345km². The coast is NW-SE in orientation which deviates slightly from the general NNW-SSE orientation of Kerala coast. This makes the coast unique in terms of climate, hydrodynamics, geology and land use. The geological setting of the coast comprises Archaean crystalline rocks in patches around Veli, Kovalam and Vizhinjam; and the Tertiary and Quaternary sediments in elongated lateritic and sandy coastal plains (GSI, 1986). About 47% of the area belongs to less than 10m elevation which is the Low Elevation Zone (LEZ). Well developed beaches, spits, bars, ridge-runnels, estuaries, sandy plains, sedimentary and crystalline cliffs/headlands are the important landforms present in the coast. Neyyar, Karamana and Vamanapuram the three important rivers originating from the Western Ghats pass through the coast and empties into the Lakshadweep Sea. The coast has a chain of backwaters (locally known as 'Kayal') interconnected each other through canals. Poovar Kayal Poonthura Kayal, Veli Kayal, Kadinamkulam Kayal, Anchuthengu (encompassing Moongottu Kayal and Kozhithottom Kayal) and Edava-Nadayaara Kayal are the important backwaters present in the coast.

Administratively, the coast spreads over 15 rural panchayats and 2 urban centres. Thiruvananthapuram City, the capital of Kerala and Varkala, a class-II urban centre are located within the coast. The coast is densely populated with an average of 2942 persons/ km² (2011, Census). As many as 42 fishing villages and 51 fish landing centres are located in the coast (Dept of Fisheries, 2012). Apart from fisheries, tourism significance of the coast is known across the globe. Kovalam beach, a world class tourist destination of Asia is positioned on the coast. Techno Park, a multinational IT firm, Vikram Sarabai Space Centre, and Travancore Titanium Products Ltd are also based within the premise of the coast. Agriculture, mining and quarrying, construction works, coir manufacturing, and fish processing are the other important economic activities prevalent in the coast. The proposed Vizhinjam International Seaport Ltd is under the first stage of its building in the coast and is expected to accelerate the economic development in the coast.

2. MATERIAL AND METHODS

The present study analyses land use change within Thiruvananthapuram coast using various geo-spatial materials and techniques. The extent of the coastal zone has been delineated with the help of salinity measurement data of coastal water bodies published by National Centre for Earth Science Studies (2014). The salinity measurement data showed that salinity value ≥ 5 ppt recorded up to a maximum distance of 5km from the shoreline in the coast. Hence the width of the coastal zone has been fixed as 5km from the shoreline uniformly throughout the coast. Survey of India Topographical Sheets of 1988 (1:25,000 scale), Landsat image of 2000 (30 m resolution), and Quickbird image of 2010 (0.5m resolution) are the base materials used for land use change analysis. Due to high scale variability, the land use categories delineated has limited up to the Level -I category of Coastal Land use classification scheme of Nayak (2001). However, the

nomenclature of each land use type has been assigned as per the standard scheme developed for land use mapping of Kerala coast by Centre for Earth Science Studies (1995). The Toposheets are georeferenced with its co-ordinates embedded and projected to Universal Transverse Mercator Projection 43 1/2°N and WGS 1984 Co-ordinate System using ArcGis Version 10.2.1. Both the satellite images are also georeferenced and projected using the same co-ordinate system. Six land use categories are delineated with the help of onscreen digitisation of SOI Toposheets (1988) and Quickbird image (2010). The growth and reduction in area of each land use category are demarcated in the Landsat image (2000) with the help of visual interpretation and onscreen digitization. Agricultural lands, clustered settlement area, settlement with vegetation, water body, sandy area, rocky area are the six landuse categories delineated in all three sets of data. The area under each land use types is extracted for 1988, 2000, and 2010 using ArcGis software and their change in percentage has been calculated using the following formula:

$$LUC = (A_o - A_1)/A_o \times 100\%$$

Where,

LUC = area change of each landuse type

A_o = area of Present Period

A₁ = area of Past Period

3. RESULTS AND DISCUSSION

Agricultural land, clustered settlement area, settlement with vegetation, water body, sandy area and rocky area are the broad six landuse categories delineated in the Thiruvananthapuram coast. Besides, the land has also been given for infrastructural development like airport, highway bypass, and other roads meeting the demand of local settlements and tourism. Area under these categories has not been computed separately. The area under airport was computed under clustered settlement category. Landuse change under each category during the 1988-2010 is discussed in the following paragraphs.

3.1 Agricultural lands

Agricultural land spreads over the flood plains, runnels and low valleys. Both seasonal and perennial crops are cultivated in the coastal tract. Coconut is the most important perennial crop cultivated in the coast. Availability of well drained coastal alluvial soil, salt - laden wind, high rainfall and temperature conditions support the flourishing of coconut farming in the area. Paddy, plantain and vegetables are also cultivated in the coastal tracts of Thiruvananthapuram. However, a drastic reduction in the agricultural land area is observed. The agricultural land constituted 9% of the coastal zone in 1988 (Fig. 2), 6% in 2000 and has further decreased to 4% in 2010 (Table 1). On analysing the percentage of growth rate, it is evident that maximum reduction in agricultural land (-39%) takes place during 1988 - 2000 periods. Indiscriminate and accelerated reclamation of lowlands to accommodate settlements and other built up areas after 1990's has shrunk the agricultural land in the coast. This tendency has continued in the coast and shown a growth rate of -31% for the period 2000-2010. The reduction in agricultural land area for about -58% during short 22 years time limit in the coastal zone would have a multiple adverse effect on the ecosystems and society at this era of global climate change and accelerated sea level rise. Incidents of severe water logging even for small downpours in the coast are posing serious threat to life and property in the coast. Management measures are required to conserve the remaining 13km² agricultural land as part of Coastal Zone Management Programme.

3.2 Clustered Settlement Area

Settlement patterns are not uniform in the Thiruvananthapuram coast. They were clustered and concentrated mainly in and around urban centres of Thiruvananthapuram and Varkala along with small patches at Poovar, Vizhinjam and Anchuthengu in 1988 (Fig. 2). Category of landuse characterised with built-up area of above 50% are included in this category. Area under clustered settlement covered only 6% of the total area in 1988 which has increased to 13% in 2000 (Fig. 3) and 26% by 2010 (Fig. 4). The percentage of growth corresponds to these two decades are 116% and 106% respectively. Settlement area doubled within a decade although rate of population growth is low. This trend calls for deeper probe to understand the family structure vis-a-vis dwelling requirements. Clustered settlement area has grown at the cost of agricultural lands and areas of settlement with vegetation. The inter settlement space within the clustered settlements is decreasing. The dominant concentration of clustered settlements in the central sector between Veli and Panathura is due to the phenomenon of urban sprawl (Fig. 3 and Fig. 4). This is clear from the study that the Thiruvananthapuram city has grown laterally along the coastal tract invading the nearby lands. Nearly 70 km² land area of the coastal zone is converted into built-up land over the period of 22 years indicating a positive increase of 346%. This type of landuse change comes under the irreversible category as conversion of this landuse type

back to its original state is almost impossible. Hence, strategies and plans are to be worked out to efficiently use this land for future settlement requirements and to minimize its encroachments to surrounding lands.

Table 1. Area under different landuse categories, Thiruvananthapuram coast

Sl. No.	Landuse Categories	1988		2000		2010		Rate of Change (%)		
		Area (km ²)	Area (%)	Area (km ²)	Area (%)	Area (km ²)	Area (%)	1988-2000	2000-2010	1988-2010
1	Agricultural Lands	32.23	9.34	19.35	5.66	13.31	3.89	-39.4004	-31.27	-58.35
2	Waterbody	18.13	5.25	18.13	5.30	18.11	5.29	0.952381	-0.19	0.76
3	Rocky Area	0.66	0.19	0.31	0.09	0.40	0.12	-52.6316	33.33	-36.84
4	Sandy Area	1.09	0.32	2.46	0.72	4.17	1.22	125	69.44	281.25
5	Clustered Settlement	19.83	5.75	42.53	12.43	87.76	25.65	116.1739	106.36	346.09
6	Settlement with Vegetation	273.15	79.15	259.32	75.80	218.35	63.83	-4.23247	-15.79	-19.36
Total		345.10	100.00	342.10	100.00	342.10	100.00	-	-	

3.3 Settlement with Vegetation

As in the case of Kerala coast, Thiruvananthapuram has also its major share of land for settlement with vegetation landuse category where half of the area is built-up or half falls under vegetative cover. This type of landuse is predominant in the rural areas of the coast in the southern and northern sectors where population density is comparatively low and per capita availability of land is high.

Majority of the land area belonged to settlement with vegetation (79%) in 1988, 75% in 2000 (Fig. 3) and further decreased to 64% in 2010 (Table 1). The percentage of growth rate for both the periods registers negative values of -4% and -16% respectively. The massive change has took place during 2000-2010 periods indicates that land scarcity in the urban coastal areas lead to encroachments of rural lands for settlement purposes and related land reclamations. Hereditary fragmentation of landholdings associated with population increase and increasing demands for settlements in rural areas have also contributed to the reduction of area under clustered settlements with vegetation at a faster rate. Since this category of land use is based on the tertiary sediments with varying relief in the coast, management plans are required for their sustainable use.

3.4 Water bodies

Water bodies including backwaters, lakes, estuaries and rivers together account for about 5% of the total area of the coast and maintain their status except for the period 2000-2010. Encroachment of water bodies particularly backwaters and estuaries are prevalent in the coastal zone around Akkulam and Kadinamkulam. The beds of these water bodies are mined indiscriminately for sand and gravel. This is degrading the rivers and its surrounding lands. This problem is not addressed in this study and this requires field measurements of bed characteristics of water bodies. Since coastal water bodies supports highly sensitive ecosystems like mangroves and islands, special attention has to be given in the ICZM.

3.5 Sandy Area

The area under sand comprises beaches, spits and bars. About 61% of coastline is sandy in Thiruvananthapuram and nearly half of this coastline is eroding. However, area under sand increased from 0.32% in 1988 to 0.72% in 2000 and further increased to 1.22% in 2010 (Table 1). This increasing trend is mainly due to the accretional activity, accelerated locally, due to the influence of coastal structures like groynes and breakwaters. The major increase in the sand area is noticed from South of Panathura breakwater and at Adimalathura where accretion is taking place at a faster rate. Panathura breakwater constructed during 2002-2003 has caused severe accretion in its updraft side and erosion in its downdraft side. However, strategies to conserve the new accreted sandy areas of Thiruvananthapuram coast are to be worked out as part of ICZM.

3.6 Rocky Area

Rocky area includes cliffs, rock exposure and rock outcrops. Rocky area has decreased to 0.12% in 2010 from 0.19 in 1988 showing a reduction of about 37% (Table 1). About 19% of coastline in Thiruvananthapuram coast is rocky. Rocky coastline is mainly seen in the northern and southern portions around Varkala, Vizhinjam and Kovalam. Area under rocky

188 area category shows decrease trend mainly due to the cliff slumping, erosion, mining and quarrying. Rocky coastline of
189 Thiruvananthapuram supports unique intertidal biodiversity. Management actions are also required to protect this
190 ecosystem as part of ICZM.

191 Landuse zonation and planning are two essential aspects of Integrated Coastal Zone Management. Field validation of
192 land use maps helps in analysing the impact of landuse change on environment. Landuse is essentially a footprint of
193 human use of land and land use change reflects the change of human activities within land spatially and temporally. The
194 present inventory derived shall help in framing out efficient management strategies for coastal land use planning in
195 Thiruvananthapuram Coast as part of evolving ICZM plan.




Fig. 1. Study Area - Location




Fig. 2. Landuse (1988) of Thiruvananthapuram Coast

244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298





Fig. 3. Landuse (2000) of Thiruvananthapuram Coast



299 **Fig. 4. Landuse (2010) of Thiruvananthapuram Coast**

300
301 **4. CONCLUSION**

302 Agricultural land, clustered settlement area, settlement with vegetation, sandy area and rocky area are the broad landuse
303 categories delineated in the coastal zone of Thiruvananthapuram. Change in landuse for a period of twenty two years has
304 evaluated using geo-spatial data and techniques. The analysis shows that agricultural land area has decreased from
305 9.34% to 3.89% while area under clustered settlement increased from 5.75 % to 25.65% during 1988-2010. Clustered
306 settlement area has increased at the cost of agricultural lands and area under settlement with vegetation. Sandy area
307 showed an increasing trend while a decreasing trend is noticed for rocky area. Significance of landuse management as
308 part of ICZM is also highlighted. This study is expected to provide a baseline database to develop land use zonation
309 schemes and management strategies for Thiruvananthapuram coast as part of ICZM.
310

311
312
313 **COMPETING INTERESTS**

314 "Authors have declared that no competing interests exist."
315

316
317
318 **CONSENT (WHERE EVER APPLICABLE)**

319 Not applicable
320

321
322 **ETHICAL APPROVAL (WHERE EVER APPLICABLE)**

323 NOT APPLICABLE
324

325
326 **REFERENCES**

- 327
328 1. Costanza R, d' Arge R, De Groot R, Farber S, Grasso M, Hannon B, Naeem S, Limburg K, Parueto J, O' Neill RV,
329 Raskin R, Sutton P, Van den Belt M. The value of world's ecosystem services and natural capital. *Nature*. 1997;
330 387: 253-260.
331 2. Small C, Nicholls RJ. A global analysis of human settlement in coastal zones, *Journal of Coastal Research*, 2003;
332 19 (3): 584-599.
333 3. Howari FM, Benjamin RJ, Naima Bouhouche, Sandy Wyllie-Echeverria. Field and remote sensing assessment of
334 mangrove forests and sea grass beds in the North western parts of the United Arab Emirates. *Journal of Coastal*
335 *Research*, 2009; 25 (1): 48-56.
336 4. Senapati S, Gupta V. Climate change and coastal ecosystem in India-Issues in perspectives. *International Journal*
337 *of Environmental Sciences*, 2014; 4(3): 530-543.
338 5. Weismiller RA, Kristof SJ, Scholz DK, Anuta PE, Momin SA. Change detection in coastal zone environments.
339 *Photogrammetric Engineering and Remote Sensing*, 1977; 43(12): 1533-1539.
340 6. United States Geological Survey-USGS. A landuse/landcover classification system for use with Remote Sensor
341 Data. *Geological Survey Professional*, 1997; Paper 964:19.
342 7. Nayak S. Use of satellite data in coastal mapping. *Indian Cartographer*, 2001; CMMC-1: 147-156.
343 8. Chua Thia – Eng. The Integrated Management Plan for Segara Anakan-Cilacap, Central Java, Indonesia.
344 International Centre for Living Aquatic Resources Management on behalf of Association of Southeast Asian
345 Nations/United States Coastal Resources Management Project (ASEAN/US CRMP), 1992; 30.
346 9. Zhao B, Kreuter U, Li B, Ma Z, Chen J, Nakagoshi N. An ecosystem value service assessment of landuse change
347 on Chongming Island, China. *Landuse Policy*, 2004; 21(2):139-148. doi.org/10.1016/j.landusepol.2003.10.003
348 10. Sharma VVLN, Krishna GM, Malini BH, Rao KN. Landuse/Land cover change detection through remote sensing
349 and its climatic implications in the Godavari delta region. *Journal of Indian Society of Remote Sensing*, 2001;
350 29(1):85-91.

- 351 11. Lacono MJ, Levinson DM. Predicting landuse change, how much does transportation matter? Transportation
352 Research Record, 2009; 2119:130-136. DOI: 10.3141/2119-16
- 353 12. Yagoub MM, Kolan GR. Monitoring coastal zone landuse and landcover changes of Abu Dhabi using Remote
354 Sensing. Journal of the Indian Society of Remote Sensing, 2006; 34(1):57-68.
- 355 13. Shalaby A, Tateshi R. Remote sensing and GIS for monitoring land cover and land use changes in the North
356 Western coastal zone of Egypt. Applied Geography,2007; 27(1): 28-41. [doi:10.1016/j.apgeog.2006.09.004](https://doi.org/10.1016/j.apgeog.2006.09.004)
- 357 14. Baba M, Samsuddin M, Thomas KV, John Mathai, Vasudevan V, John Paul. Coastal landuse and resources.
358 Centre for Earth Science Studies, Thiruvananthapuram, 1997:108-147.
- 359 15. Census of India. Panchayat wise demographic data for Kerala State. Directorate of Census, Thiruvananthapuram,
360 2011.
- 361 16. Department of Fisheries. Facts and Figures 2012, Statistical Cell, Directorate of Fisheries, Government of Kerala,
362 Thiruvananthapuram.2012; 45-56.
- 363 17. National Centre for Earth Science Studies. Coastal zone management plan of Thiruvananthapuram District,
364 Kerala (as per CRZ Notification 2011 in 1:25000 Scale), prepared for Kerala State Council for Science and
365 Technology and Environment, Government of Kerala, 2014: 18-19.

366
367
368
369
370
371
372

UNDER PEER REVIEW