

# Changes in adiposity and dietary intake during Nowruz holiday in university students

## ABSTRACT

**Aims:** This study aimed to assess possible alterations in adiposity and dietary intake during holiday periods in healthy subjects.

**Study Design:** Four hundred and fifty-two healthy students attending the Mashhad University of Medical Sciences in all courses Iran were examined before and after Nowruz holiday (15-day-New Year holiday) in this observational, longitudinal study in February 2014.

**Methodology:** Anthropometric parameters and body composition were measured using standard protocols. Individual nutritional intakes were assessed using Dietplan6 software. SPSS software version 11.5 was used for statistical analysis.

**Results:** Of the 452 students who attended the pre-holiday visit, 433 returned for the post-holiday appointments. Of these participants, 82 (18.1 %) were men. The mean age was 24.4 years for men and 23.8 years for women ( $p=0.171$ ). All the adiposity and body composition variables significantly changed over the follow-up except for fat free mass ( $p=0.074$ ) and trunk fat free mass ( $p=0.935$ ). Mean weight of participants increased from 63.5 kg to 63.9 kg at the end of holiday (absolute change 0.37 kg,  $p<0.001$ ). Total energy intake increased by 16.4%, carbohydrate by 15.9% and total fat by 25.4%, and significant association were observed between changes in some dietary variables and change in adiposity during follow-up (15 days).

**Conclusion:** In conclusion, holidays like Nowruz play a significant role in periodical weight gain and obesity in the college students of Mashhad University. Moreover we have found that dietary intake of carbohydrate and total fat increased through Nowruz holiday. Understanding times when people are more likely to gain weight including holidays is important for the development of prevention strategies.

*Keywords: Obesity, Body composition, Body weight, Nowruz holiday, Dietary intake*

## 1. INTRODUCTION

Obesity is a potent predictor of poor health status [1]. Obesity and overweight are leading risk factors for many adverse health outcomes including type 2 diabetes, hypertension, dyslipidaemia, cardiovascular diseases and some types of cancers [2-5]. Although genetic predisposition may play an important role in the occurrence of obesity, high caloric diets and sedentary lifestyles are major contributors [6]. Weight gain during holidays is thought to be a significant determinant of obesity occurrence [7, 8]; with the incidence of a large weight gain during holiday increasing with the starting body mass index (BMI) [9]. It has been indicated that only a few weeks of overeating and reduced physical activity causes many harmful effects on the body including increased fat percentage, waist circumference, body mass index, total cholesterol and triglycerides [10, 11]. Studies recommended that a significant percentage of the average annual body weight gain in US adults occurs during the winter holiday quarter and is sustained throughout the following year [12]. Summer vacations have also been a time period of focus. Summer weight gain has been stated in most studies, with the greatest weight gain occurring in overweight children and adolescents [13-15]. These changes in anthropometric indices can be attributed to higher dietary intake and lower physical activity during holidays periods compared to non-holiday days. However, existing evidence is still conflicting. Some studies reported no significant difference in body weight during holiday [8, 16, 17].

Nowruz indicates the first day of spring, beginning of the year in the Persian calendar. It is celebrated by people from diverse ethnic and religious backgrounds in many other countries including Afghanistan, Albania, Azerbaijan, the Former Yugoslav Republic of Macedonia, India, Kazakhstan, Kyrgyzstan, Tajikistan, Turkey and Turkmenistan as well as Iran (15-day-New Year holiday) [18]. Since during this period (15-day-New Year holiday) people visit each other much more than at other times of the year, and it is traditional custom for people to prepare staple food, snacks,

34 fruits, nuts, candies, and get these ready for the visitors to eat, people tend eat much more than  
35 in usual time.

36 Regarding the long-term negative health consequences of overweight and obesity, and considering  
37 that the Nowruz holiday in Iran may contribute to annual body weight gain, it is necessary to  
38 understand factors affecting this body weight gain.. The aim of this study was to evaluate alterations  
39 in dietary intake, body weight and body composition in the healthy subjects during Nowruz.

40

## 41 **2. MATERIAL AND METHODS**

42

### 43 **2.1 Study Population**

44

45 In this observational, longitudinal study, 452 participants were recruited from among students of  
46 Faculty of Medicine, Mashhad University of Medical Sciences via internet and on campus  
47 announcements in February 2014. Eligibility criteria included an age of at least 18 years, being  
48 healthy, and willingness to attend all study visits. Subjects were excluded (12 person) if they suffered  
49 from any particular diseases affecting body composition or metabolism, were pregnant, were severely  
50 underweight (body mass index [BMI]<18.5 kg/m<sup>2</sup>), were on restriction diets and were using drugs that  
51 affect appetite and weight. Sample size were estimated based on according to the study  
52 conducted by Yanovski et al. with confidence interval of 95%, and power of 80% was calculated  
53 [19].

54

### 55 **2.2 Protocol**

56

57 Baseline measurements were obtained two weeks before the Nowruz holiday (pre-holiday and follow-  
58 up measurements after the holiday (early post-holiday). Dietary information was collected using a  
59 three different days, 24-hour food record questionnaire [20]. Individual nutritional intakes (all macro  
60 and micro nutrients intake) were assessed with the use of the Dietplan6 software (Forestfield  
61 Software Ltd., UK). All the data collected by one person (A specialist in nutrition, ZG).

62 Height (to the nearest 0.1 cm) was measured using a portable stadiometer (Seca 213, Seca Corp.,  
63 Hamburg, Germany) in a participant without shoes, stretching to the maximum height with the head  
64 positioned in the Frankfort plane. The weight, BMI, and body composition were measured by a bio-  
65 impedance analyzer (BIA) (Tanita BC-418 MA, Tanita Corp.,Japan) and participants were dressed in  
66 light clothing (i.e. no shoes, sweaters or jackets, with 0.1Kg accuracy, frequency range 50-60 Hertz)  
67 [21]. The BIA was calibrated according to the manufacturer's guidelines before each testing and the  
68 participants were informed in advance not to use any substance affecting their body composition (e.g.  
69 alcohol and coffee) 24 hours before the test [21] to evaluate the fat percentage, fat mass, fat free  
70 mass, trunk fat percentage, trunk fat mass, trunk fat free mass. All evaluations were performed by a  
71 single trained nutritionist (ZG). Waist circumference was measured by a flexible anthropometric tape  
72 measure, at the narrowest part of the trunk between the last rib and the iliac crest [21].

73 Ethical considerations

74 The study aims and methods were described to the participant and the signed informed consent  
75 obtained from them prior to participation. Moreover, the study protocol was approved by Ethical  
76 Committee in Research of Mashhad University of Medical Sciences (No: A545).

77

### 78 **2.3 Statistical Analysis**

79

80 SPSS software (version 11.5, Chicago, IL, USA) was used for statistical analysis. The Kolomogrov-  
81 Smirnov test was used to confirm the normal distribution of continuous variables. Data are expressed  
82 as mean and standard deviation (SD) for normally distributed variables and median and 25th-75th  
83 percentiles for skewed variables. Change in adiposity markers and dietary intake during follow-up  
84 were investigated using the paired t-test and Wilcoxon test in the overall cohort. Linear regression  
85 models (adjusted for sex and age) were used to assess the association between change in  
86 anthropometric variables and change of dietary intake during holiday. A p-value ≤0.05 was considered  
87 statistically significant.

88

89

## 90 **3. RESULTS**

91

### 92 **3.1 Baseline Characteristics**

93  
94  
95  
96  
97  
98  
99  
100  
101  
102

Of the 452 students who attended the pre-holiday visit, 433 returned for the post-holiday appointments and completed the study (4% drop-out rate). Of these participants, 82 (18.1 %) were men and 370 (81.9%) were women. The mean age of the participants was 23.9 years overall, 24.4 years for men and 23.8 years for women (p=0.171). With respect to adiposity factors, significant men vs. women differences were found for all the factors, except for trunk fat mass (p=0.131). Energy intake was lower in men than women (1618.9 vs. 1639.1; p=0.030), while no significant difference was found for all other dietary variables (Table 1).

**Table1. Participants' characteristics at baseline**

Variables	Overall (n=433)	Male (n=96)	Female (n=357)	P-value
Age, years	23.9±4.2	24.4±4.7	23.8±4.0	0.171
Education status, n (%)				0.863
freshman	146 (32.2)	32 (33.3)	114 (31.8)	
Sophomore	88 (19.4)	19 (19.8)	69 (19.3)	
Junior	79 (17.4)	16 (16.7)	63 (17.6)	
Senior	71 (15.6)	12 (12.5)	59 (16.5)	
graduate	70 (15.4)	17 (17.7)	53 (14.8)	
Marital status, n (%)				0.352
single	368 (81.1)	81 (84.4)	287 (80.2)	
married	86 (18.9)	15 (15.6)	71 (19.8)	
Residence, n (%)				0.175
dormitory	204 (44.9)	49 (51.2)	155 (43.3)	
own room	250 (55.1)	47 (49.3)	203 (56.7)	
Height (cm)	166.4±8.6	177.3±5.9	163.4±6.6	<0.001
Weight (Kg)	63.5±12.2	78.9±12.3	59.5±8.3	<0.001
Waist circumference (cm)	76.8 ±10.2	89.4±10.8	73.4±6.4	<0.001
Body mass index (kg/m <sup>2</sup> )	22.8±3.2	25.0±3.3	22.2±2.9	<0.001
Fat percentage (%)	24.5±6.6	17.8±6.2	26.3±5.5	<0.001
Fat mass(kg)	15.7±5.6	14.5±6.2	16.0±5.3	0.023
Fat free mass (kg)	47.7±10.0	64.4±7.9	43.2±4.0	<0.001
Trunk fat percentage (%)	21.6±7.1	17.8±6.4	22.6±6.9	<0.001
Trunk fat mass (kg)	7.5±3.2	7.9±3.4	7.3±3.2	0.131
Trunk fat free mass (kg)	26.5±5.6	35.0±4.5	24.2±3.1	<0.001
Energy (Kcal)	1639.1 (1400.9-1803.3)	1618.9 (1462.5-1838.0)	1639.1 (1382.7-1787.2)	0.030
Protein (g)	201.2 (156.9-227.9)	216.4 (183.3-231.4)	191.7 (152.9-227.6)	0.808
Carbohydrates (g)	229.2 (191.2-257.6)	240.1 (191.5-255.4)	218.2 (187.6-259.4)	0.549
Fat total (g)	50.1 (42.3-64.8)	50.1 (45.9-63.7)	50.1 (41.5-64.8)	0.353
Cholesterol (mg)	175.8 (139.8-255.6)	201.4 (147.9-270.3)	160.7 (131.3-247.3)	0.021
Saturated fatty acids (g)	15.7 (12.8-18.0)	16.1 (12.7-20.9)	15.7 (12.8-17.8)	0.194
Monounsaturated fatty acids (g)	17.0 (13.8-19.6)	18.2 (16.2-21.4)	16.1 (13.5-19.5)	0.262
Polyunsaturated fatty acids (g)	10.1 (7.6-13.8)	10.8 (8.6-14.3)	10.07 (7.51-13.5)	0.211
Fiber total (g)	21.0 (13.8-27.9)	21.2 (16.6-25.2)	20.8 (13.5-28.1)	0.211
Sugar (g)	72.3 (51.5-88.2)	60.9 (45.4-102.3)	72.3 (51.7-88.2)	0.149
Glucose (g)	8.7 (5.3-14.1)	10.2 (6.4-14.2)	8.6 (5.2-14.1)	0.160
Galactose (g)	1.4 (1.0-2.7)	1.4 (1.1-3.3)	1.4 (1.0-2.5)	0.175
Fructose (g)	11.2 (7.2-17.6)	14.1 (7.0-18.0)	10.2 (7.2-17.1)	0.165
Sucrose (g)	14.5 (8.6-20.8)	15.9 (7.9-18.5)	14.4 (9.3-20.9)	0.183

<b>Lactose (g)</b>	9.6 (5.6-16.2)	8.5 (6.3-11.3)	9.6 (5.6-16.2)	0.125
<b>Maltose (g)</b>	0.7 (0.4-1.2)	0.8 (0.4-1.3)	0.7 (0.4-1.1)	0.164

103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118

### 3.2 Changes in Adiposity, Body Composition and Dietary Intake during Follow-Up

Changes in adiposity and body composition during holiday are described in Table 2. Body weight increased by 0.4 kg (relative change 0.6%), BMI by 0.1 Kg/m<sup>2</sup> (0.6%), waist circumference by 0.3 cm (0.4%), fat percentage by 0.6% (relative change 3.3%), total fat by 0.4 kg (3.6%), trunk fat percent by 0.9% (relative change 6.6%) and absolute trunk fat mass by 0.4 kg (14.4%); with all these changes being statistically significant (all p<0.001). Contrariwise, marginal decreased in total fat free mass (absolute change -0.2 kg [relative change -0.4%], p=0.502) and trunk fat free mass (-0.01 kg [7.8%], p=0.499) did not reach statistical significance. When changes in adiposity and body composition during holiday were compared between men and women, the following differences were observed: Body weight changes were higher in men than women (0.8±1.3 g vs. 0.2±1.4 g for men and women, respectively) and also for BMI (0.2±0.4 vs. 0.09 ± 0.5 for BMI respectively for men and women).

**Table 2. Anthropometric, adiposity and dietary intake baseline**

Variables		Absolute change (immediate holiday baseline - post-holiday)	Relative change (immediate holiday baseline - post-holiday)	p-value	P effect of gender
<b>Anthropometric factors</b>	Weight (kg)	0.32±1.44	0.62±2.63	<0.001	<0.001
	Waist circumference (cm)	0.32±0.01	0.31±0.02	<0.001	<0.001
	Body mass index (kg/m <sup>2</sup> )	0.1±0.5	0.6±2.6	<0.001	0.005
	Fat percentage (%)	0.5±1.6	3.3±9.0	<0.001	0.234
<b>Adiposity factors</b>	Fat mass (kg)	0.3±1.1	3.6±9.9	<0.001	0.002
	Fat free mass (kg)	-0.2±2.5	-0.3±7.3	0.074	0.502
	Trunk fat percentage (%)	0.9±2.3	6.6±16.7	<0.001	0.159
	Trunk fat mass (kg)	0.4±1.4	14.4±80.4	<0.001	0.129
<b>Dietary factors</b>	Trunk fat free mass (kg)	-0.01±2.5	7.8±92.2	0.935	0.499
	Energy (Kcal)	246.4 (-169.7 to 65.8)	16.4 (-11.7 to 45.9)	<0.001	0.310
	Protein (g)	19.5 (-37.0 to 80.6)	11.1 (-16.2 to 47.3)	0.315	0.244
	Carbohydrates (g)	29.1 (-49.3 to 71.0)	15.9 (-23.5 to 40.7)	0.011	0.938
	fat total (g)	13.8 (-3.5 to 36.5)	25.4 (-6.6 to 81.9)	<0.001	0.331
	Cholesterol (mg)	6.2 (-61.8 to 142.4)	3.9 (-26.8 to 81.8)	<0.001	0.222
	Saturated fatty acids (g)	6.5 (-0.6 to 12.9)	37.9 (-3.9 to 85.0)	0.342	0.150
	Monounsaturated fatty acids (g)	3.8 (-2.1 to 11.6)	22.7 (-12.9 to 67.8)	0.598	0.230
	Polyunsaturated fatty acids (g)	0.4 (-2.2 to 6.4)	5.4 (-18.1 to 60.5)	0.034	0.219
	Fiber total (g)	-1.3 (-12.2 to 8.6)	-8.1 (-45.2 to 63.9)	0.004	0.124
	Sugar (g)	15.8 (-13.3 to 47.6)	22.9 (-16.9 to 75.1)	0.005	0.073
	Glucose (g)	5.5 (-2.7 to 10.4)	61.3 (-21.4 to 168.0)	0.137	0.123
	Galactose (g)	0.9 (-1.1 to 2.5)	55.6 (-73.5 to 208.6)	0.009	0.124
Fructose (g)	4.6 (-5.3 to 11.5)	43.0 (-32.5 to 149.0)	0.100	0.124	

Sucrose (g)	-1.4 (-8.4 to 8.9)	-8.3 (-48.6 to 86.0)	0.007	0.220
Lactose (g)	0.2 (-7.5 to 6.5)	1.2 (-59.4 to 99.8)	0.006	0.058
Maltose (g)	0.1 (-0.3 to 0.7)	23.0 (-38.5 to 146.5)	0.006	0.163

*P-values are from paired sample t-test and Wilcoxon test. Data expressed as a mean ± standard deviation for anthropometric and adiposity factors, for dietary factors data are based on mean and confidence interval.*

Changes in dietary intake are also described in Table 2. Significant increased were observed for total energy (relative increased 16.4%,  $p<0.001$ ), carbohydrate (16%,  $p=0.011$ ), total fat (25.4%,  $p<0.001$ ), cholesterol (4%,  $p<0.001$ ), polyunsaturated fatty acids (5.5%,  $p=0.034$ ), while fibre intake significantly decreased (-8.2%,  $p=0.004$ ). Change in protein intake was not significant ( $p=0.315$ ). The following differences were observed when changes in dietary intakes were compared between men and women: energy intake in men was higher in men than women (244±804 kcal vs. 331±669g for men and women respectively) and also for sugar intake (24.3±59.7g vs. 7.2 ± 80 g for sugar intake respectively for men and women).

### 3.3 Association between Changes in Dietary Intake and Changes in Adiposity during Holiday

The age and sex adjusted regression coefficients for the association between changes in dietary intake and changes in adiposity are described in Table 3. Change in saturated fatty acids, galactose and lactose (all  $p<0.001$ ) were significantly and positively associated with changes in weight and BMI, while change in monounsaturated fatty acids ( $p=0.011$ ), cholesterol ( $p=0.009$ ), fructose, sucrose and lactose (all  $p<0.01$ ) were significantly associated with change in fat mass.

**Table 3. Change in anthropometric, adiposity and dietary intake factors during follow-up**

Variables (absolute change)	Weight (absolute change)		Body mass index (absolute change)		Fat mass (kg) (absolute change)		Trunk mass (kg) (absolute change)		Fat (Kg)
	$\beta$	p-value	$\beta$	p-value	$\beta$	p-value	$\beta$	p-value	
Energy(Kcal)	0.29	0.842	0.14	0.920	0.33	0.824	1.78	0.235	
Protein (g)	-1.29	0.045	-1.29	0.045	-0.60	0.359	-1.77	0.007	
Carbohydrates (g)	-0.22	0.795	-0.10	0.903	-0.19	0.829	-1.07	0.227	
fat total (g)	-0.52	0.740	-0.47	0.766	-0.90	0.577	-1.95	0.222	
Cholesterol (mg)	0.10	0.127	0.11	0.089	-0.18	0.009	-0.19	0.006	
Saturated fatty acids (g)	1.64	0.038	1.64	0.037	0.84	0.293	2.30	0.004	
Monounsaturated fatty acids (g)	2.05	0.085	2.07	0.080	3.07	0.011	3.13	0.009	
Polyunsaturated fatty acids (g)	1.24	0.275	1.26	0.265	-0.17	0.879	-2.45	0.033	
Fiber total (g)	-0.03	0.894	-0.04	0.865	-0.50	0.084	-0.22	0.451	
Sugar (g)	0.31	0.228	0.35	0.184	0.26	0.321	-0.19	0.475	
Glucose (g)	-0.57	0.661	-0.80	0.537	-0.86	0.516	1.49	0.256	
Galactose (g)	-6.98	<0.001	-7.08	<0.001	-2.13	0.214	-2.62	0.125	
Fructose (g)	1.18	0.214	1.29	0.173	2.49	0.010	0.50	0.603	
Sucrose (g)	-0.77	0.081	-0.82	0.064	-1.31	0.004	-0.07	0.874	
Lactose (g)	2.36	<0.001	2.41	<0.001	1.67	<0.001	2.00	<0.001	
Maltose (g)	1.43	0.677	1.44	0.674	-1.63	0.642	0.55	0.874	

*Models were adjusted for sex and age.*

## 4. DISCUSSION

The present study investigated the effect of holiday on anthropometric, adiposity and dietary intake among medical students in Iran. We observed that during this relatively short holiday, total and regional adiposity significantly increased, in parallel with increases in energy intake, fat, carbohydrate intake, and a reduction in fibre intake. There was further evidence from regression analysis that

150 changes in adiposity were correlated with changes in dietary intakes. Although the magnitude of  
151 changes was inconsiderable, cumulative effect over successive holiday will likely have profound effect  
152 on obesity development and maintenance.

153 We observed increased in adiposity during holiday are in line with the findings of Cooper et al. [13],  
154 Montero et al. [11], Yanovski et al. [19], Costa et al. [7] and Branscum et al. [22]. Similarly, Payab et  
155 al. recently reported significant weight gain during Nowruz holiday among the staff of a hospital in  
156 Tehran, Iran [23]. Cooper et al. conducted a prospective study to determine if a 1- to 3-week vacation  
157 in adults leads to weight gain and whether that gain persists 6 weeks later. Their results showed that  
158 holidays causes important weight gain, and this weight gain persisted at the 6-week follow-up period.  
159 In accordance with our findings, they reported that the weight gain seems to be determined by  
160 increased energy intake above energy requirements [13].

161 However, some studies reported no significant difference in body weight during holiday [8, 16, 17].  
162 The difference in body weight and BMI following holiday in our study seems inconsiderable. However,  
163 the accumulating such small gains over successive holidays across years is likely to translate into  
164 sizable gains, which in turn can explain some of the rise in obesity among adults [19].

165 Other body composition variables including fat mass and fat percent were significantly different  
166 between pre- and post-holiday visits in present study which is in accordance with Hull et al. [8] and  
167 Cristi-Montero et al. [11]. Significant increase in trunk fat mass of the participants during holiday in  
168 this study is in line with the findings of Costa et al. and Hull et al. [7, 8]. In contrast, Wagner et al.  
169 found no significant difference in fat percent [17], while Payab et al. [23] observed a significant  
170 decrease in fat percent, as well as a significant increase in fat free mass among the hospital staff  
171 during Nowruz holiday.

172 The present study, for the first time, analysed the macro and micro nutrient intake before and after  
173 holiday. Our findings suggest that holiday had a significant effect on the components of food intake  
174 including total calories, carbohydrates and fat intakes. Grimes et al. carried out a national  
175 investigation in Australia to compare the dietary component of the student on non-school days with  
176 school days [24]. In line with our findings, they reported that intakes of total fat, sugars, saturated fat  
177 and the energy density of foods consumed is higher during holiday's days. In addition, the absolute  
178 intake of sodium was higher on non-school days. In accordance with our findings, Shahar et al.  
179 observed an increase in fat and calorie intake during winter holidays [25]. Moreover, Ma et al.  
180 demonstrated that daily calorie intake was higher during autumn holiday compared to spring and  
181 showed a slight seasonal variation with a peak for carbohydrate intake in spring and for total fat in fall  
182 in overweight/obese participants [26]. Our regression analysis also revealed some associations in the  
183 causal direction between changes in dietary intake and change in adiposity during holiday.

184

## 185 **5. STRENGTHS AND LIMITATIONS**

186

187 Follow-up the participants after holiday and assessing macro-micro nutrient intake in the mentioned  
188 times are the strengths of this study. Moreover, we collected no data for regular times (one- or two-  
189 weeks (post – pre) holiday) to figure out if college student are increasing their weight constantly.  
190 However, one limitation of the study was that the participants were informed of the aim of the study,  
191 which might possibly have biased our findings. Secondly, physical activity, which might be a  
192 confounding factor, was not assessed in our study. Lack of a third visit showing the change in weight  
193 and body composition a few months later was another limitation of the study. Generally, limiting the  
194 intake of high-calorie foods and having regular physical activity during the Nowruz holiday is strongly  
195 recommended.

196

## 197 **6. CONCLUSION**

198

199 Our study findings suggest that Nowruz holiday could play a significant role in periodical weight gain  
200 and obesity in this population of medical students, which can be contribute to weight gain and  
201 maintenance in later age. Considering the health consequences of overweight and obesity,  
202 understanding times when people are more likely to gain weight including holidays is important for the  
203 development of prevention and control strategies.

204

## 205 **REFERENCES**

206

207 1. Moreira P, Padez C, Mourao I, Rosado V. Dietary calcium and body mass index in Portuguese  
208 children. *European journal of clinical nutrition*. 2005 Jul;59(7):861-7.

- 209 2. Alharbi NS, Almutari R, Jones S, Al-Daghri N, Khunti K, de Lusignan S. Trends in the prevalence of  
210 type 2 diabetes mellitus and obesity in the Arabian Gulf States: systematic review and meta-analysis.  
211 *Diabetes Res Clin Pract.* 2014 Nov;106(2):e30-3.
- 212 3. An R. Prevalence and Trends of Adult Obesity in the US, 1999–2012. *ISRN Obesity.* 2014  
213 01/06/received 12/17/accepted;2014:185132.
- 214 4. Heidari-Bakavolia A, Esmaeilib H, Hosseinib Z, Moohebatia M, Azarpazhooaha M, Mazidig M, et al.  
215 Prevalence of obesity in Iran and its related socio-economic factors. *Mediterranean Journal of*  
216 *Nutrition and Metabolism.* 2015;8:109-18.
- 217 5. Kwagyan J, Retta TM, Ketete M, Bettencourt CN, Maqbool AR, Xu S, et al. OBESITY AND  
218 CARDIOVASCULAR DISEASES IN A HIGH-RISK POPULATION: EVIDENCE-BASED APPROACH  
219 TO CHD RISK REDUCTION. *Ethnicity & disease.* 2015 Spring;25(2):208-13.
- 220 6. Ghadiri-Anari A, Jafarizadah M, Zare A, Mozaffari-Khosravi H, Afkhami-Ardekani M, Shojaoddiny-  
221 Ardekani A. Prevalence of Obesity and Overweight among Adults in Iranian Population (Yazd  
222 Province). *Iranian Journal of Diabetes and Obesity.* [Research]. 2013;5(2):67-70.
- 223 7. Costa C, Moreira P, Teixeira VH. Holiday weight gain in university students. 2007.
- 224 8. Hull HR, Hester CN, Fields DA. The effect of the holiday season on body weight and composition in  
225 college students. *Nutrition & metabolism.* 2006;3:44.
- 226 9. Schoeller DA. The effect of holiday weight gain on body weight. *Physiology & behavior.* 2014  
227 Jul;134:66-9.
- 228 10. Cristi-Montero C. Are weight gain prevention program effective in schools? *Rev Int Med Cienc Act*  
229 *Fi's Deporte.* 2012, 12:287–298.
- 230 11. Cristi-Montero C, Munizaga C, Tejos C, Ayala R, Henriquez R, Solis-Urra P, et al. Variations of  
231 body composition, physical activity and caloric intake in schoolchildren during national holidays.  
232 *Eating and weight disorders : EWD.* 2016 Jun;21(2):251-5.
- 233 12. Cook CM, Subar AF, Troiano RP, Schoeller DA. Relation between holiday weight gain and total  
234 energy expenditure among 40- to 69-y-old men and women (OPEN study). *The American journal of*  
235 *clinical nutrition.* 2012 Mar;95(3):726-31.
- 236 13. Cooper JA, Tokar T. A prospective study on vacation weight gain in adults. *Physiology &*  
237 *behavior.* 2016 3/15;156:43-7.
- 238 14. Franckle R, Adler R, Davison K. Accelerated Weight Gain Among Children During Summer  
239 Versus School Year and Related Racial/Ethnic Disparities: A Systematic Review. *Preventing Chronic*  
240 *Disease.* 2014 06/12;11:E101.
- 241 15. Moreno JP, Johnston CA, Woehler D. Changes in weight over the school year and summer  
242 vacation: results of a 5-year longitudinal study. *J Sch Health.* 2013 Jul;83(7):473-7.
- 243 16. Reid R, Hackett AF. Changes in nutritional status in adults over Christmas 1998. *Journal of*  
244 *Human Nutrition and Dietetics.* 1999;12(6):513-6.
- 245 17. Wagner DR, Larson JN, Wengreen H. Weight and body composition change over a six-week  
246 holiday period. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity.* [journal  
247 article]. 2013;17(1):e54-e6.
- 248 18. International Day of Nowruz. The United Nations; Available from:  
249 <http://www.un.org/en/events/nowruzday/>.
- 250 19. Yanovski JA, Yanovski SZ, Sovik KN, Nguyen TT, O'Neil PM, Sebring NG. A prospective study of  
251 holiday weight gain. *The New England journal of medicine.* 2000 Mar 23;342(12):861-7.
- 252 20. Mahan LK, Raymond JL, Escott-Stump S. *Krause's Food & the Nutrition Care Process.* 13th  
253 ed2011.
- 254 21. Mazidi M, Rezaie P, Norouzy A, Saeb MH, Mehdizadeh Hakkak A, Balali S, et al. Investigating the  
255 relation between macronutrients intake and anthropometric indices. *Mediterranean Journal of*  
256 *Nutrition and Metabolism.* 2015;8(2):131-8.
- 257 22. Branscum PW, Kaye G, Succop P, Sharma M. An evaluation of holiday weight gain among  
258 elementary-aged children. *Journal of clinical medicine research.* 2010;2(4):167-71.
- 259 23. Payab M, Hasani-Ranjbar S, Zahedi H, Qorbani M, Shateri Z, Larijani B, et al. The effect of  
260 Norouz holiday on anthropometric measures and body composition. *Journal of Diabetes and*  
261 *Metabolic Disorders.* 2015 02/2509/22/received02/05/accepted;14:7.
- 262 24. Grimes CA, Riddell LJ, Nowson CA. Nutrient and core and non-core food intake of Australian  
263 schoolchildren differs on school days compared to non-school days. *Appetite.* 2014 12/1;83:104-11.
- 264 25. Shahar DR, Froom P, Harari G, Yerushalmi N, Lubin F, Kristal-Boneh E. Changes in dietary intake  
265 account for seasonal changes in cardiovascular disease risk factors. *European journal of clinical*  
266 *nutrition.* 1999 May;53(5):395-400.

267 26. Ma Y, Olendzki BC, Li W, Hafner AR, Chiriboga D, Hebert JR, et al. Seasonal variation in food  
268 intake, physical activity, and body weight in a predominantly overweight population. *European journal*  
269 *of clinical nutrition*. 2006;60.