

1 **EFFECTS OF VARYING CONCENTRATIONS OF CRUDE OIL ON SOME**  
2 **PHYSICOCHEMICAL PROPERTIES OF AGRICULTURAL SOIL**

3

4 **ABSTRACT**

5 This research investigated the effects of varying concentrations of crude oil on some  
6 physicochemical characteristics of crude oil polluted agricultural soils from Igodan- Lisa,  
7 Oba-Ile and Ido-Ani areas of Ondo State, Nigeria. The soil samples were exposed to 1- 4%  
8 (w/w) crude oil and analyzed monthly for six periods using standard physical and chemical  
9 analytical techniques. Results indicated that the physicochemical properties were altered. The  
10 physicochemical parameters varied with increase in the amount of crude oil spilled and time.  
11 The pH and moisture content (MC) progressively decreased with increase in concentration of  
12 crude oil applied to the samples. Polluted soils had lower pH values (4.91- 6.17) and MC  
13 (15.24% to 26.83%) relative to control samples. The organic matter content increased with  
14 increased amount of crude oil spilled in the range of 6.65 - 10.93%. The organic carbon  
15 contents progressively increased with concentration of crude oil and sampling days. At 4%  
16 crude oil pollution, the organic carbon content in the samples were 6.04 - 8.28%, 5.39 -  
17 7.82% and 6.05 - 8.21% for Igodan- Lisa, Oba-Ile and Ido-Ani soils respectively at 0-180  
18 days of experiment. The changes in soil physicochemical suggested that soil integrity and  
19 quality is altered by crude oil contamination. The increased acidity with time also suggested  
20 bioremediation by intrinsic microorganisms.

21 **Keywords:** Varying concentrations, Physicochemical characteristics, Agricultural soils,  
22 Crude oil, bioremediation, soil quality

23

## 24 **1. INTRODUCTION**

25 Oil and gas is a major resource and energy that has been driving the economy of Nigeria  
26 since about six decades when commercial exploitation of petroleum started. Apart from crude  
27 oil being the mainstay of the Nigeria's economy, industries also rely extensively on  
28 petroleum derivatives without which they cannot function and produce to optimal capacity.  
29 Agriculture which is the main occupation of the people of Ondo State provides the second  
30 largest support to the nation's economy. Unfortunately, the processes of exploitation,  
31 exploration, processing and storage as well as transportation of petroleum and its derivatives  
32 have resulted in enormous abuse of man's environment especially in the Niger- Delta Region  
33 of Nigeria, rendering farmlands to wastelands as a result of the toxic effects of spilled oil on  
34 agricultural lands.

35 Crude oil spillage into the environment is a common occurrence world over. The  
36 discharge of petroleum hydrocarbon and its derivatives is of greater dimension in the Niger  
37 Delta region due mainly to diverse human activities; including pipeline vandalism,  
38 negligence during production operations and fuel tanker loading processes, corrosion and  
39 pipeline leakages and oil tanker terrestrial accidents. Pollution of the environment by  
40 petroleum occurs when petroleum or its derivatives are introduced or spilled into the  
41 environment at levels harmful either directly to the environment or indirectly to the  
42 dependents of the environment [1]

43 Soil is a primary receiver of crude oil spill as well as many different types of products and  
44 chemicals such as herbicides, biocides and pesticides which are hydrocarbon products. Soil  
45 can be defined in many ways to suit different professions and purposes. To the agriculturist,  
46 soil is a medium for growth, anchorage for plants, providing nutrients (macro and micro),  
47 water and air necessary for plant growth, crop production and profitable agriculture [2]. Soil

48 also provides habitat for micro flora and micro fauna and a dynamic entity where complex  
49 interactions among its biological, chemical and physical components take place. All these  
50 components and properties determine the functioning of soil for different purposes [3]. Soil  
51 type and properties affect agricultural productivity and quality through its function as a  
52 medium for plant growth and as regulator of water flow and nutrient cycling [3]. Soil quality  
53 is the capacity of soil to function within ecosystem boundaries. Soil is made of four  
54 components; sand, silt, clay and humus (decayed organic materials). Sand is important for  
55 keeping the soil loose, aerated and well drained. Clay minerals hold water and nutrients in the  
56 soil just loosely enough to allow plant roots absorb them. The humus component provides the  
57 bulk of soil's fertility [4]. Among soil physicochemical properties normally used to evaluate  
58 soil quality include soil texture, bulk density, organic carbon content, soil reaction (pH),  
59 cation exchange capacity while soil respiration, earthworm presence and microbial  
60 biodiversity are biological factors.

61 Soil pH is the degree of acidity or alkalinity of soil. The pH value is a very important  
62 property that affects many other physicochemical and biological properties. Soil reaction  
63 (pH) measures acidity of soil on a scale of 0 – 14. A pH value of 6.5-7.5 is considered  
64 optimum for growth of many plants [5]. Extreme pH values decrease microbial activity in  
65 soils, thereby affecting many soil processes such as organic matter decomposition,  
66 nitrification and the biological nitrogen fixation. Water content or moisture content is the  
67 quantity or amount of water contained in a material. Oyem and Oyem [5] asserted that water  
68 in the soil, in term of volume and movement, is the single factor determining plant growth  
69 and the solvent in which all chemical reactions take place as well as the most important factor  
70 determining remediation of salt water and hydrocarbon spills. Microbial activity in soil is  
71 generally greatest at water contents ranging between 50-80% of the maximum water holding

72 capacity [1]. Other soil characteristics affecting soil quality include organic matter and  
73 organic carbon contents. Soil organic matter is principal soil property affecting biological  
74 activity in soil. It is composed of organic compounds from decomposed remains of living  
75 organisms and their waste products in the environment. It functions as the carbon source for  
76 many soil organisms including soil micro biota [3]. It has also been reported that the  
77 interactions between organic pollutants and soil particles are largely determined by soil  
78 organic matter content [6].

79 Crude oil is a complex mixture of different kinds of hydrocarbons, liquid in their natural  
80 state and composed of aliphatics, aromatics and asphaltene fractions along with nitrogen,  
81 sulphur and oxygen containing compounds. Many of these compounds are known to be  
82 highly toxic to humans, animals, plants and microorganisms. The sources of crude oil spill  
83 into the environment differ and the amount spilled varies from minor to disaster. Crude oil  
84 destroys soil richness, causes alterations in soil physicochemical and microbiological  
85 properties [7] and cause severe damages to the environment and all forms of life dependent  
86 on the environment [1]. The release of crude oil causes enormous damages to the  
87 environment due to the presence of many toxic compounds such as polycyclic aromatic  
88 hydrocarbons, benzene and its substituted and cycloalkane rings in relatively high  
89 concentration [8]. Crude oil spillage on agricultural land and the attendant fouling effect can  
90 render the soil (especially the biologically active surface layer) toxic and unproductive [9,  
91 10]. The overall effects of crude oil spillage on agricultural land may be due to the nutritional  
92 imbalance (especially of carbon and nitrogen) created by the spilled oil (9, 11], causing  
93 reduced agricultural yield and consequently adversely affecting the socio-economic lives of  
94 the people residing in the affected area due to high levels of unemployment and poverty rates  
95 and hence increased hunger.

96 The high incidence and frequency of crude oil spill and the concomitant consequences on  
97 both biotic and abiotic components of the ecosystem are of great concern to Environmental  
98 Researchers. These concerns have consequently resulted in the development of many  
99 remediation options towards returning crude oil polluted environment to its pre-  
100 contamination status in order to restore soil quality to support agriculture. Remediation of  
101 crude oil contaminated site refers to removing or transforming contaminants to harmless or  
102 less harmful substances [12]. Several physicochemical and biological approaches have been  
103 applied to remediate polluted soil and water environments. The effects and time for  
104 reclamation of crude oil polluted soil depend on the quantity and the concentration of the  
105 pollutant (13, 14]. Among the remediation techniques, bioremediation as a contaminant  
106 removal strategy relies on the metabolic capabilities of microorganisms to detoxify or remove  
107 organic pollutants from the environment. It is considered a safe, ecosystem friendly and cost  
108 effective approach relative to the physicochemical methods [15]. The effectiveness of  
109 bioremediation technologies applied to hydrocarbon polluted soil is dependent upon physical  
110 and chemical conditions as well as the presence of native microbial population (primarily  
111 bacteria, yeast and mold) with ability to degrade hydrocarbon pollutant and environmental  
112 conditions. In order to monitor the effectiveness of bioremediation strategy, it becomes  
113 pertinent to understand the effects of crude oil concentration on the technological parameters  
114 affecting soil quality. Therefore, this research is undertaken to evaluate the effects of varying  
115 concentrations of crude oil on some physicochemical properties of agricultural soils in order  
116 to evaluate the progress and effectiveness of bioremediation in restoration of soil quality to  
117 boost agricultural productivity and improve the livelihood the people in the risk areas of  
118 crude oil pollution.

119

## 120 **2. MATERIALS AND METHODS**

### 121 **2.1 Sample collection**

122 The soil samples used in this study were collected from Igodan- Lisa (6° 27' 0''N, 4°  
123 47'0''E), Oba- Ile (7° 16' 0''N, 5° 15' 0''E) and Ido-Ani (7° 17' 0''N, 5° 52'0''E) all in Ondo  
124 State, Nigeria. According to Ikuesan [16] the soil texture and total petroleum hydrocarbon  
125 contents of the soils were as follows; Igodan- Lisa (Sand; 51.32%, Silt; 36.34%, Clay;  
126 13.21%, TPH; 13.27mg/kg) Oba- Ile (Sand; 67.35%, Silt; 20.35%, Clay; 12.67%, TPH;  
127 10.57mg/kg) and Ido-Ani (Sand; 68.36%, Silt; 15.68%, Clay; 14.66%, TPH; 11.96mg/kg).

128 The samples were collected using the hand auger at depth of 15-20cm into sterile black  
129 cellophane bags. The crude oil used in this study was standard - grade crude oil (Bonny light)  
130 collected from Bille Flow Station in Port Harcourt, Nigeria.

### 131 **2.2 Soil Treatment**

132 The samples were then partially air-dried at  $28 \pm 2^\circ\text{C}$  and passed through a 2mm mesh to  
133 remove large particles, debris and stones. A total of 45 plastic buckets were filled with sieved  
134 experimental soil and then used for this study to prepare triplicate samples of each  
135 concentration (0-4% w/w). Crude oil was then added to the soil in the plastic buckets at  
136 different concentrations to obtain 0-4% (w/w) contamination according to the method of [17,  
137 18]. The crude oil was then thoroughly mixed with the soil using the spatula. The untreated  
138 samples (0% w/w) were the controls.

### 139 **2.3 Analysis of Soil Physicochemical Properties**

140 The physicochemical characteristics of control and polluted soil samples were determined as  
141 follows. The pH of samples was determined by the glass electrode pH meter (Jenway 3051)  
142 method in 1:1w/v soil: water slurry which was standardized at pH 7.0 using phosphate buffer  
143 solution [19]. The weight loss method was used to determine the moisture content [20, 21].

144 The moisture content was adjusted to 25% by adding water in order to enhance microbial  
145 activities. Organic matter (OM) and Organic carbon (OC) were measured by dichromate-  
146 oxidation of Walkley-Black method [19, 21].

147 The effects of crude oil contamination on the physicochemical properties of soil was  
148 then determined following the methods described above. The physicochemical parameters  
149 evaluated were pH, moisture content, percentage organic matter and organic carbon.  
150 Triplicate samples of the various treatment containers were tilled weekly with spatula for  
151 necessary aeration and proper mixing of the oil with the soils. Analysis of the  
152 physicochemical characteristics of each agricultural soil sample (0-4% w/w) from Igodan -  
153 Lisa, Oba-Ile and Ido-Ani were carried out at 7 days post contamination as day zero [14] and  
154 then periodically evaluated at intervals of 30 days using standard physical and chemical  
155 methods as earlier described.

156


### 157 **3. RESULTS**

158 The results obtained in this study revealed that crude oil pollution of soil at all levels of  
159 contamination resulted in a remarkable alterations in soil physicochemical properties  
160 affecting soil quality. The physicochemical characteristics of crude oil polluted (varying  
161 concentration of 1 - 4%) and unpolluted (controls) agricultural soils over a study period of  
162 180 days are shown in figures 1-3 and tables 1 (a - c). The observed changes were much more  
163 noticed in the polluted soils as the concentration of applied crude oil increased. The soil  
164 acidity (pH), organic matter (OM) and organic carbon (OC) contents increased as the level of  
165 crude oil contamination increased. Conversely, the moisture content decreased with increase  
166 in concentration of crude oil applied to soil. The result of the pH of the soil samples is shown  
167 in figures 1(a-c). The pH for all treated samples were lower at the end of the study period

168 compared with the unpolluted samples, thus, the soil acidity increased. The pH of the control  
169 samples was almost stable at near neutral over the study period. The pH of crude oil  
170 contaminated soils progressively decreased both with increase in concentration of crude oil  
171 applied and sampling days. The experimentally polluted samples had acidic pH of 4.91-6.17  
172 at 1- 4% (w/w) levels of contamination compared with 7.03 – 7.28 in the control samples  
173 during the study period of 180 days.

174

175



176


177 Figure 1a: Effect of crude oil concentration on the pH of Igodan- Lisa soil

178

179

180






181

182 Figure 1b: Effect of crude oil concentration on the pH of Oba- Ile soil

183




184

185 Figure 1c: Effect of crude oil concentration on the pH of Ido- Ani soil

186


187            Conversely, there was also a significant reduction in moisture content (MC) of crude  
 188 oil contaminated soils with increase in the amount of applied crude oil, but showed  
 189 progressive and gradual increase with time in all the samples. The MC values of polluted  
 190 soils rangebetween 15.24% -26.83% relative to control samples (23.03 - 27.26%) as shown in  
 191 figures 2(a - c).

192




193

194            Figure 2a: Effect of crude oil concentration on the moisture content of Igodan- Lisa soil



195

196 Figure 2b: Effect of crude oil concentration on the moisture content of Oba-Ile soil



197

198 Figure 2c: Effect of crude oil concentration on the moisture content of Ido- Ani soil

199

200 In contrast to the observed decrease in pH and MC, the organic matter (OM) and organic  
 201 carbon (OC) contents respectively showed significant increases (figures 3a-c) and (tables 1a-  
 202 c) contents with increase in the concentration of crude oil applied. However, while the OM

203 values which increased with concentration decreased as the sampling days progressed, the  
204 OC of samples progressively increased. Results revealed lower OM (3.13 – 7.72) at the end  
205 of study in polluted soil samples compared to 5.73 – 8.11 in unpolluted soils, whereas, the  
206 polluted samples showed significant increase in OC (3.26 – 8.28) relative the control (2.25 –  
207 4.45) samples. The OC contents shown in tables 1(a - c) progressively increased with  
208 concentration of crude oil applied and sampling days. Tables 1(a - c) revealed that at 4%  
209 (w/w) crude oil contamination of the soil samples, the OC content in samples were 6.04 -  
210 8.28%, 5.39 - 7.82% and 6.05 - 8.21% respectively for Igodan- Lisa, Oba- Ile and Ido Ani at  
211 0 - 180 days sampling periods.

212

213

214

215


216

217

218

219


220



221


222 Figure 3a: Effect of crude oil concentration on the organic matter content of Igodan- Lisa soil

223



224

225 Figure 3b: Effect of crude oil concentration on the organic matter content of Oba- Ile soil



226

227 Figure 3c: Effect of crude oil concentration on the organic matter content of Ido - Ani

228 Table 1a: Effect of crude oil concentration on the organic carbon content of Igodan- Lisa soil

Time (Days)	SS <sub>3</sub>	SS <sub>3A</sub>	SS <sub>3B</sub>	SS <sub>3C</sub>	SS <sub>3D</sub>
0	2.80 ± 0.01 <sup>a</sup>	3.74 ± 0.02 <sup>a</sup>	4.38 ± 0.00 <sup>a</sup>	4.99 ± 0.01 <sup>a</sup>	6.04 ± 0.01 <sup>a</sup>
30	3.29 ± 0.01 <sup>b</sup>	4.190 ± 0.00 <sup>b</sup>	4.45 ± 0.01 <sup>b</sup>	5.12 ± 0.00 <sup>b</sup>	6.20 ± 0.01 <sup>b</sup>
60	3.79 ± 0.01 <sup>c</sup>	4.49 ± 0.01 <sup>c</sup>	4.78 ± 0.00 <sup>c</sup>	5.46 ± 0.02 <sup>c</sup>	6.61 ± 0.01 <sup>c</sup>
90	3.91 ± 0.01 <sup>d</sup>	4.66 ± 0.01 <sup>d</sup>	4.97 ± 0.02 <sup>d</sup>	5.67 ± 0.01 <sup>d</sup>	6.89 ± 0.01 <sup>d</sup>
120	4.11 ± 0.01 <sup>c</sup>	4.86 ± 0.00 <sup>e</sup>	5.17 ± 0.01 <sup>e</sup>	5.90 ± 0.01 <sup>e</sup>	7.16 ± 0.00 <sup>e</sup>
150	4.33 ± 0.06 <sup>f</sup>	5.18 ± 0.01 <sup>f</sup>	5.60 ± 0.02 <sup>f</sup>	6.42 ± 0.01 <sup>f</sup>	7.77 ± 0.01 <sup>f</sup>
180	4.44 ± 0.02 <sup>g</sup>	5.89 ± 0.02 <sup>g</sup>	6.53 ± 0.02 <sup>g</sup>	7.15 ± 0.01 <sup>g</sup>	8.28 ± 0.00 <sup>g</sup>

229 Legend:

230 SS<sub>3</sub> = 0% Contamination (control)

231 SS<sub>3A</sub> = 1% Crude Oil Contamination

232 SS<sub>3B</sub> = 2% Crude oil Contamination

233 SS<sub>3C</sub> = 3% Crude Oil Contamination

234 SS<sub>3D</sub> = 4% Crude Oil Contamination

235

236 Table 1b: Effect of crude oil concentration on the organic carbon content of Oba-Ile soil

Time (Days)	SS <sub>4</sub>	SS <sub>4A</sub>	SS <sub>4B</sub>	SS <sub>4C</sub>	SS <sub>4D</sub>
0	2.25 ± 0.01 <sup>a</sup>	3.26 ± 0.00 <sup>a</sup>	3.87 ± 0.01 <sup>a</sup>	4.49 ± 0.01 <sup>a</sup>	5.39 ± 0.01 <sup>a</sup>
30	2.78 ± 0.01 <sup>b</sup>	3.18 ± 0.00 <sup>b</sup>	4.14 ± 0.01 <sup>b</sup>	4.67 ± 0.01 <sup>b</sup>	5.63 ± 0.02 <sup>b</sup>
60	3.25 ± 0.01 <sup>c</sup>	4.30 ± 0.02 <sup>c</sup>	4.67 ± 0.01 <sup>c</sup>	5.02 ± 0.00 <sup>c</sup>	6.10 ± 0.01 <sup>c</sup>
90	3.35 ± 0.00 <sup>d</sup>	4.45 ± 0.01 <sup>d</sup>	4.82 ± 0.01 <sup>d</sup>	5.28 ± 0.00 <sup>d</sup>	6.39 ± 0.01 <sup>d</sup>
120	3.53 ± 0.01 <sup>e</sup>	4.66 ± 0.00 <sup>e</sup>	5.04 ± 0.01 <sup>e</sup>	5.59 ± 0.02 <sup>e</sup>	6.62 ± 0.01 <sup>e</sup>
150	3.69 ± 0.01 <sup>f</sup>	4.83 ± 0.00 <sup>f</sup>	5.28 ± 0.00 <sup>f</sup>	6.18 ± 0.01 <sup>f</sup>	7.19 ± 0.01 <sup>f</sup>
180	3.90 ± 0.00 <sup>g</sup>	5.09 ± 0.01 <sup>g</sup>	6.19 ± 0.02 <sup>g</sup>	7.03 ± 0.01 <sup>g</sup>	7.82 ± 0.00 <sup>g</sup>

237 Legend:

238 SS<sub>4</sub> = 0% Contamination (control),

239 SS<sub>4A</sub> = 1% Crude Oil Contamination

240 SS<sub>4B</sub> = 2% Crude oil Contamination

241 SS<sub>4C</sub> = 3% Crude Oil Contamination

242 SS<sub>4D</sub> = 4% Crude Oil Contamination

243

244

245

246

247

248

249

250 Table 1c: Effect of crude oil concentration on the organic carbon content of Ido - Ani soil

Time (Days)	SS <sub>5</sub>	SS <sub>5A</sub>	SS <sub>5B</sub>	SS <sub>5C</sub>	SS <sub>5D</sub>
0	2.79 ± 0.01 <sup>a</sup>	3.77 ± 0.02 <sup>a</sup>	4.37 ± 0.01 <sup>a</sup>	5.02 ± 0.02 <sup>a</sup>	6.05 ± 0.01 <sup>a</sup>
30	3.29 ± 0.01 <sup>b</sup>	4.20 ± 0.02 <sup>b</sup>	4.47 ± 0.01 <sup>b</sup>	5.15 ± 0.01 <sup>b</sup>	6.23 ± 0.01 <sup>b</sup>
60	3.78 ± 0.02 <sup>c</sup>	4.50 ± 0.01 <sup>c</sup>	4.77 ± 0.06 <sup>c</sup>	5.49 ± 0.01 <sup>c</sup>	6.65 ± 0.01 <sup>c</sup>
90	3.91 ± 0.00 <sup>d</sup>	4.63 ± 0.02 <sup>d</sup>	5.03 ± 0.02 <sup>d</sup>	5.71 ± 0.01 <sup>d</sup>	6.93 ± 0.01 <sup>d</sup>
120	4.10 ± 0.01 <sup>e</sup>	4.85 ± 0.02 <sup>e</sup>	5.24 ± 0.05 <sup>e</sup>	5.92 ± 0.02 <sup>e</sup>	7.17 ± 0.01 <sup>e</sup>
150	4.29 ± 0.01 <sup>f</sup>	5.22 ± 0.01 <sup>f</sup>	5.69 ± 0.01 <sup>f</sup>	6.45 ± 0.01 <sup>f</sup>	7.83 ± 0.07 <sup>f</sup>
180	4.45 ± 0.05 <sup>g</sup>	5.83 ± 0.12 <sup>g</sup>	6.55 ± 0.01 <sup>g</sup>	7.27 ± 0.01 <sup>g</sup>	8.21 ± 0.01 <sup>g</sup>

251 Legend:

252 SS<sub>5</sub> = 0% Contamination (control)

253 SS<sub>5A</sub> = 1% Crude Oil Contamination

254 SS<sub>5B</sub> = 2% Crude oil Contamination

255 SS<sub>5C</sub> = 3% Crude Oil Contamination

256 SS<sub>5D</sub> = 4% Crude Oil Contamination

257

## 258 DISCUSSION

259 It has been reported that the effects and time for reclamation of polluted soil depend on  
 260 the quantity and concentration of the pollutant [13, 14]. A thorough knowledge of the impact  
 261 of oil pollution on the physicochemical properties of the soil as a technological parameters  
 262 for its elimination is very critical. In this study, the effects of varying concentrations of crude  
 263 oil on some physicochemical properties of soil were evaluated. Contamination of the three  
 264 arable experimental soil samples of Igodan-Lisa, Oba- Ile and Ido- Ani at 1-4% (w/w) caused  
 265 alterations in the physicochemical properties of the soils. This finding is in line with the



266 findings of [7, 20] who stated that oil spills cause alterations in the physicochemical and  
267 microbiological properties of soils.

268 The results of varying concentrations of crude oil on some physicochemical properties of  
269 the agricultural soils revealed observable changes in pH, contents of moisture, organic matter  
270 and organic carbon. All of these parameters are significant in determining soil quality and  
271 also influence the efficiency of bioremediation as strategy for hydrocarbon pollutant removal.  
272 The degree of acidity and alkalinity of soil is a very important property affecting many other  
273 physicochemical and biological properties and can as well be used as index to assess soil  
274 quality and suitability of the environment for bioremediation of polluted soil.

275 In this study, results revealed that the pH status of the polluted soils varied and extent of  
276 this depended on the concentration of spilled oil. The values of pH in the control samples  
277 ranged 7.03- 7.28, suggesting that the pH of the control samples were almost neutral or  
278 slightly alkaline compared to the acidic pH of 4.91 - 6.17 obtained for crude oil contaminated  
279 soil samples at 1- 4% (w/w) contamination levels. The observed decrease in pH which  
280 implies increased acidity agrees with the reports of [22, 23] who observed increased acidity  
281 as following increased crude oil pollution of soil. The decrease in pH with increase in levels  
282 of oil in soil samples, however, deviates from the reports of [18, 24] who observed increase  
283 in pH as the level of pollution increased. The decrease in pH with increase in the amount of  
284 crude oil used in the treatments implies that at all levels of crude oil contamination, the pH of  
285 the samples were altered becoming more acidic as concentration and study period increased.  
286 This finding agrees with the reports of [21, 25, 26] who ascribed the progressive decrease in  
287 pH of crude oil polluted soil with time to the accumulation of acidic metabolites resulting  
288 from microbial degradation or metabolism of the spilled soils. A pH of 6.5-7.5 is considered  
289 optimum for the growth of many plants [5]. Oyem and Oyem [5] reported that pH affects

290 plant growth primarily by its effects on nutrient availability and that high or low pH causes  
291 deficiencies of essential nutrients that plants need to grow. The results of this study therefore  
292 show that the effect of 1-4% oil pollution of soil is a fall in pH below the limit favorable for  
293 plant and crop growth and survival. Therefore, plants growth in this adverse pH condition  
294 may be stunted in growth for reasons of deficiencies of nutrients and may as well be more  
295 prone to disease and fungal attack and consequently the destruction of vegetation. Also, pH  
296 affects microbial activities, growth and survival. Different microbial strains exhibit their  
297 maximum growth potentials in a limited pH range [27]. The values of pH obtained in this  
298 study for crude oil polluted soils fall below the optimum pH (6.0 – 8.0) for microbial growth  
299 and bioremediation of crude oil polluted soil [1]. This implies that the efficiency of soil  
300 microbes in breaking down organic pollutants will be limited or slow.

301 All soil microorganisms require moisture for growth and other metabolic activities. The  
302 effective transport of soluble nutrients, food and waste metabolic products in and out of the  
303 microbial cells depends on available moisture. In the present study, the moisture content  
304 (MC) of crude oil polluted soils decreased with increase in the level of pollution. The  
305 moisture content (MC) of polluted soils reduced compared with the control samples (figures  
306 2a - c). The observation in this study supports the reports of [19, 26, 28]. Essien and John  
307 [28] asserted that moisture content per unit weight of soil sample was less in crude oil  
308 polluted soils than in unpolluted (control) soil samples. The reduction in MC of polluted soil  
309 was ascribed to coating of the soil surface by hydrophobic hydrocarbon that reduces the water  
310 holding capacity of the soil and reduction in the binding property of clay soil [29]. The  
311 progressive increase in moisture content with increased sampling days may also be attributed  
312 to insufficient aeration of the soil that might have arisen from the displacement of air in the  
313 soil; this probably encouraged water logging and reduced rate of evaporation [21]. Also, the

314 increase in MC with time may be the result of degradation by microorganisms during which  
315 organic compounds in crude oil are converted to carbon dioxide and water as products of  
316 microbial degradation and therefore suggesting reclamation of the crude oil spilled soils.

317 The data from this study revealed appreciable increases in organic matter contents  
318 following the increase in the level of applied petroleum oil against the control soils, thereby  
319 agreeing with the reports of [30, 31, 32] who reported a surge in organic matter content of  
320 contaminated soil. The observed response to increase in the level of crude oil spilled was  
321 thereafter followed by a remarkable gradual decrease in the percentage organic matter with  
322 time (figures 3a - c). The continual decrease in organic matter content of contaminated soil  
323 might have resulted from crude oil mineralization by native microbial population. This  
324 research revealed that the percentage organic carbon in crude oil contaminated soils were  
325 higher than in uncontaminated control soils. This is in line with the findings of [33] that  
326 pollution of sandy loam soil by crude oil led to an increase in soil organic carbon. This  
327 increase in organic carbon also agrees with the report of [23] which suggested that crude oil  
328 pollution adversely affects the ecosystem through the provision of excess carbon that might  
329 be unavailable for microbial use. The increased organic carbon will consequently create  
330 nutritional imbalance especially of carbon and nitrogen since crude oil contains large amounts  
331 of carbon-containing compounds. The progressive increase in carbon content over the study  
332 period could also be attributed to the accumulation of organic acids resulting from  
333 degradation. Osuji and Nwoye [21] however, reported a slightly lower total organic carbon  
334 and total organic matter in polluted soils than in the control. Their report asserted that severe  
335 hydrocarbon contamination is indicated by high soil acidity (low pH) and high MC, low TOC  
336 and TOM all implying low soil fertility which in turn implies low agricultural productivity  
337 and reduced source of livelihood in the affected area.

338 The impacts of crude oil pollution on agricultural lands have generated great concern  
339 among the people living in the oil producing areas of Niger Delta region where residents  
340 depend solely on sales from farms as means of livelihood as well as government and  
341 environmental researchers. The hitherto agricultural lands have become wastelands and  
342 unproductive for profitable agriculture. On the basis of the results obtained in this study, it  
343 can be concluded that crude oil pollution no matter the quantity and size (minor, medium,  
344 major and catastrophic) caused impaired changes in physicochemical properties of soil,  
345 thereby destroying soil integrity and quality and hence, agricultural productivity. The extent  
346 of this depended on the amount of crude oil spill. The pH of soil samples was reduced below  
347 the limit favorable for plant and microbial growth and bioremediation. Hence, crude oil  
348 polluted soil could be limed. It is suggested to deal with crude oil spillage than dealing with  
349 the consequences. This implies the need for increased public awareness on the prospective  
350 environmental consequences of crude oil spill and enforcement of regulatory environmental  
351 laws. Prompt response in term of contingency fund to meet the needs and concerns of those  
352 affected by spill and application of appropriate remediation strategies to oil spilled sites is  
353 recommended in order to restore soil quality and improve agriculture and the socio-  
354 economic lives of the residents of the Niger – Delta areas where crude oil spill is a common  
355 occurrence.

356

## 357 REFERENCES

- 358 1. Odeyemi, O. Two centuries of oil and gas (1860-2060) [www.universalacademicservices.org](http://www.universalacademicservices.org)  
359 2014
- 360 2. Ibitoye, A. A.. Laboratory manual on basic soil analysis (2<sup>nd</sup> ed). Foladvae Publishing  
361 Company, Akure, Nigeria. 2006

- 362 3. Delgado, A, Gomez, J. A. The Soil, Physical, Chemical and Biological Properties. In: F. J.  
363 Villalobos, E. Fereres (eds). Principles of Agronomy for sustainable Agriculture. Springer  
364 International Publishing. AG 2016: 15-26.
- 365 4. European Commission. Soil international year of soils, Director – General for environment,  
366 [env-soil@ec.europa.eu](mailto:env-soil@ec.europa.eu) 2015
- 367 5. Oyem, I. L. R., Oyem, I. L.. Effects of crude oil spillage on soil physicochemical properties  
368 in Ugborodo Community. *International Journal of Modern Engineering Research*, 2013;  
369 3(6): 3336-3342
- 370 6. Santhaveerana ,G, Mahendra, B. G.. Influence of soil characteristics on bioremediation of  
371 hydrocarbons contaminated soil. *Journal of Environmental Research and Development*,  
372 2010; 4(3): 734 – 741
- 373 7. Sharma, S., Pathak, H.. Physicochemical properties of petroleum contaminated soil collected  
374 from Arid region of Rajasthan (Churu). *International Journal of Pharma and Bio Sciences*,  
375 2017; 8(2):  
376 (B) 926 – 932.
- 377 8. Agarry, S. E., Ogunleye, O. Factorial designs application to study enhanced  
378 bioremediation of soil artificially contaminated with weathered Bonny light  
379 crude oil through biostimulation and bioaugmentation strategy. *Journal of*  
380 *Environmental Protection* 2012; 3: 748-759
- 381 9. Chorom, M., Shariffi, H . S., Mutamedi, H. Bioremediation of a crude – oil polluted soil by  
382 application of fertilizers. *Iran Journal of Health Science Engineering*, 2010; 7(4): 319 -326  
383

- 384 10. Digha, O. N., Ambah, B., Jacob, E. N. 2017. The effect of crude oil spillage on farmland in  
385 Gokana Local Government Area of Rivers State. *European Journal of Basic and Applied*  
386 *Sciences*, 4(1), 76-96
- 387 11. Ijah, U. J. J., Safiyanu, H., Abioye, O. P. Comparative study of Biodegradation of crude oil  
388 in soil amended with chicken Droppings and NPK fertilizer. *Science World Journal*, 2008;  
389 3(2): 63-67.
- 390 12. Jain, P. K., Gupta, V. K., Gaur, R. K., Lowry, M., Jaroli, D. P., Chauhan, U. K..  
391 Bioremediation of Petroleum Oil Contaminated Soil and Water. *Research. Journal of*  
392 *Environmental Toxicology*, 2011; 5(1): 1-26
- 393 13. Ikpeme, E. M., Nfongeh, J. F., Etim, L. Comparative bioremediation enhancement  
394 procedures on kerosene polluted utisol from Niger Delta Region, Southern Nigeria. *Research*  
395 *Journal of Microbiology*, 2007; 2(11): 826-860.
- 396 14. Onuoha SC, Chukwura EI, Fatokun K. Stimulated biodegradation of spent lubricating motor  
397 oil in soil amended with animal droppings. *American Journal of BioScience*, 2014; 2(1): 19-  
398 27.
- 399 15. Marinescu M, Dumitru M, Lăcățusu A, Marinescu M. Evolution of Maize Biomass in a  
400 crude oil polluted soil according to the treatment applied. *Scientific papers, UASVM.*  
401 *Bucharest, Series A, Liv.* 2011; 287-292.
- 402 16. Ikuesan FA, Boboye BE, Adetuyi FC. The microbiological and physicochemical properties  
403 of some crude oil contaminated and uncontaminated agricultural soils in Ondo State, Nigeria.  
404 *Pyrex Journal of Microbiology and Biotechnology Research*, 2016;2(1): 1 – 8
- 405 17. Ekpo MA, Ebeagwu CJ. The effect of crude oil on microorganisms and dry matter of fluted  
406 pumpkin (*Telfairia occidentalis*) *Scientific Research and Essays*, 2009 ;4(8): 733 -739

- 407 18. Njoku KL, Akinola MO, Oboh BO. Phytoremediation of crude oil contaminated Soil. The  
408 effect of Growth of *Glycine max* on the physico-chemistry and crude oil contents of soil.  
409 *Nature and Science*, 2009; 7(10): 79 – 87
- 410 19. Akubugwo EI, Ogbuji GC, Chinyere CG, Amadike EB. physicochemical properties and  
411 enzymes activities studies in a refined oil contaminated soil in Isiukwuato, Abia State,  
412 Nigeria. *Nigeria Society for Experimental Biology*, 2009; 21(2): 79-84
- 413 20. Ijah UJJ, Abioye OP. Assessment of physicochemical and microbiological properties of soil  
414 30 months after kerosene spill. *Journal of Research in Science and Management*, 2003; 1:  
415 24-30.
- 416 21. Osuji LC, Nwoye I. An appraisal of impact of petroleum hydrocarbons on soil fertility: the  
417 Owaza experience. *African Journal of Agricultural Research*. 2007; 2(7): 318-324
- 418 22. Amadi EN, Okolo JC, Odu, CTI. Optimizing crude oil degradation in a sandy soil: Effects of  
419 urea – nitrogen and phosphoric acid. *J. Niger Environ. Soc.* 2005; 2(3): 322 – 329.
- 420 23. Obasi NA, Eze E, Anyanwu DI, Okorie UC. Effects of organic manures on the  
421 physicochemical properties of crude oil polluted soils. *Academic Journal of Biochemistry*  
422 *Research*, 2013; 7(6): 67-75.
- 423 24. Onuh MO, Madukwe DK, Ohia GU. Effects of poultry manure and cow dung on the  
424 physical and chemical properties of crude oil polluted soil. *Science World Journal*, 2008;  
425 3(2): 45-50.
- 426 25. Merkl N, Schutze-Kraft R, Infante C. Assessment of tropical grasses and legumes for  
427 phytoremediation of petroleum contaminated soils, *Water, Air and Soil pollution* , 2005 ;165  
428 (1-4): 195-209.

- 429 26. Barua D, Buragohain J, Sarma SK. Certain physiochemical changes in the soil brought about  
430 by contamination of crude oil in two oil fields of Assam, NE. India. *European Journal of*  
431 *Experimental Biology*, 2011;1(3): 154 - 161
- 432 27. Pathak H, Bhatnagar K, Jaroli DP. Physico-chemical properties of petroleum polluted soil  
433 collected from transport Nagar (Jaipur) *Indian Journal of Fundamental and Applied life*  
434 *Sciences*. 2011; 1(3): 84-89
- 435 28. Essien OE, John IA. Impact of Crude Oil Spillage Pollution and Chemical Remediation on  
436 Agricultural Soil Properties and Crop Growth. *J. Appl. Sci. Environ. Management*, 2010;  
437 14(4): 147-154
- 438 29. Dibble J, Bartha R. Effects of Environmental parameters on biodegradation of oil sludges.  
439 *Applied Environmental Microbiology*, 1989; 37: 729 – 730
- 440 30. Etuk EA, Ogboi KC, Nwadinigwe CA. Bioremediation of hydrocarbon polluted soil in the  
441 lowland forest ecosystem in the Niger Delta through enhanced Natural Attenuation Process  
442 (ENAP). *International Journal of Applied Science and Technology*, 2013;3(8): 128 – 137
- 443 31. Obayori OS, Ilori MO, Adebusoye SA, Amund OO, Oyetibo GO. Microbial population  
444 changes in tropical agricultural soil experimentally contaminated with crude oil. *African*  
445 *Journal of Biotechnology*, 2008; 7(24); 4512 – 4520.
- 446 32. Shukry WM, Al-Hawas GHS, Al-Moaikal RMS, MA El-Bendary Effect of petroleum crude  
447 oil on mineral nutrient elements, soil properties and bacterial biomass of the Rhizosphere of  
448 Jojoba. *British Journal of Environment and Climate Change*, 2013; 3(1): 103 – 118.
- 449 33. Okolo JC, Amadi EN, Odu CTI. Effects of soil treatments containing poultry manure on  
450 crude oil degradation in sandy loam soil. *Appl. Biol. Environ. Res.*, 2005; 3(1):  
451 47-53