

Original Research Article

PHYSICO-CHEMICAL QUALITY RESPONSES OF MANGO CHIPS DRIED TO DIFFERENT MOISTURE CONTENTS, PACKAGED AND STORED FOR SIX MONTHS

ABSTRACT

Fresh mango is a delicate fruit with high perishability and postharvest losses. Dried fruits are highly susceptible to mold infection and moisture reabsorption, must be packed well and stored immediately after drying to extend its shelf life. This study was thus to determine the effect of packaging methods on the quality of mango chips dried to three moisture content and stored for six months at tropical ambient condition. Mango chips (10%, 15% and 20% moisture contents) obtained from FriutProtech Consortium, Kintampo, were packed in different packages and stored for 6 months in a 3×5 factorial in completely randomized design and replicated 3 times. Physico-chemical attributes were studied. Results revealed that mango chips at 10% moisture content, vacuum-packed in polypropylene and polyethylene were significantly different ($p \leq 0.01$), the driest, firmest and most crispy. They also were richest in vitamin C, high in pH, highest in TSS and but high TTA. Chips at 20% moisture content in plastic pack (clamshells) were lowest in most of the measured parameters. It can be concluded that, mango chips should be dried to, at most, 15% moisture content and vacuum-packaged in polypropylene or polyethylene if they are to maintain their quality and be stored for longer periods.

Keywords: {perishability, crispy, antioxidants, hydrolysis, ripening, enzyme}

1. INTRODUCTION

Mango (*Mangifera indica* L.) fruit belongs to the family of Anacardiaceae and is grown in many parts of the world, particularly in tropical countries. According to [1], mango fruit has been reported the 2nd position as a tropical crop, behind only bananas in terms of production and acreage used. It has been well indicated that mango fruits are highly nutritious. [2], indicated that mangoes are essential to human growth, development and health. Mangoes form a 50 per-cent share of all the tropical fruits produced worldwide. Within the horticultural sector, mango cultivation can easily become a major foreign exchange earner if well developed and provided with the necessary logistics and support.

Due to high postharvest losses of fruits and vegetables, there is the need to process and preserve perishable fruits during bumper harvest to make them available throughout the year in a value-added form. Dehydrated mango fruits slices could be processed from the glut by individuals or farmer-groups to address the vitamin A and C problems experienced in certain part of the country. According to [3] vitamin A deficiency is a major public health problem in Ghana especially in the northern part of the country in children under 5 years of age. [4], reported that mango fruit is climacteric with a high rate of perishability and after harvesting at ambient conditions reaches the peak of its ripening process on the third (3rd) to fourth (4th) day. According to [5], the longevity is between four to eight days at room storage (130^oC). [6] also reported that the fruit is highly sensitive to decay, low temperature and its perishability caused by rapid ripening and softening reduce

Comment [h1]: Therefore, the fruits must be packaged well...

Comment [h2]: ...3 times before the physiochemical attributes were studied.

Comment [h3]: Vitamin C. The 'C' is always in capital letter. Please adjust it.

Comment [h4]: ...and or but high in TTA

31 the storage, handling and transport potential. Mango fruits have been processed into chips and other forms to extend the
32 shelf-life and improve its commercial potentials through value-addition. Dried fruits are highly susceptible to mold infection
33 and moisture reabsorption and must be well- packaged and stored immediately after drying to maintain its quality [7]. The
34 type of packaging material used has been reported to have effect on nutrient content during storage [8]. Packaging is an
35 inevitable component of food processing, for assuring the safe handling and delivery of fresh and processed agricultural
36 products from producer to the consumer [9]. It is therefore important to assess the packaging method to ascertain their
37 performance in ensuring product quality during storage. Postharvest management is essential for extending the
38 consumption period of fruits, for regulating their supply to the market and for transporting them over long distances. The
39 objective of the study was to determine the effect of packaging methods (sealed polyethylene, vacuumed polyethylene,
40 sealed polypropylene, vacuumed polypropylene and plastic pack-clamshell) on the physico-chemical properties of mango
41 chips dried to three different moisture contents (10%, 15% and 20%) and stored for six (6) months at tropical ambient
42 temperature.

43 44 **2.0 MATERIALS AND METHODS**

45 **2.1 STUDY SITE**

46 The study was conducted at the laboratories of the Department of Horticulture, Faculty of Agriculture, and the Department
47 of Theoretical and Applied Biology, Kwame Nkrumah University of Science and Technology (KNUST), Kumasi.

48 **2.2 EXPERIMENTAL MATERIAL**


49 The following materials were used in the experiment: dried mango chips at three different moisture contents (10%, 15%
50 and 20%), five different packaging methods [sealed polyethylene (0.085mm thick) (PolyEthyseal), vacuumed polyethylene
51 (0.085mm thick) (PolyEthovac), sealed polypropylene (0.055mm thick) (PolyPropseal), vacuumed polypropylene
52 (0.055mm thick) (PolyPropvac) and plastic pack (clamshell) (PP)], vacuum machine, sealer, a pair of scissors, top-loading
53 electronic balance.

54 **2.3 SOURCE OF MANGO CHIPS AND PRODUCTION**

55 The mango chips were obtained from the FriutProtech Consortium at Kintampo.

56 **2.3.1 Mango Chips Production**

57 Keitt variety of mango fruits at a stage of fully ripe were weighed and washed thoroughly under running water after which
58 they were reweighed. The mango fruits were then peeled using a sharp knife and cut into two equal halves and the seeds
59 removed. The pulp was then cut into slices (approximately 0.5cm thick) and dried in a solar-LPG hybrid oven at between
60 65^oC- 70^oC for at most 8 hours. Moisture contents of mango chips were tested periodically to ensure that the three
61 moistures (10%, 15% and 20%) were achieved.



63 Plate 1: Bulk mango chips at different moisture contents packaged at FruitProtech consortium, Kintampo before being
64 sent to the laboratory for further studies.

65 **2.4. SAMPLE PREPARATION**

66 Forty (40) grams each of dried mango chips (10%, 15% and 20%) were weighed into each of the five different packaging
67 methods [sealed polyethylene (0.085mm thick) (PolyEthyseal), vacuumed polyethylene (0.085mm thick) (PolyEthyvac),
68 sealed polypropylene (0.055mm thick) (PolyPropseal), vacuumed polypropylene (0.055mm thick) (PolyPropvac) and
69 plastic pack (clamshell) (Plasticpack)], stored and observed for six (6) months under tropical ambient conditions (27°C
70 temperature at 70% relative humidity).



71
72 Plate 2: Vacuuming of packaged mango chips in vacuum machine.
73



74
75 Plate 3: Differently processed, packaged and labeled mango chips stored at ambient conditions (L-R: sealed polyethylene
76 (0.085mm thick) (PolyEthyseal), vacuumed polyethylene (0.085mm thick) (PolyEthyvac), sealed polypropylene (0.055mm
77 thick) (PolyPropseal), vacuumed polypropylene (0.055mm thick) (PolyPropvac) and plastic (clamshell) pack
78 (PLASTICPACK)).

79 **2.5 EXPERIMENTAL DESIGN**

80 A 3×5 Factorial Completely Randomized Design (CRD) was used in the experiment. The three-different moisture content
81 (10%, 15% and 20%) against five packaging methods [sealed polyethylene (0.085mm thick) (PolyEthyseal), vacuumed
82 polyethylene (0.085mm thick) (PolyEthyvac), sealed polypropylene (0.055mm thick) (PolyPropseal), vacuumed
83 polypropylene (0.055mm thick) (PolyPropvac) and plastic pack (clamshell) (PLASTICPACK)]. Each treatment was
84 replicated three times

85 **2.6. PHYSICO-CHEMICAL PROPERTIES OF MANGO CHIPS DETERMINED.**

86 **2.6.1 Determination of moisture content (%)**

87 Moisture content was determined by following the procedures of AOAC, 2005 [10].

88 **2.6.2 Firmness determination (N)**

89 Pieces of mango chips of each treatment were tested using a type-C digital durometer (LX-C durometer, China) and
90 readings recorded initially and subsequently, monthly.

91 **2.6.3 Vitamin C determination**

92 Vitamin C was determined by following the procedures of AOAC, 2005 [10].

93 **2.6.4 pH determination**

94 -Five grams of blended oven dried sample was weighed into a 50ml beaker. Twenty-five (25) ml of distilled water was
95 added and stirred vigorously for 20 minutes. Sample water suspension was allowed to stand for 30 minutes by which time
96 most of the suspended ions would have settled out from the suspension. A pH meter-ELICO (L1617) was calibrated blank
97 at pH of 7 and 4 respectively. The electrode of the pH meter was inserted into the partly settled suspension. The pH value
98 was read from the pH meter and recorded.

99 **2.6.5 Total titratable acid determination**

100 TTA was determined by following the procedures of AOAC, 2005 [10].


101 **2.6.6 Total soluble solids (°Brix) determination**

102 The total soluble solid was determined by using HANNA refractometer (HI9680). Before determining the sugar content,
103 the refractometer was first calibrated using 25% sucrose solution and distilled water. A 10ul of the prepared sample
104 solution was placed on the prism surface of the refractometer. The reading on the prism scale was noted and recorded to
105 one decimal in degree Brix [11].

106 **3.0 RESULTS**

107 **3.1 Changes in moisture content during storage**

108 There was significant difference ($P \leq 0.01$) between the moisture content over the storage period. Moisture reduced
109 consistently over the storage period except for month five (5) where 15% moisture content increased and decrease the
110 following month (Figure 1).



111
112 Figure 1: Changes in moisture content of mango chips (10%, 15% and 20% moisture contents) during a six (6) months
113 storage period

114 There was significant difference ($P \leq 0.01$) between different packaging methods on mango chips during storage period.
115 Moisture content was reduced by PolyEthylvac throughout the storage period (Figure 2)




Figure 2: Moisture content of mango chips in different packages during a six (6) months storage period.

There were significant difference in ($P \leq 0.01$) moisture content and different packaging methods interaction for changes in moisture during six (6) months storage period (Table 1). Mango chips processed at 10% moisture content and packaged using Polyethylene vacuum recorded the highest (13.76) moisture content and the least (3.27) was recorded by mango chips processed at 20% moisture content and packaged using plastic pack (clamshell). Across the different packaging methods, Polyethylene vacuum recorded the highest (12.79) and plastic pack (clamshell) recorded the least (4.89) moisture. Across the moisture content, highest (10.53) moisture content was recorded at 20% moisture content and the least (9.47) was recorded by moisture content at 10%.

Table 1: Means of moisture content (%) of mango chips dried to different moisture content and packaged differently using different methods and stored for six (6) months.

Moisture content of chips (%)	Packaging Methods					Means
	Plastic _{pack}	PolyProp _{seal}	PolyProp _{vac}	PolyEth _{seal}	PolyEth _{vac}	
10	3.27 ⁱ	8.52 ^h	11.05 ^{def}	10.77 ^{efg}	7.46 ⁱ	8.21 ^b
15	3.93 ^j	10.52 ^{fg}	11.38 ^{cde}	10.17 ^d	12.79 ^b	9.76 ^b
20	13.76 ^a	11.24 ^{cdef}	10.02 ^d	12.09 ^c	11.82 ^{cd}	11.81 ^a
Mean	6.99 ^d	10.09 ^c	10.82 ^b	10.98 ^a	10.69 ^b	

CV=2.21 HSD(0.01) mc=0.25 pm=0.37 mc*pm=0.77

*Means followed by the same letter are not significantly different at $p=0.01$

3.2 Changes in firmness during storage

There was significant difference ($P \leq 0.01$) between firmness and moisture content from month one to month four (Figure 3). Mango chips processes at 10% moisture content recorded the highest firmness and moisture content at 20% recorded the least. In month five and month six, moisture content at 10% recorded the highest firmness and the least, moisture content at 15%.




Figure 3: Firmness of mango chips processed at three moisture content over six (6) months storage period.

There was significant difference ($P \leq 0.01$) between Firmness and the different packaging methods (Figure 4). Vacuum packaging increases hardness hence scored highest firmness while Polyethylene seal packaging recorded least for firmness throughout storage periods except for month five (5) where plastic pack (clamshell) recorded the highest (43.61) firmness and Polyethylene seal (33.78) recorded the least firmness.




Figure 4: Firmness of mango chips in different packages stored over storage period.

There were significant ($P \leq 0.01$) moisture content and different packaging methods interaction for Firmness during six (6) months storage period (Table 2). Mango chips processed at 10% moisture content and packaged using Polypropylene vacuum recorded the highest (12.67) firmness and the least (1.73) was recorded by mango chips processed at 20% moisture content and packaged using Polyethylene vacuum. Across the different packaging methods, Polypropylene vacuum recorded the highest (9.38) and plastic pack (clamshell) recorded the least (7.89) firmness. Across the moisture content, highest (11.67) firmness was recorded at 10% moisture content and the least (3.63) was recorded by moisture content at 20%.


Table 2: Means of Firmness (N) of mango chips dried to different moisture content and packaged differently using different methods and stored for six (6) months.

Moisture content of chips (%)	Packaging Methods					Means
	Plastic _{pack}	PolyProp _{seal}	PolyProp _{vac}	PolyEthy _{seal}	PolyEthy _{vac}	
10	11.77 ^b	11.23 ^d	12.67 ^a	11.20 ^d	11.50 ^c	11.67 ^a
15	9.70 ^f	9.77 ^f	10.17 ^e	9.50 ^g	11.77 ^b	10.18 ^b
20	2.20 ^k	5.70 ^h	5.30 ⁱ	3.20 ^j	1.73 ^l	3.63 ^c
Mean	7.89 ^d	8.90 ^b	9.38 ^a	7.97 ^d	8.33 ^c	
CV =0.63	HSD (0.01)	mc=0.06	pm=0.90	mc*pm=0.19		

*Means followed by the same letter are not significantly different at $p=0.01$


151 **3.3 Vitamin C (mg/100ml) content during the storage period.**

152 There were no significant differences in the Vitamin C content for mango chips processed at moisture content level in the
 153 early months of storage (Figure 5.). Meanwhile, in month four, mango chips processed at 10% moisture content had the
 154 highest vitamin c (3.48) and moisture content at 20% recorded the least (2.62). Mango chips processed at (10%) moisture
 155 content, maintained a higher vitamin C content while moisture content at 20% lost vitamin C content considerably
 156 throughout the storage period.



157 Figure 5: Vitamin C content of mango chips processed at three moisture content over storage period.

159 There were significant differences ($P \leq 0.01$) Vitamin C content between the different packaging methods during storage
 160 (Figure 6). Polyethylene vacuum recorded the highest vitamin c (9.70) and Polypropylene vacuum recorded the least
 161 (5.20) in month one. In subsequent months, vitamin c reduced with time yet mango chips packaged in Polyethylene seal
 162 packaging methods recorded the highest vitamin c (7.30) compared to Polyethylene vacuum which recorded the least
 163 (2.86) in the final month.



164 Figure 6: Vitamin C content of mango chips in different packages stored over storage period.

167 There were significant ($P \leq 0.01$) packaging methods and moisture content interaction for Vitamin C during six (6) months
 168 of storage (Table 3). Highest Vitamin C (8.61) was indicated by mango chips processed at 10% moisture content and
 169 packaged using Polyethylene vacuum. The least (1.67) was recorded by mango chips proceeded at 20% moisture content
 170 and packaged using plastic pack (clamshell). Across the moisture content, moisture content at 10% recorded the highest
 171 (6.00) and moisture content at 15% recorded the least (4.81). Across the different packaging method, Polyethylene seal
 172 recorded the highest (6.80) and plastic pack (clamshell) recorded the least (2.76).

173 Table 3: Means of Vitamin c (mg/100ml) of mango chips dried to different moisture content and packaged using different
 174 methods and stored for six (6) months.

Moisture content of chips (%)	Packaging Methods					Means
	Plastic _{pack}	PolyProp _{seal}	PolyProp _{vac}	PolyEthyl _{seal}	PolyEthyl _{vac}	
	4.31 ^{de}	5.16 ^{cd}	5.36 ^c	6.56 ^b	8.61 ^a	6.00 ^a

10						
15	2.30 ^f	5.41 ^c	5.43 ^c	6.74 ^b	4.15 ^e	4.81 ^b
20	1.67 ^f	4.83 ^{cde}	5.53 ^c	7.11 ^b	5.57 ^c	4.94 ^b
Mean	2.76 ^d	5.13 ^c	5.44 ^c	6.80 ^a	6.11 ^b	

CV =5.30 HSD (0.01). mc=0.32 pm=0.47 mc*pm=1.0

*Means followed by the same letter are not significantly different at p=0.01

3.4 Acidity (pH) of mango chips during storage.

There were no significant differences in pH for mango chips processed at different moisture content) during storage (Figure 7). Mango chips processed at 10% moisture content recorded highest pH (3.70) while mango chips processed at 20% moisture content increased with time and recorded least pH (2.77) at the later stage of storage.




Figure 7: pH of mango chips processed at three moisture content over storage period.

There were significant differences ($P \leq 0.01$) between the packaging methods over the storage period (Figure 8). Mango chips packaged using Polypropylene seal recorded the highest (3.33) pH and the least (3.22) was recorded by mango chips packaged using Polypropylene vacuum at the initial stage of storage. Meanwhile in the final month of storage, Polyethylene seal packaging method recorded the highest pH (4.38) and Polyethylene vacuum recorded the least (1.45)




Figure 8: pH of mango chips in different packages stored over storage period.

There were significant ($P \leq 0.01$) moisture content and different packaging methods interaction for acidity (pH) during six (6) months of storage (Table 4). Mango chips processed at 20% moisture content and packaged using Polyethylene seal recorded the highest (5.93) pH and the least (0.89) was recorded by mango chips processed at 20% moisture content and package using plastic pack (clamshell). Across the packaging methods, highest (5.83) pH was recorded by mango chips packaged using Polyethylene seal and the least (1.19) was recorded by the plastic pack (clamshell) packaging method. Across the moisture, highest (4.67) pH between the moisture content was recorded at 10% and the least (4.36) at 15%.

194 Table 4: pH of mango chips dried to different moisture content and packaged using different methods and stored for six
 195 (6) months.


Moisture content of chips (%)	Packaging Methods					Means
	Plastic _{pack}	PolyProp _{seal}	PolyProp _{vac}	PolyEthy _{seal}	PolyEthy _{vac}	
10	1.78 ^{g*}	4.47 ^e	5.51 ^d	5.72 ^{bc}	5.85 ^{ab}	4.67 ^a
15	0.91 ^h	5.77 ^{abc}	5.63 ^{cd}	5.83 ^{ab}	3.69 ^f	4.36 ^c
20	0.89 ^h	5.70 ^{bc}	5.92 ^a	5.93 ^a	3.81 ^f	4.45 ^b
Mean	1.19 ^e	5.31 ^c	5.69 ^b	5.83 ^a	4.45 ^d	

CV =1.18 HSD (0.01). mc=0.06 pm=0.09 mc*pm=0.19

196 *Means followed by the same letter are not significantly different at p=0.01


197 **3.5 Total Titratable Acid (TTA) Content During Six (6) Storage Periods.**

198 There were no significant differences ($P \leq 0.01$) in total titratable acid for mango chips at different moisture content during
 199 six (6) months of storage (Figure 9). Mango chips processed at 20% moisture content recorded highest (0.29) total
 200 titratable acid and lowest total titratable acid (0.11) of chips were recorded by moisture content at 15%. At the sixth (6)
 201 month of storage, mango chips processed at 10% and 15% moisture content recorded the highest total titratable acid
 202 (0.11) while mango chips processed at 20% moisture content recorded the least total titratable acid (0.08).



203
 204 Figure 9: Total titratable acid of mango chips processed at three moisture content over storage period.

205 There was significant difference ($P \leq 0.01$) in total titratable acid for mango chips packaged at different packaging methods
 206 during storage (Figure 10). Plastic pack (clamshell) recorded the highest total titratable acid (1.10) and Polyethylene seal
 207 recorded the least (0.67) at the beginning of storage. Polyethylene seal recorded the highest (0.14) total titratable acid
 208 and Polyethylene vacuum recorded the least (0.05) in the last month.



209

210 Figure 10: Total titratable acid content of mango chips in different packages stored over storage period.

211 There were significant ($P < 0.01$) moisture content and different packaging method interaction for Total titratable acidity
 212 over the six (6) storage period (Table 5). Highest total titratable acid (0.53) was recorded by mango chips processed at
 213 10% moisture content and packaged using Polyethylene vacuum. The least (0.14) was recorded by mango chips
 214 processed at 10% moisture content and packaged using Polypropylene seal. Across the moisture content, moisture
 215 content at 10% recorded the highest (0.32) total titratable acid and the least (0.29) by 15% moisture content. Across
 216 packaging method, highest (0.37) total titratable acid was recorded by Polypropylene vacuum. The least (0.23) was
 217 recorded by plastic pack (clamshell).

218 Table 5: Means of Total titratable acid (%) of mango chips dried to different moisture content and packaged using different
 219 methods and stored for six (6) months.


Moisture content of chips (%)	Packaging Methods					Means
	Plastic _{pack}	PolyProp _{seal}	PolyProp _{vac}	PolyEth _{seal}	PolyEth _{vac}	
10	0.27 ^{de*}	0.14 ^f	0.32 ^{cd}	0.32 ^{cd}	0.53 ^a	0.32 ^a
15	0.19 ^f	0.34 ^c	0.34 ^c	0.34 ^c	0.25 ^e	0.29 ^b
20	0.24 ^e	0.28 ^{de}	0.46 ^b	0.32 ^{cd}	0.25 ^e	0.31 ^a
Mean	0.23 ^c	0.25 ^c	0.37 ^a	0.33 ^b	0.34 ^b	

CV =5.19 HSD (0.01). mc=0.2 pm=0.3 mc*pm=0.06

220 *Means followed by the same letter are not significantly different at $p=0.01$

221 **3.6 Total soluble solids (^oBrix) during storage period.**

222 There were significant differences in Total soluble solids for mango chips processed at different moisture content over the
 223 storage period (Figure 11). Mango chips processed at 15% moisture content recorded highest total soluble solids (11.77)
 224 and mango chips processed at 10% moisture content recorded lowest total soluble solids (9.47) in the initial stage of
 225 storage. Total soluble solids reduced at the later stage of storage. Processed chips at 10% moisture content recorded
 226 highest total soluble solids (9.53) and mango chips processed at 20% moisture content recorded least (6.93)




227

Figure 11: Total soluble solids of mango chips processed at three moisture content over storage period.

228

229 Total soluble solids of Mango chips packaged using different packaging methods declined and increased during storage
 230 (Figure 12). Polyethylene seal packaging method lost the highest (12.29) total soluble solids to Polypropylene seal in
 231 month six (6) yet Polyethylene vacuum maintained the least (4.19) total soluble solid.



232

Figure 12: Total soluble solids content of mango chips in different packages stored over storage period.

233

234 There were significant ($P \leq 0.01$) packaging methods and moisture content interaction for Total soluble solids during six (6)
 235 months storage (Table 6). Highest total soluble solids (10.86) were recorded by mango chips processed at 10% moisture
 236 content and packaged using Polypropylene seal. The least (1.87) was recorded by mango chips processed at 20%
 237 moisture content and packaged using plastic pack (clamshell). Across the different packaging methods, highest (9.98)
 238 total soluble solids was recorded by Polypropylene seal and the least (2.96) by plastic pack (clamshell). Across the
 239 moisture content, processed mango chips at 10% moisture content recorded the highest (8.91) total soluble solids and
 240 least (7.09) by moisture content at 20%.

241 Table 6: Means of Total soluble solids ($^{\circ}$ Brix) of mango chips dried to different moisture content and packaged using
 242 different methods and stored for six (6) months.

Moisture content of chips (%)	Packaging Methods					Means
	Plastic _{pack}	PolyProp _{seal}	PolyProp _{vac}	PolyEthy _{seal}	PolyEthy _{vac}	
10	4.97 ^h	10.86 ^a	9.98 ^{bc}	9.03 ^g	9.73 ^{cd}	8.91 ^a
15	2.05 ⁱ	9.73 ^{cd}	9.52 ^{de}	9.18 ^{fg}	4.98 ^h	7.13 ^b
20	1.87 ⁱ	9.35 ^{ef}	9.09 ^{fg}	10.07 ^b	5.24 ^h	7.09 ^b

Means	2.96 ^d	9.98 ^a	9.53 ^b	9.43 ^b	6.65 ^c
CV =0.19 HSD (0.01).		mc=0.09	pm=0.14	mc*pm=0.29	

*Means followed by the same letter are not significantly different at p=0.01

4.0 DISCUSSIONS

4.1 Moisture content

There were significant ($P \leq 0.01$) moisture content and different packaging methods interaction on changes in moisture during six (6) months storage period. This result illustrated a general decrease in moisture over the storage period. Also, different packaging method, had a significant effect on the moisture content of mango chip due to the fact that, packaging method had a moisture reabsorption barrier properties (PolyEthylvac). According to [12], dried products absorb moisture from its surrounding which increases moisture content, decreasing shelf-life. Deterioration and chemical reactions could be higher in plastic pack (clamshell) with higher moisture content (20%) due to the proliferation characteristics of the packaging method during storage time [12].

4.2 Firmness

Low moisture content (10%) makes the chips quite brittle and crispy while the higher moisture (20%) renders it flabby. Mango chips processed at 10% moisture content and packaged using polypropylene vacuum recorded the highest firmness and the least was recorded by mango chips processed at 20% moisture content and packaged using polyethylene vacuum. This suggests that moisture content at which the chips were processed had significant impact on firmness. According to [13], firmness is the result of complex interactions among food components at micro- and macro-structural levels. Again, [14] indicated that during drying of fruits and vegetables, several changes in texture and firmness are common (e.g.; hardness, cohesiveness, springiness and chewiness). The results of the present study corroborate their findings.

4.3 Vitamin C content during the storage period.

Results indicated that vitamin C content reduced when chips were processed at 20% moisture content and packaged using plastic pack (clamshell) while those processed at 10% moisture content and packaged using polyethylene vacuum increased. Furthermore, higher reduction in Vitamin C content occurred when storage duration increased. This could be due to the fact that the vacuum polyethylene methods prevented exchange of gases between the dried chips at 10% moisture content and the storage environment. However, for the plastic pack (clamshell), there was a possibility of gas exchange between the chips and the environment which enhanced rapid oxidation of abundant Vitamin C leading to its breakdown. Oxidation becomes faster when dried products absorb moisture at higher temperature. According to [15] oxidation causes the disruption of the cell membrane leading to the release of membrane bound phytochemicals. The presence of oxygen could also initiate the conversion of vitamin C to dehydroascorbic acid and other oxidized products. [16] indicated that light has a significant influence on the stability of vitamin C during storage. This could also be a contributory factor for the loss in vitamin C in the plastic pack (clamshell). The results of the present study agree with findings of [17] who reported that the degradation rate of ascorbic acid (Vitamin C) of dried tomato pulp increased with high temperature, longer storage period and higher moisture content. Reduction in the vitamin C could be attributed to the fact that, increasing moisture content increases water activity a condition suitable for oxidative degradation of vitamins C as noted by (1995) [18] and [8]. Vitamin C is sensitive to air, light and heat.

4.4 pH during the storage period.

Results showed that pH of dried mango chips processed at 20% moisture content and stored in Polyethylene seal was the highest and the least was recorded by mango chips processed at 20% moisture content and packaged using plastic pack (clamshell) probably due to the effect of organisms responsible for the spoilage, some of which can release basic substances into the samples. This corroborates with the work of [19] who reported that certain organisms were responsible for spoilage by releasing basic substances into food products. The plastic pack (clamshell) undoubtedly

284 enhanced rapid absorption of moisture by the chips creating a conducive environment for microorganisms to proliferate.
285 Furthermore, [20] also explained that the pH values of vegetables and fruits being weakly acidic allow growth of certain
286 microorganisms. The food acidity (pH) is an important parameter in food. Food acidity not only affects flavor but also the
287 growth and survival of bacteria and other microorganism in foods. Water ionizes as temperature rises, so hydrogen ion
288 concentration rises which means that pH decreases.

289 4.5 Total titratable acidity during the storage period.

290 Titratable acidity gives a measure of the amount of acid present in a fruit [21]. Major acid in mango is known to be caused
291 by Citric acid [22]. The decline in acidity by mango chips processed at 10% moisture content and packaged using
292 polypropylene seal was probably due to [23]. Again, the reduction in acidity in ripening process was as a result of starch
293 hydrolysis which led to a rise in total sugars and a decline in acidity as opined by [24]. Chips produced in this study had
294 reduced sourness with potential taste improvement as a result of decreased acidity.

Comment [h5]: Due to what property?

295 4.6 Total soluble solids during the storage period.

296 Highest total soluble solids were recorded by mango chips processed at 10% moisture content and packaged using
297 Polypropylene seal. The least was recorded by mango chips processed at 20% moisture content and packaged using
298 plastic pack (clamshell). This could be due to the oxidation of dried mango chips kept in plastic pack (clamshell) and
299 processed at 20% which possibly absorbed moisture from its environment thereby increasing the breakdown or hydrolysis
300 of sugars. As the storage time increased this rate of sugar hydrolysis was also increased thereby reducing the available
301 sugars in the product. A study done by [25], showed that mango pulps are mainly made up of fructose, with about 30%
302 sucrose and 20% glucose. In addition, sucrose is known to be the major sugar in mango [26]. The high increase in total
303 soluble solids during ripening is reported to be the major cause of significant increase in the amount of sucrose. This
304 could probably due to the conversion of starch to soluble sugars as the carbohydrates in the fruit are broken down into
305 simple sugars with the action of phosphorylase enzyme during ripening [27]. Conversely, the amylase enzyme is closely
306 associated with hydrolysis of starch in the ripening of mango fruit. The extent of sweetening was due to the increase in
307 total soluble solids during ripening [28].

308 **COMPETING INTERESTS**

309
310 "Authors have declared that no competing interests exist."

Comment [h6]: This section should come after the conclusion. Please adjust accordingly.

313 5.0 CONCLUSION

314 The study revealed that mango chips processed at 10% moisture content and vacuum packaged using polypropylene
315 were and recommended drier, firm and crispy during the storage period. For vitamin C, mango chips processed at 10%
316 moisture content and vacuum packaged using polyethylene had the highest vitamin C content while mango chips
317 processed at 20% moisture content and packaged using plastic pack (clamshell) had the least. Mango chips processed at
318 20% moisture content and packaged in plastic pack (clamshell) had the least pH and mango chips processed at 20%
319 moisture content and seal packaged using polyethylene recorded the least. Mango chips processed at 10% moisture
320 content and packaged in polypropylene sealed had the highest total soluble solids while mango chips processed at 20%
321 moisture content and packaged using plastic pack (clamshell) had the least. Total titratable acid of mango chips
322 processed at 10% moisture content and packaged in polypropylene seal had the lowest and highest recorded by 10%
323 vacuumed in polyethylene.

324 **REFERENCES**

- 325
326
327 1. Muchiri D R, Mahungu SM, Gituanja SN. Studies on mango (*Mangifera indica*, L.) kernel fat of some Kenyan
328 varieties in Meru. *Journal of the American Oil Chemists' Society*. 2012. 89(9): 1567-1575.
329 2. Tharanathan RN, Yashoda HM, Prabha TN. Mango (*Mangifera indica* L.), "The king of fruits"—An overview. *Food*
330 *Reviews International*. 2006; 22(2): 95-123.

- 331 3. Benamba ER. Nutrition in Northern Ghana. *Vitamin A deficiency. The Northern Health Monitor*. 2005. 4(1): 25.
- 332 4. Narayana CK, Pal RK, Roy SK. Effect of pre-storage treatments and temperature regimes on shelf-life and
- 333 respiratory behavior of ripe Baneshan mango. *Journal of food science and technology*. 1996; 33(1): 79-82.
- 334 5. Carillo Lopez A, Ramirez-Bustamante F, Valdez-Torres JB, Rojas-Villegas R, Yahia EM. Ripening and quality
- 335 changes in mango fruit as affected by coating with an edible film. *Journal of Food Quality*. 2000 23(5): 479-486.
- 336 6. Hoa TT, Ducamp MN, Lebrun M, Baldwin EA. Effect of different coating treatments on the quality of mango fruit.
- 337 *Journal of food quality*. 2002; 25(6): 471- 486.
- 338 7. Chassery S, Gormley TR. Quality and shelf life of pre-peeled vacuum-packed potatoes. *Farm and Food*.1994;
- 339 4(2): 30-32.
- 340 8. Salunkhe DK, Bolin HR, Reddy NR. *Storage, processing, and nutritional quality of fruits and vegetables. Volume I.*
- 341 *Fresh fruits and vegetables* (No. Ed. 2). CRC press. 1991.
- 342 9. Mditshwa A, Fawole OA, Al-Said F, Al-Yahyai R, Opara, UL. Phytochemical content, antioxidant capacity and
- 343 physicochemical properties of pomegranate grown in different microclimates in South Africa. *South African*
- 344 *Journal of Plant and Soil*. 2013; 30(2): 81-90.
- 345 10. Association of Official Analytical Chemists (AOAC) Official Methods of Analysis. 2005.
- 346 11. Hrazdina G, Parsons GF, Mattick LR. Physiological and biochemical events during development and maturation
- 347 of grape berries. *American Journal of Enology and Viticulture*.1984 35(4): 220-227.
- 348 12. Nyame B. *Quality of keitt mango chips as affected by method of drying, packaging and storage periods* (MPhil
- 349 dissertation, KNUST). 2015.
- 350 13. Guihe L Jian G, Cuijun Y, Nutritional evaluation of fresh and wilted mixed silage of naked oats (*Avena nuda*) and
- 351 alfalfa (*Medicago sativa*). *International Journal of Agriculture and Biology*, 2015. 17(4).
- 352 14. Kotwaliwale N, Bakane P, Verma A. Changes in textural and optical properties of oyster mushroom during hot air
- 353 drying. *Journal of Food Engineering*. 2007. 78(4): 1207-1211.
- 354 15. Lemmens L, Van Buggenhout S, Oey I, Van Loey A, Hendrickx M. Towards a better understanding of the
- 355 relationship between the β -carotene in vitro bio-accessibility and pectin structural changes: a case study on
- 356 carrots. *Food Research International*. 2009. 42(9): 1323-1330.
- 357 16. Alzamora SM, Gómez PL, Castro MA, Salvatori DM. Effect of ultraviolet-C light dose on quality of cut-apple:
- 358 Microorganism, color and compression behavior. *Journal of Food Engineering*. 2010. 98(1): 60-70.
- 359 17. Goula AM, Adamopoulos KG, Chatzidakis PC, Nikas VA. Prediction of lycopene degradation during a drying
- 360 process of tomato pulp. *Journal of Food Engineering*. 2006. 74(1): 37-46.
- 361 18. Baldwin EA, Nisperos-Carriedo MO, Baker RA. Use of edible coatings to preserve quality of lightly (and slightly)
- 362 processed products. *Critical Reviews in Food Science and Nutrition*. 1995. 35(6): 509-524.
- 363 19. Garangyo T, Thiombiano T, Werem A, Drawara B, Sawadogo L. Etude de l'impact du sechage solaire direct sur
- 364 la teneur en vit c de la tomato. *Science Technology*. 1992. 20, 53-61.
- 365 20. Matazu IK, Umar RA, Ebbo AA, Akpeji AJ, Hassan SW. Hepatoprotective effect of leaf extracts of *Parkinsonia*
- 366 *aculeata* L. against CCl 4 intoxication in albino rats. *Int J Biol Chem*. 2008. 2(2): 42-48.
- 367 21. Dadzie BK, Orchard JE. *Routine post-harvest screening of banana/plantainhybrids: criteria and methods* (Vol. 2).
- 368 Bio-versity International. 1997.
- 369 22. Ihekoronye AI, Ngoddy PO. *Integrated food science and technology for the tropics*. Macmillan. 1985.
- 370 23. Aina JO. Physico-chemical changes in African mango (*Irvingia gabonensis*) during normal storage ripening. *Food*
- 371 *Chemistry*. 1990. 36(3): 205-212.
- 372 24. Fuchs Y, Pesis E, Zauberman G. Changes in amylase activity, starch and sugars contents in mango fruit pulp.
- 373 *Scientia Horticulturae*. 1980. 13(2): 155-160.
- 374 25. Faure P, Corticelli P, Richard MJ, Arnaud J, Coudray C, Halimi S, Roussel AM. Lipid peroxidation and trace
- 375 element status in diabetic ketotic patients: influence of insulin therapy. *Clinical Chemistry*. 1993. 39(5): 789-793.
- 376 26. Gil AM, Duarte IF, Barros A, Delgadillo I, Almeida C. Application of FTIR spectroscopy for the quantification of
- 377 sugars in mango juice as a function of ripening. *Journal of agricultural and food chemistry*. 2002. 50(11): 3104-
- 378 3111.
- 379 27. Mitra SK, Baldwin EA. Mango. *Postharvest physiology and storage of tropical and subtropical fruits*. 1997. Pp 85-
- 380 122.
- 381 28. Jha SN, Chopra S, Kingsly ARP. (Determination of sweetness of intact mango using visual spectral analysis.
- 382 *Biosystems engineering*. 2005. 91(2): 157-161.