

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

**Effects of Different Types of Organic Fertilizers on Growth Performance of
Amaranthus caudatus (Samaru Local Variety) and *Amaranthus cruentus*
(NH84/452)**

31

ABSTRACT

32
33
34
35
36
37
38
39
40
41
42
43
44
45

To evaluate the effect of different types of organic fertilizers on growth performance of *Amaranthus caudatus* (Samaru local variety) and *Amaranthus cruentus* (NH84/452). A randomized complete block design (RCBD) was used for the experiment. The field experiment was carried out in the nursery of a homestead garden at No 20, Isaiah Balat Street, Sabo GRA, Kaduna State, Nigeria. The study consists of seven treatments which includes control (no fertilizer), 5 tons/ha and 10 tons/ha poultry manure, 5 tons/ha and 10 tons/ha sewage sludge, 35kg/ha and 70kg/ha NPK compound fertilizer and also with *Amaranthus caudatus* (Samaru local variety) and *Amaranthus cruentus* (NH84/452) in factorial arrangement fitted into a randomized complete block design (RCBD) and replicated three times. Growth performance data were collected on plant height, number of leaves, leaf length, leaf width, leaf area and leaf area index from 2 weeks after transplanting (WAT) to 6 weeks after transplanting (WAT). The plant height and number of leaves of the two varieties were found in the range of 18.30-135.67cm and 13.33-78.33cm respectively. Leaf area and leaf area index of the two varieties had values in the range of 41.71-258.29cm² and 1.76-41.72 respectively. At 6WAT, 10tons/ha poultry manure recorded the highest value for all the growth parameters for both varieties except for leaf length, leaf width and leaf area of *Amaranthus caudatus* (Samaru local variety), where 10 tons/ha sewage sludge and 70kg/ha NPK compound fertilizer were highest. The experimental results of this study have shown that poultry manure had higher growth performance on the two varieties of Amaranth when compared with sewage sludge and NPK compound fertilizer. The application of poultry manures at 10 tons/ha is therefore recommended for farmers to use to obtain higher yields of Amaranth.

31
32

Keywords: Growth, Organic and Inorganic fertilizers, Amaranth, Soil and Insecticides

33

1. INTRODUCTION

34
35
36
37
38
39
40
41
42
43
44
45

Increasing population of the world has doubled the food demands and inundated the available land resources. Alongside other food alternatives, vegetables are considered cheap source of energy (11). Vegetables are rich sources of essential biochemicals and nutrients such as carbohydrates, carotene, protein, vitamins, calcium, iron, ascorbic acid and palpable concentration of trace minerals (28). Amaranth has been one of the most important vegetables of Amaranthaceae family. Amaranth has been naturalized in central parts of Asia and possibly Iran (14) and has cultivation history of more than 2000 years (6). Cultivation of the various *Amaranthus* species is acquiring increasing importance in Nigeria and other parts of African continent where the available species are grown for their leaves (4) Organic and inorganic fertilizers are essential for plant growth as it supplies plants with the nutrients needed for optimum performance. Organic fertilizer has been used for many centuries whereas chemically synthesized inorganic fertilizers were only widely developed during the industrial revolution. Inorganic fertilizers have significantly supported global population growth, as it has been estimated that

46 almost half the people on the earth are currently fed as a result of artificial nitrogen fertilizer use (10).
47 Commercial and subsistence farming has been and is still relying on the use of inorganic fertilizers for
48 growing crops (18). This is because they are easy to use, quickly absorbed and utilized by crops. The
49 continued dependence of developing countries on inorganic fertilizers has made prices of many
50 agricultural commodities to skyrocket (18).
51 Moreover, most vegetable farmers in tropical Africa are small holders who cannot afford the cost of
52 inorganic fertilizers, although soil fertility limits yield of vegetables especially in urban centres (17). In
53 Nigeria, fertilizers, being costly and sometimes scarce can make farmers not apply enough for good
54 growth (4). Fertilizer application rates in intensive agricultural systems have increased drastically during
55 recent years in Nigeria. Farmers depend largely on locally sourced organic fertilizers (17). In Nigeria,
56 huge amount of organic wastes such as poultry waste, animal excreta, sewage sludge, refuse soil and
57 palm oil mill effluent are generated and heaped on dump sites, posing potential environmental hazard.
58 Incorporating these waste materials into the soil for crop production is expected to be beneficial to the
59 buildup of organic matter layer that is needed for a steady supply of nutrients by tropical soils (3).
60 Oyediji *et al.* (24) reported that NPK and poultry manure improved the growth and yield of three different
61 species of amaranth (*Amaranthus hybridus*, *Amaranthus deflexus* and *Amaranthus cruentus*) but
62 influenced proximate composition differently. Emede *et al.* (9) reported that poultry manure influenced the
63 plant growth and yield of *Amaranthus cruentus* L. positively. Therefore, the objective of this study was to
64 determine the effect of different types of organic fertilizers on the growth performance of *Amaranthus*
65 *caudatus* (Samaru local variety) and *Amaranthus cruentus* (NH84/452)

67 2. MATERIALS AND METHODS

68 2.1 Seeds

69 The seeds of *Amaranthus caudatus* (Samaru local variety) were obtained from local farmers in Samaru,
70 Zaria, Nigeria while the seeds of *Amaranthus cruentus* (NH84/452) were obtained from National
71 Horticultural Research Institute (NIHORT), Ibadan, Nigeria. *Amaranthus caudatus* samples collected were
72 authenticated at the herbarium unit of Biological Sciences Department, Ahmadu Bello University Zaria,
73 Nigeria and a voucher specimen was deposited.

74 2.2 Study Area

75 The field experiment was carried out in the nursery of a homestead garden at No 20, Isaiah Balat Street,
76 Sabo GRA, Sabo Tasha, Kaduna State, Nigeria. Kaduna metropolis has a tropical savanna climate with
77 dry winters characterized by maritime air and rainfall is between April and October with annual raining
78 days ranging from 81 to 103 mm. During the reference period, the annual mean rainfall values range from
79 145.37mm to 318.67mm. From the figures above, ample rains are available for the production of many
80 agricultural crops. During harmattan, dry desert wind blows between December and mid February while
81 night temperature is very low. The geographical location of Kaduna metropolis is Latitude $9^{\circ}03'N$ and
82 $11^{\circ}32'N$ north of the equator and Longitudes $6^{\circ}05' E$ and $8^{\circ}38' E$ East of the Greenwich meridian.
83 Kaduna metropolis has a sub-humid semi arid tropical climate with maximum annual mean temperature
84 ranging from $25.30^{\circ}C$ to $36.20^{\circ}C$ while the minimum annual mean temperature range of $28.45^{\circ}C$ to
85 $34.38^{\circ}C$ (2).

86 2.3 Soil sampling

87 Surface soil sample was taken from the experimental site at a depth of 0 – 15cm at land preparation (after
88 ploughing and harrowing) using the zigzag method. The sample was collected from twenty points and
89 bulked to form a composite sample. The composite sample was air-dried, crushed and sieved through a
90 2mm mesh sieve and stored for chemical analysis (19).

91 **2.4 Fertilizers**

- 92 i. Poultry manure: The poultry manure was collected at Ishaya's poultry farm in Sabo GRA,
93 Sabo Tasha Kaduna State, Nigeria.
- 94 ii. Sewage sludge: The dried packed sewage sludge was collected at the sewage site of
95 Ahmadu Bello University Zaria, Nigeria.
- 96 iii. NPK compound fertilizer: NPK compound fertilizer (15:15:15) was bought at Kawo market
97 Kaduna State, Nigeria.

98 **2.5 Soil analysis**

99 The sampled soil was analyzed at the Soil Science Department of the Institute of Agricultural Research,
100 Ahmadu Bello University Zaria, Nigeria. The following parameters were analyzed in the sampled soil;
101 particle size, pH (in water), organic carbon, available phosphorus, total nitrogen, cation exchange
102 capacity (CEC) and exchangeable bases (5).

103 **2.6 Experimental design and fertilizer treatment**

104 The experiment included seven (7) fertilizer treatments for each of the two (2) varieties of Amaranth which
105 are in factorial arrangement fitted into a randomized complete block design (RCBD) and replicated three
106 times. Hence, the experiment had a total of 42 experimental plots. The treatments were: Control (no
107 fertilizer), 5 tons/ha poultry manure, 10 tons/ha poultry manure, 5 tons/ha sewage sludge, 10 tons/ha
108 sewage sludge, 35kg/ha NPK compound fertilizer, 70kg/ha NPK compound fertilizer (19).
109

110 **2.7 Planting and nursery management**

111 Prior to planting, the amaranth seeds were soaked in water for about 24 hours in order to enhance
112 germination. The soaked seeds were first sown in the nursery of about 1.9 cm deep and were watered
113 twice daily. Appropriate nursery management practices were carried out as at when needed to obtain
114 healthy and uniform seedlings. The experimental site was ploughed, harrowed and prepared into slightly
115 raised beds (plots) of 25cm width × 80cm length dimension preparatory to transplanting the crop
116 seedlings. Poultry manure and sewage sludge were incorporated according to treatment level to specific
117 plots during land preparation, thoroughly mixed with the soil and then left for two weeks to allow for
118 mineralization. Half of the NPK Compound fertilizer was applied at day of transplanting while the balance
119 was applied one week later. After two weeks in the nursery, randomly picked seedlings were transplanted
120 to the well prepare beds (plots). The seedlings were watered twice daily using watering can and the
121 surrounding areas were weeded regularly. The experimental area and the surroundings were kept clean
122 to prevent harbouring of pests. Insects were controlled by using "Dime Force Insecticide" with
123 concentration of 1.5 L/ha (22).
124

125 **2.8 Data collection for growth performance**

126 Data were first collected two weeks after transplanting (WAT) and subsequently at one week interval for
127 up to six weeks after transplanting. Two randomly selected plants were tagged and used in each plot for
128 data collection. Data collected included plant height, number of leaves, leaf length and leaf width, while
129 the leaf area and leaf area index were computed (19).

130 **2.8.1 Determination of plant height**

131 Plant height is the length of the plant from the base of the stem (surface of the soil) to the apex of the
132 leaves. Plant height was measured using a measuring tape for the two tagged plants per plot and the
133 average computed (19).

134 **2.8.2 Determination of number of leaves**

135 The number of leaves was counted from the two tagged plants and the average computed (19).

136 **2.8.3 Determination of leaf area**

137 The Leaf Area (LA) was computed by multiplying the Leaf Length (LL) by the Leaf Width (LW) and the
138 product multiplied by the correction factor (19).

139 Calculation;

140 Leaf Area = (Leaf Length × Leaf Width) 0.578.

141 **2.8.4 Determination of leaf area index**

142 The leaf area index (LAI) was computed using this formula (21)

$$143 \text{ LAI} = Y \times N \times \text{LA} \times (\text{AP})^{-1}$$

144 Where: Y = Population of plants per plot (5 plants), N = Average number of leaves, LA = Leaf area, AP =
145 Area of plot (25cm width * 80cm length = 2000cm²)

146

147 **2.9 Statistical analysis**

148 Data was analyzed using the Statistical Package for Social Sciences (SPSS) version 21.0 computer
149 package. Descriptive statistics was used to determine the measures of central tendency. Means were
150 separated using Duncan Multiple Range test. Values with different superscripts down the column are
151 significantly different at $p < 0.05$.

152

153 **3. RESULTS AND DISCUSSION**

154 **3.1 Soil analysis results**

155 Results of analyses of the soil used for this experiment are shown in Table 1. The texture class of the soil
156 is sandy clay loam in which sand was highest with value of $66 \pm 2.0\%$, followed by clay with $24 \pm 3.0\%$
157 and silt was the lowest with value of $10 \pm 1.0\%$. The soil organic carbon, total nitrogen and available
158 phosphorus were $0.46 \pm 0.02\%$, $0.32 \pm 0.01\%$ and $7.4 \pm 0.30\text{ppm}$ respectively. The exchangeable bases
159 of Sodium, magnesium, calcium, potassium and cation exchange capacity (CEC) contents were $0.34 \pm$
160 0.02Cmol/Kg , $0.84 \pm 0.02\text{Cmol/Kg}$, $3.26 \pm 0.05\text{Cmol/Kg}$, $0.65 \pm 0.03\text{Cmol/Kg}$ and $5.7 \pm 0.20\text{Cmol/Kg}$
161 respectively. Soil pH value was 7.7 ± 0.2 .

162

163 **3.2 Organic fertilizer analysis results**

164 Results of analyses of the organic fertilizers used for this experiment are shown in Table 2. Poultry
165 manure showed a pH of 7.62 ± 0.04 , while the concentrations for total nitrogen, available phosphorus and
166 potassium were found to be $3.53 \pm 0.02\%$, $0.71 \pm 0.05\%$ and $1.61 \pm 0.03\%$ respectively. Sewage sludge
167 pH was found to be 8.25 ± 0.09 ; the concentrations of total nitrogen, available nitrogen and potassium
168 were gotten as $2.44 \pm 0.03\%$, $0.97 \pm 0.02\%$ and $1.33 \pm 0.05\%$ respectively.

169

170

171 **3.3 Plant Height**

172 Plant height was significantly ($P = .05$) higher in plants derived from poultry manure treated plots applied
173 at 10 tons/ha treatment and lowest in plants derived from no fertilizer treatment plots for both *Amaranthus*
174 *caudatus* (Samaru local variety) and *Amaranthus cruentus* (NH84/452) as shown in Fig 1 and Fig 2. At 2
175 Weeks After Transplanting (2WAT), the plant height was $26.28 \pm 1.07\text{cm}$ and $45.97 \pm 0.88\text{cm}$ from poultry
176 manure applied at 10 tons/ha treatment for *Amaranthus caudatus* (Samaru local variety) and *Amaranthus*
177 *cruentus* (NH84/452) respectively, which was also consistently highest till maturity (6WAT) as shown in
178 Fig1 and Fig 2. The highest plant height for the two varieties of amaranth at 6WAT were both observed in
179 plants treated with 10tons/ha poultry manure which was significantly ($p < 0.05$) different from the other
180 treatments. This position was earlier reported by Egharevba and Ogbe (7) and Okokoh and Bisong (23).
181 The highest plant height exhibited by plants treated with 10tons/ha poultry manure might have been due
182 to the presence of the primary nutrients plus other minerals found in inorganic manure, and also it may be
183 probably due to favourable nutrient mineralization of poultry manure as a result of the influence of the
184 mineral component on the organic content of the manure (1). The control plants had the lowest height as
185 they had to depend mainly on the intrinsic soil fertility as exhibited by the soil chemical analysis to be low.

186 A similar effect for control was reported for *Amaranthus caudatus* by Abayomi and Adebayo (1) and on
187 radish stems amaranth-indian spinach by Islam *et al.* (12). The height of the plant is an important growth
188 character directly linked with the productive potential of plants. An optimum plant height is claimed to be
189 positively correlated with productivity of plants (27).

190 **3.4 Number of leaves**

191 The number leaves were highest for plants treated with 10tons/ha poultry manure for both varieties of
192 amaranth and were not significantly ($P= .05$) different among the treatments except for the plants in the
193 control group as shown in Fig 3 and Fig 4. At 6WAT, poultry manure applied at 10tons/ha gave the
194 highest number of leaves with values of 78.67 ± 5.03 and 64.50 ± 3.50 for *Amaranthus caudatus* (Samaru
195 local variety) and *Amaranthus cruentus* (NH84/452) respectively. At the start of the experiment, the
196 average number of leaves was highest for poultry manure and NPK compound fertilizers for both varieties
197 of Amaranth. However, between 2WAT and 4WAT, the highest development of new leaves was observed
198 in 70kg/ha NPK compound fertilizer but not significantly different from 35kg/ha NPK and poultry manure.
199 Relatively high content of nitrogen in the NPK compound fertilizer increase the growth and development
200 of new leaves. Normally inorganic fertilizer nutrients are soluble, so the nitrogen was quickly released into
201 the soil leading to fast leaf growth and development. Although, during maturity leaf development declined
202 because the nutrients were probably exhausted in the soil; however, the reason for the high number of
203 leaves for plants treated with poultry manure compared to the sewage sludge at the early stages was
204 attributed to the high amount of nitrogen in the poultry manure than sewage sludge from chemical
205 analysis, and also due to faster mineralization and release of nutrients from the poultry manure than
206 sewage sludge. At maturity, the 10tons/ha poultry manure showed the highest average number of leaves
207 for both varieties of Amaranth, which was also reported by Law-Ogbomo and Ajayi (16) for *Amaranthus*
208 *cruentus*. This also agrees with reports by previous workers such as Sanwal *et al.*(29) in turmeric
209 (*Curcuma longa*); Premesekhar and Rajashree (26) in Okra (*Abelmoschus esculentus*) who separately
210 attributed higher leaf yield to released nutrients from organic manure application which improved
211 chemical, physical and biological properties of soil. This high leaves development in the poultry manure
212 compared to the sludge is due to the higher amount of nitrogen in poultry manure and continuous release
213 of the nutrients. However, the reason behind the higher number of leaves for plants treated with organic
214 fertilizers than the NPK compound fertilizer may be due to availability of nutrients as affected by the water
215 holding capacity of the soil (13). Most probably because as the manure quantities increased the water
216 holding capacity of the soil and subsequent nutrient release increases, while the NPK compound fertilizer
217 nutrients have been exhausted as the early stages due to the solubility of the nutrients.

218

219 **3.5 Leaf Area**

220 At maturity, leaf area which is a measure from the leaf length and leaf width was significantly ($P= .05$)
221 higher in plants derived from plots treated with 10 tons/ha sewage sludge with area of $127.36 \pm 3.40\text{cm}^2$
222 and 10 tons/ha poultry manure with area of $258.29 \pm 23.96\text{cm}^2$ for *Amaranthus caudatus* (Samaru local
223 variety) and *Amaranthus cruentus* (NH84/452) respectively and was lowest in plants derived from plots
224 with no fertilizer (control treatment) as shown in Fig 5 and Fig 6. Leaf area were found to be highest in
225 plants from the 70kg/ha NPK compound fertilizer for green type and the 10tons/ha poultry manure
226 recorded the highest for *Amaranthus cruentus* (NH84/452) but there no significant ($P= .05$) difference
227 among the treatments except for the control treatment. Similar work also reported by Mshelia and Degri
228 (20) on effect of different levels of poultry manure on performance of *Amaranthus caudatus* L. Okokoh
229 and Bisong (23) reported similarly in a research in Calabar that application of poultry manure significantly
230 influenced performance of amaranth. The increase in leaf area had been claimed to be directly influence
231 by nitrogen supply in fertilizer applied (8). The insignificant difference among the treatments suggests that
232 the nutrients in both the organic and inorganic fertilizers increases leaf width but higher amount of
233 nutrients in individual fertilizers may not necessary influence noticeable difference in the width of the
234 plants.

235 **3.6 Leaf Area Index**

236 Leaf area index which indicates the photosynthetic ability of the plants was significantly ($P= .05$) higher in
 237 plants derived from plots treated with 10 tons/ha poultry manure for both *Amaranthus caudatus* (Samaru
 238 local variety) and *Amaranthus cruentus* (NH84/452) with values of 23.74 ± 1.96 and 41.72 ± 5.48
 239 respectively and was lowest for plants derived from plots with no fertilizer added for both varieties as
 240 shown in Fig7 and Fig 8. The 10tons/ha poultry manure treatment resulted in the highest leaf area index
 241 for both varieties of amaranth which is consistent with report on red lettuce (18). The positive effect of
 242 poultry manure increasing leaf area index of amaranth was earlier reported by Egharevha and Ogbe, (7).
 243 Law-Ogbomo and Ajayi (16) also reported similar results on *Amaranthus cruentus*. Leaf area index and
 244 number of leaves follow the same pattern as both are directly related. The higher leaf area index in
 245 poultry manure was caused by the relatively higher nutrient availability which increased the leaf length,
 246 number of leaves and leaf width per unit area of the plot. Normally, inorganic chemical fertilizer nutrients
 247 are soluble, so the nitrogen was quickly released into the soil thus leading to fast leaf growth and
 248 development. However, during maturity, leaf development declined because the nutrients were probably
 249 exhausted in the soil. This resulted in the leveling of the leaf growth and development between NPK
 250 compound fertilizer, sewage sludge and poultry manure at maturity as sewage sludge and poultry manure
 251 was continuously releasing nitrogen. Organic manures like cattle manure and poultry manure have been
 252 reported to release both micro and macro nutrients slowly resulting in subsequent promotion of vegetable
 253 growth (30;25;15).

254

255

256

257 **Table 1: Physical and Chemical Properties of Soil used in this Experiment.**

Particulars	Value	Methods
Particle size		
• Clay	$24 \pm 3.0 \%$	USDA
• Silt	$10 \pm 1.0 \%$	USDA
• Sand	$66 \pm 2.0 \%$	USDA
Texture Class	Sandy Clay Loam	USDA
pH (in Water)	7.70 ± 0.20	
Organic Carbon	$0.46 \pm 0.02 \%$	Walkley-Black method
Available Phosphorus	7.40 ± 0.30 ppm	Bray and Kurts method
Total Nitrogen	$0.32 \pm 0.01 \%$	Kjeldahl method
Exchangeable bases		
• Calcium (Ca)	3.26 ± 0.05 Cmol/Kg	AAS
• Magnesium (Mg)	0.84 ± 0.02 Cmol/Kg	AAS
• Potassium (K)	0.65 ± 0.03 Cmol/Kg	AAS
• Sodium (Na)	0.34 ± 0.02 Cmol/Kg	AAS
• Cation Exchange Capacity (CEC)	5.70 ± 0.20 Cmol/Kg	Ammonium saturation

258 Values are mean \pm standard deviation of triplicate analysis.

259

260

261


262 **Table 2: Chemical Properties of Organic Fertilizers used in this Experiment**

Chemical Properties of the Organic Fertilizers used in the Experiment

	pH (in H ₂ O)	Total Nitrogen (%)	Available Phosphorus (%)	Potassium (%)
Poultry manure	7.62± 0.04	3.53 ± 0.02	0.71 ± 0.05	1.61± 0.03
Sewage sludge	8.25± 0.09	2.44 ± 0.03	0.97 ± 0.02	1.33 ± 0.05

263 Values are mean ± standard deviation of triplicate analysis.

264
265



266

267 **Fig 1: Effect of Organic Fertilizers on Plant Height of *Amaranthus caudatus* (Samaru local variety)**

268 Mean values ± standard deviation of triplicate analysis

269 **WAT**=Week after transplanting **Control**= No fertilizer **PM1**=Poultry manure at 5 tons/ha **PM2**=


270 Poultry manure at 10 tons/ha **SS1**= Sewage sludge at 5 tons/ha **SS2**= Sewage sludge at 10 tons/ha

271 **NPK1**= NPK compound fertilizer at 35kg/ha **NPK2**= NPK compound fertilizer at 70kg/ha


272

273


274




275
 276 **Fig 2: Effect of Organic Fertilizers on Plant Height of *Amaranthus cruentus* (NH84/452)**
 277 Mean values \pm standard deviation of triplicate analysis
 278 **WAT**=Week after transplanting **Control**= No fertilizer **PM1**=Poultry manure at 5 tons/ha **PM2**=
 279 Poultry manure at 10 tons/ha **SS1**= Sewage sludge at 5 tons/ha **SS2**= Sewage sludge at 10 tons/ha
 280 **NPK1**= NPK compound fertilizer at 35kg/ha **NPK2**= NPK compound fertilizer at 70kg/ha
 281
 282
 283




284
 285 **Fig 3: Effect of Organic Fertilizers on Number of Leaves of *Amaranthus caudatus* (Samaru local**
 286 **variety)**
 287 Mean values \pm standard deviation of triplicate analysis
 288 **WAT**=Week after transplanting **Control**= No fertilizer **PM1**=Poultry manure at 5 tons/ha **PM2**=
 289 Poultry manure at 10 tons/ha **SS1**= Sewage sludge at 5 tons/ha **SS2**= Sewage sludge at 10 tons/ha
 290 **NPK1**= NPK compound fertilizer at 35kg/ha **NPK2**= NPK compound fertilizer at 70kg/ha
 291
 292
 293
 294



295
 296 **Fig 4: Effect of Organic Fertilizers on Number of Leaves of *Amaranthus cruentus* (NH84/452)**
 297 Mean values \pm standard deviation of triplicate analysis
 298 **WAT**=Week after transplanting **Control**= No fertilizer **PM1**=Poultry manure at 5 tons/ha **PM2**=
 299 Poultry manure at 10 tons/ha **SS1**= Sewage sludge at 5 tons/ha **SS2**= Sewage sludge at 10 tons/ha
 300 **NPK1**= NPK compound fertilizer at 35kg/ha **NPK2**= NPK compound fertilizer at 70kg/ha
 301



302
 303 **Fig 5: Effect of Organic Fertilizers on Leaf Area of *Amaranthus caudatus* (Samaru local variety)**
 304 Mean values \pm standard deviation of triplicate analysis
 305 **WAT**=Week after transplanting **Control**= No fertilizer **PM1**=Poultry manure at 5 tons/ha **PM2**=
 306 Poultry manure at 10 tons/ha **SS1**= Sewage sludge at 5 tons/ha **SS2**= Sewage sludge at 10 tons/ha
 307 **NPK1**= NPK compound fertilizer at 35kg/ha **NPK2**= NPK compound fertilizer at 70kg/ha
 308
 309




310
311
312
313
314
315
316
317
318

Fig 6: Effect of Organic Fertilizers on Leaf Area of *Amaranthus cruentus* (NH84/452)

Mean values \pm standard deviation of triplicate analysis

WAT=Week after transplanting **Control**= No fertilizer **PM1**=Poultry manure at 5 tons/ha **PM2**=
Poultry manure at 10 tons/ha **SS1**= Sewage sludge at 5 tons/ha **SS2**= Sewage sludge at 10 tons/ha
NPK1= NPK compound fertilizer at 35kg/ha **NPK2**= NPK compound fertilizer at 70kg/ha




319
320
321
322
323
324
325
326
327
328

Fig 7: Effect of Organic Fertilizers on Leaf Area Index of *Amaranthus caudatus* (Samaru local variety)

Mean values \pm standard deviation of triplicate analysis

WAT=Week after transplanting **Control**= No fertilizer **PM1**=Poultry manure at 5 tons/ha **PM2**=
Poultry manure at 10 tons/ha **SS1**= Sewage sludge at 5 tons/ha **SS2**= Sewage sludge at 10 tons/ha
NPK1= NPK compound fertilizer at 35kg/ha **NPK2**= NPK compound fertilizer at 70kg/ha



329 **Fig 8: Effect of Organic Fertilizers on Leaf Area Index of *Amaranthus cruentus* (NH84/452)**
 330 Mean values \pm standard deviation of triplicate analysis
 331 **WAT**=Week after transplanting **Control**= No fertilizer **PM1**=Poultry manure at 5 tons/ha **PM2**=
 332 Poultry manure at 10 tons/ha **SS1**= Sewage sludge at 5 tons/ha **SS2**= Sewage sludge at 10 tons/ha
 333 **NPK1**= NPK compound fertilizer at 35kg/ha **NPK2**= NPK compound fertilizer at 70kg/ha
 334
 335

336 4. CONCLUSION

337 The rapidly rising cost of chemical fertilizers has forced small scale vegetative farmers to look for
 338 alternatives such as organic fertilizers; the result of this present study justifies the use of some organic
 339 fertilizers over chemical fertilizer due to high vegetative growth and plant development and increased
 340 level of nutrients but fails to justify based on elevated level of anti-nutrients and heavy metals seen in the
 341 organic fertilizer treatments. The present study revealed that application of poultry manure at 10 tons/ha
 342 on average effected the highest growth and development of *Amaranthus Amaranthus caudatus* (Samaru
 343 local variety) and *Amaranthus cruentus* (NH84/452) with highest values for plant height, number of
 344 leaves, leaf length, leaf width, leaf area and leaf area index. This effect can be as a result of the slow
 345 decomposition and release of nutrients from the organic fertilizers.

346 COMPETING INTERESTS

347 No competing interests

348 REFERENCES

- 349 1. Abayomi, O.A. and Adebayo, O.J. (2014). Effect of fertilizer types on the growth and yield of
 350 *Amaranthus caudatus* in Ilorin, Sothern Guinea, Savanna Zone of Nigeria. *Advances in*
 351 *Agriculture* 14:5
 352 2. Abdullahi, J., Shaibu-Imodagbe, E.M., Mohammed, F., Said, A. and Idris, U.D. (2009). Rural –
 353 Urban migration of the Nigerian work populace and climate change effects on food supply. A case
 354 study of Kaduna city in northern Nigeria.
 355 3. Agboola, A.A. and Omueti, J.A. (1982). Soil fertility problem and its management in the tropical
 356 Africa. Paper presented at the International Institute of Tropical Agriculture, Ibadan, Nigeria. pp
 357 25.

- 358 4. Alonge, S.O., Alonge, F.O., Bako, S.P., Olarewaju, J.D. and Adeniji, O.B. (2007). Effects of rates
359 and split application of compound NPK fertilizer on the growth and yield of three *Amaranthus*
360 species in Nigeria guinea savanna. *Asian J. Plant Sci.*, **6**: 906-912.
- 361 5. Burt R. (2014). Soil survey field and laboratory methods manual. *Soil survey investigations report*.
362 **2**(51): 181-387.
- 363 6. Daneshvar, H. (2000). Vegetable Farming. Chamran University Press Martyr, **190**: 461.
- 364 7. Egharevba, R.K.A. and Ogbe, F.M. (2002). The effects of different levels of organic and mineral
365 fertilizers on the yield performance of two *Amaranthus* (*A. cruentus*) cultivars. *The Plant*
366 *Scientists*. **3**:62-72.
- 367 8. Ehigiator, J. O. (1990). Farm yard manure: Need for its adoption as an alternative to chemical
368 fertilizer uses in Nigeria. *Nigerian Journal of Horticultural Science* **3**: 1– 9.
- 369 9. Emede, T.O., Law-Ogbomo, K.E. and Osaigbovo, A.U. (2012). Effects of poultry manure on the
370 growth and herbage yield of amaranth (*amaranthus cruentus* l.). *Nigerian Journal of Agriculture,*
371 *Food and Environment*. **8**(4):26-31.
- 372 10. Erisman, J.W., Sutton, M.A., Galloway, J., Klimont, Z. and Winiwarter, W. (2008). How a century
373 of ammonia synthesis changed the world. *Nat. Geosci.*, **1**: 636-639.
- 374 11. Hussain, J., Khan, A.L., Rehman, N., Hamayun, M., Shah, T., Nisar, M. and Lee, I. (2009).
375 Proximate and nutrient analysis of selected vegetable species: A case study of Karak region,
376 Pakistan. *African Journal of Biotechnology*, **8**(12).
- 377 12. Islam, M.M., Karim, A.J.M., Jahiruddin, M., Majid, N.M., Miah, M.G., Ahmed, M.M. and Hakim
378 M.A. (2011). Effects of organic manure and chemical fertilizers on crops in the radish-stem
379 amaranth-indian spinach cropping pattern in homestead area. *Australian journal of crop science*.
380 **5**(11):1370-1378.
- 381 13. Jacobs, R.D., Loan, D. and Jacob, J.(2003). Cage Layer Manure: An Important Resource for
382 Land use. Retrieved from: <http://edis.ifas.ufl.edu/PS005>, (Accessed on: May 19, 2009).
- 383 14. Kawazu, Y., Okimura, M., Ishii, T. and Yui, S. (2003). Varietal and seasonal differences in oxalate
384 content of spinach. *Scientia Horticulturae*, **97**(3), 203-210.
- 385 15. Kuntashula, E., Shileshi, G., Mafongoya, P.L. and Banda, J.(2006). Farmer participatory
386 evaluation of the potential for organic vegetable production in the wetlands of Zambia. *Outlook*
387 *Agr.*, **35**(4): 299-305.
- 388 16. Law-Ogbomo, K. A. and Ajayi, S. O. (2009). Growth and yield performance of *Amaranthus*
389 *cruentus* influenced by planting density and poultry manure application. *Natulae Botanicae Horti*
390 *Agrobotanici Cluj-Napoca* **37**(1): 195-199.
- 391 17. Makinde, E.A., Ayeni, L.S., Ojeniyi, S.O. and Odedina, J.N. (2010). Effect of organic,
392 organomineral and NPK fertilizer on nutritional quality of *Amaranthus* in Lagos, Nigeria.
393 *Researcher*, **2**: 91-96.
- 394 18. Masarirambi, M.T., Hlawe, M.M., Oseni, O.T. and Sibiya, T.E.(2010). Effects of organic fertilizers
395 on growth, yield, quality and sensory evaluation of red lettuce (*Lactuca sativa* L.) 'Veneza Roxa'.
396 *Agric. Biol. J. North America*, **1**(6): 1319-1324.
- 397 19. Masarirambi, M.T., Mbokazi, M.B., Wahame, P.K. and Oseni, T.O. (2012). Effect of kraal manure,
398 chicken manure and inorganic fertilizer on growth and yield of lettuce (*Lactuca sativa* L. var
399 Commander) in a semi-arid environment. *Asian Journal of Agricultural Sciences***4**(1): 56-64.
- 400 20. Mshelia, J.S. and Degri, M.M. (2014). Effect of different levels of poultry manure on the
401 performance of *Amaranthus caudatus* L. in Bama, Nigeria. *International journal of science and*
402 *nature*.**5**(1):121-125
- 403 21. Msibi, B.M., Mukabwe, W.O., Manyatsi, A.M., Mhazo, N. and Masarirambi, M.T. (2014). Effect of
404 liquid manure on growth and yield of spinach (*Beta vulgaris* var Cicla) in a sub-tropical
405 environment in Swaziland. *Asian Journal of Agricultural Sciences***6**(2): 40-47.
- 406 22. Musa, A., Ogbadoyi, E.O., Oladiran, J.A., Ezenwa, M.I.S. and Akanya, H.O. (2011). Effect of
407 fruiting on micronutrients, antinutrients and toxic substances in *Tilfairia occidentalis* grown in

- 408 Minna, Niger State, Nigeria. *African Journal of Environmental Science and Technology*. **5**(9):710-
409 716
- 410 23. Okokoh, S.J. and Bisong, B.W. (2011) Effect of Poultry Manure and Urea- N on Flowering
411 Occurrence and Leaf Productivity of *Amaranthus cruentus* in Calabar. *Journ.of Apl. Sic.*
412 *Environmental Management*. **15**(1):13-15.
- 413 24. Oyedeji, S., Animasaun, D.A., Bello, A.A. and Agboola, O.O. (2014). Effect of NPK and Poultry
414 Manure on Growth, Yield, and Proximate Composition of Three Amaranths. Hindawi Publishing
415 Corporation *Journal of Botany*. **10**: 11 – 55.
- 416 25. Pimentel, D., Hepperly, P., Hanson, J., Douds, D. and Sidel, R. (2005). Environmental, energetic
417 and economic comparisons of organic and conventional farming systems. *BioSci.*, **55**(7): 573-582.
- 418 26. Premsekhar, M. and Rajashree, V. (2009). Influence of organic manures on growth, yield and
419 quality of okra. *American-Eurasian Journal of sustainable Agriculture*, **3**(1): 6-8.
- 420 27. Saeed, I.N., Abbasi, K. and Kazim, M. (2001). Response of maize (*Zea mays*) to nitrogen and
421 phosphorus fertilization under agro-climatic condition of Rawalokot Azad Jammu and Kashmir.
422 *Pak. J. Biol. Sci.* **4**:53-55.
- 423 28. Salunkhe, D.K. and Kadam, S.S.(1998). Handbook of Vegetable Science and Technology:
424 Production, Composition, Storage and Processing. Marcel Dekker, INC, pp: 721.
- 425 29. Sanwal, S. K., Lakminaragana, K., Yadav, R. K., Yadav, D. S. and Mousumi, B. (2007). Effect of
426 organic manures on soil fertility, growth, physiology, yield and quality of turmeric. *Indian Journal*
427 *of Horticulture*, **64** (4): 444-449.
- 428 30. Van-Averbeke, W. and Yoganathan, S.(2003). Using Kraal Manure as Fertilizer. Department of
429 Agriculture Cape Town, University of Fort Hare, Republic of South Africa.