

Lung Cancer: A Chronic Disease Epidemiology; Prevalence Study.

Abstract

Chronic lung diseases (CLD) including asthma or chronic obstructive pulmonary disease (COPD) are a leading cause of morbidity and mortality worldwide and their occurrence in multiple sclerosis (MS) remains of interest. Increasing awareness of the possible adverse effect of CLD on outcomes in MS, such as disability progression and mortality, has heightened the need to understand the relationship between these chronic conditions. Prevalence of Lung Cancer was discussed in this paper, with intend to; Investigate the number of patients and deaths affected with lung cancer, test the effect of sex on lung cancer incidence, test the effect of environment and educational level on lung cancer incidence, examine the trend in lung cancer, and measure the relative risk associated with lung cancer. Secondary data sourced from the records units of five different hospitals was used. Cross tabulation, Chi-square test for independence, Regression Analysis, Correlation Analysis and Odds Ratio were applied on the three year study. From the study, it was found that lung cancer cases are independent on environmental factor, educational level and sex. A strong linear relationship exists between Lung Cancer and death from such disease, implying that increase in the number of lung cancer cases has very high positive effect on the occurrence of death ($r = 0.783$), 61.4% of the variation in death occurrence is explained by lung cancer. The probability of dying from lung cancer is higher in patients 50 years and above than in younger patients (age < 50 yrs).

Keywords: Lung cancer, chronic disease epidemiology, prevalence study, odds ratio, relative risk.

1. Introduction

Lung cancer, also known as lung carcinoma,^[1-3] is a malignant lung tumor characterized by uncontrolled cell growth in tissues of the lung. If left untreated, this growth can spread beyond the lung by process of metastasis into nearby tissue or other parts of the body^[4-5]. Most cancers that start in the lung, known as primary lung cancers, are carcinomas that derive from epithelial cells. The main primary types are small-cell lung carcinoma (SCLC) and non-small-cell lung carcinoma (NSCLC). The most common symptoms are coughing (including coughing up blood), weight loss, shortness of breath, and chest pains.^[6] The vast majority (85%) of cases of lung cancer are due to long-term exposure to tobacco smoke.^[7] About 10–15% of cases occur in people who have never smoked.^[8] These cases are often caused by a combination of genetic factors^[9] and exposure to radon gas,^[10] asbestos,^[11] or other forms of air pollution,^[12] including second-hand smoke.^[11] Lung cancer may be seen on chest radiographs and computed tomography (CT) scans. The diagnosis is confirmed by biopsy^[12] which is usually performed by bronchoscopy or CT-guidance. Treatment and long-term outcomes depend on the type of cancer, the stage (degree of spread), and the person's overall health, measured by performance status. Common treatments include surgery, chemotherapy, and radiotherapy. NSCLC is sometimes treated with surgery, whereas SCLC usually responds better to chemotherapy and radiotherapy.^[13] Overall, 16.8% of people in the United States diagnosed with lung cancer survive five years after the diagnosis,^[14] while outcomes on average are worse in the developing world. Worldwide, lung cancer is the most common cause of cancer-related death in men and women, and was responsible for 1.56 million deaths annually, as of 2012.^[14] Signs and symptoms which may suggest lung cancer include; Respiratory symptoms: coughing, coughing up blood, wheezing, or shortness of breath, Systemic symptoms: weight loss, weakness, fever, or clubbing of the fingernails and Symptoms due to the cancer mass pressing on adjacent structures: chest pain, bone pain, superior vena cava obstruction, or difficulty swallowing If the cancer grows in the airways, it may obstruct airflow, causing breathing difficulties. The obstruction can lead to accumulation of secretions behind the blockage, and predispose to pneumonia.^[15] Depending on the type of tumor, paraneoplastic phenomena—symptoms not

50 due to the local presence of cancer—may initially attract attention to the disease.^[16] In lung cancer, these
 51 phenomena may include hypercalcemia, syndrome of inappropriate antidiuretic hormone (SIADH,
 52 abnormally concentrated urine and diluted blood), ectopic ACTH production, or Lambert–Eaton myasthenic
 53 syndrome (muscle weakness due to autoantibodies). Tumors in the top of the lung, known as Pancoast
 54 tumors, may invade the local part of the sympathetic nervous system, leading to Horner's syndrome
 55 (drooping of the eyelid and a small pupil on that side), as well as damage to the brachial plexus.^[17] Many of
 56 the symptoms of lung cancer (poor appetite, weight loss, fever, fatigue) are not specific.^[18] In many people,
 57 the cancer has already spread beyond the original site by the time they have symptoms and seek medical
 58 attention.^[19] Symptoms that suggest the presence of metastatic disease include weight loss, bone pain and
 59 neurological symptoms (headaches, fainting, convulsions, or limb weakness).^[20] Common sites of spread
 60 include the brain, bone, adrenal glands, opposite lung, liver, pericardium, and kidneys.^[20] About 10% of
 61 people with lung cancer do not have symptoms at diagnosis; these cancers are incidentally found on routine
 62 chest radiography.^[21-22] Therefore in this paper, we intend to;

- 63 i. Investigate the number of patients and deaths affected with lung cancer
- 64 ii. Test the effect of sex on lung cancer incidence
- 65 iii. Test the effect of environment and educational level on lung cancer incidence
- 66 iv. Examine the trend in lung cancer.
- 67 v. Measure the relative risk associated with lung cancer.

69 2. Methodology

70 To achieve the set objectives, data pertaining the subject matter was obtained from the records unit of five
 71 different hospitals.

73 2.1 Chi-Square Test for Independence

74 This test was applied to investigate the agreement between the observed and expected frequencies;

$$X^2 = \sum_{i=1}^r \sum_{j=1}^c \frac{(o_{ij} - e_{ij})^2}{e_{ij}}$$

75 And to test the hypothesis of independence

76 H_0 : The Classification is independent

77 H_1 : The Classification is dependent

80 2.2 Regression Model

81 Here we shall make use of the estimated model given by;

$$\hat{y} = a + bx$$

82 To determine the relationship between the number of lung cancer patients and their death cases where,

$$\hat{b} = \frac{(n \sum xy - \sum x \sum y)}{n \sum x^2 - (\sum x)^2}$$

$$\hat{a} = \bar{y} - \hat{b}\bar{x}$$

90 2.3 Correlation Coefficient 'R' and Coefficient Of Determination 'R²'

$$\hat{b} = \frac{(n \sum xy - \sum x \sum y)}{(n \sum x^2 - (\sum x)^2)(n \sum y^2 - (\sum y)^2)}$$

$$R^2 = \frac{SS_Y - SS_E}{SS_{YY}} = 1 - \frac{SS_E}{SS_{YY}} \quad \text{for } 0 < R^2 < 1$$

93
94 2.4 Odds Ratio

95
96 We employed this ratio to measure the risk of experiencing the outcome under study when the antecedent
97 factor is present.

98 Table 1: Odd Ratio

	B	\bar{B}	Total
A	P_{11}	P_{12}	$P_{1.}$
\bar{A}	P_{21}	P_{22}	$P_{2.}$
Total	$P_{.1}$	$P_{.2}$	$P_{..}$

100
101 Therefore,

$$O_A = \frac{P_{11}}{P_{12}}$$

$$O_{\bar{A}} = \frac{P_{21}}{P_{22}}$$

$$O = \frac{O_A}{O_{\bar{A}}}$$

$$S.E(O) = \frac{O}{(n)^{1/2}} = \left(\frac{1}{P_{11}} + \frac{1}{P_{12}} + \frac{1}{P_{21}} + \frac{1}{P_{22}} \right)^{1/2}$$

102
103 Thus, the estimated odds ratio is;

$$RR = \frac{P(B/A)}{P(\bar{B}/A)}$$

104
105 **3. Data Analysis and Result**

106
107 3.1 Chi-Square Test for Independence of Sex on Lung Cancer Cases.

108
109 Table 2: Data Showing Age and Sex on Lung Cancer

Age	Sex		Total
	Male	Female	
< 50	26	5	31
≥ 50	22	8	30
Total	48	13	61

111
112 H_0 : Lung Cancer cases are independent on Sex
113 H_1 : Lung Cancer cases are dependent on Sex

114
115 Table 3: Age * Sex Cross tabulation

			Sex		Total
			Male	Female	
Age < 50	Count	26	5	31	
	Expected Count	24.4	6.6	31.0	

116

≥ 50	Count	22	8	30
	Expected Count	23.6	6.4	30.0
Total	Count	48	13	61
	Expected Count	48.0	13.0	61.0

117
118
119

Table 4: Chi-Square Test

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.010 ^a	1	.315	.363	.245
Continuity Correction ^b	.479	1	.489		
Likelihood Ratio	1.016	1	.313		
Fisher's Exact Test					
Linear-by-Linear Association	.993	1	.319		
N of Valid Cases ^b	61				

120
121
122
123
124
125
126
127

From Table 4, we see that " $\chi_{cal}^2 = 1.010$ " this χ_{cal}^2 value is less than the " $\chi_{0.05,1}^2 = 3.841$ " thus, we do not reject the null hypothesis and therefore conclude that lung cancer cases are independent on Gender.

3.2 Chi-Square Test for Independence Of Environment on Lung Cancer Cases.

Table 5: Data Showing Age and environment on lung Cancer

Age	Environment		Total
	Urban	Rural	
< 50	22	9	31
≥ 50	17	13	30
Total	39	22	61

128
129
130
131
132
133
134
135
136
137
138

H_0 : Lung Cancer cases are independent on Environmental factor

H_1 : Lung Cancer cases are dependent on Environmental factor

Table 6: Age * Environment Cross tabulation

Age	< 50	Count	Environment		Total
			Urban	Rural	
		Count	22	9	31
		Expected Count	19.8	11.2	31.0

≥ 50	Count	17	13	30
	Expected Count	19.2	10.8	30.0
Total	Count	39	22	61
	Expected Count	39.0	22.0	61.0

139
140
141

Table 7: Chi-Square Test

	Value	Df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.352 ^a	1	.245		
Continuity Correction ^b	.803	1	.370		
Likelihood Ratio	1.358	1	.244		
Fisher's Exact Test				.293	.185
Linear-by-Linear Association	1.330	1	.249		
N of Valid Cases ^b	61				

142

143 From Table 7, we see that " $\chi_{cal}^2 = 1.352$ ", this χ_{cal}^2 value is less than the " $\chi_{0.05,1}^2 = 3.841$ " thus, we do not
144 reject the null hypothesis and therefore conclude that lung cancer cases are independent on environmental
145 factor.

146

147

148 3.3 Chi-Square Test for Independence of Educational Level on Lung Cancer Cases.

149

Table 8: Data Showing Age and Educational Level on Lung Cancer

150

151

Age	Educational Level			Total
	Tertiary	Secondary	Primary	
< 50	12	13	6	31
≥ 50	5	12	13	30
Total	17	25	19	61

152

153 H_0 : Lung Cancer cases are independent on Educational Level

154 H_1 : Lung Cancer cases are dependent on Educational Level

155

156

157

Table 9: Age * Educational Level Cross tabulation

		Educational Level			Total
		Tertiary	Secondary	Primary	
Age < 50	Count	12	13	6	31
	Expected Count	8.6	12.7	9.7	31.0
> 50	Count	5	12	13	30

	Expected Count	8.4	12.3	9.3	30.0
Total	Count	17	25	19	61
	Expected Count	17.0	25.0	19.0	61.0

Table 10: Chi-Square Test

	Value	Df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.486 ^a	2	.064
Likelihood Ratio	5.634	2	.060
Linear-by-Linear Association	5.392	1	.020
N of Valid Cases	61		

From Table 10, we see that " $\chi_{cal}^2 = 5.486$ ", this χ_{cal}^2 value is less than " $\chi_{0.05,2}^2 = 5.991$ " thus, we do not reject the null hypothesis and therefore conclude that lung cancer cases are independent on educational level.

3.4 Regression Analysis on the Total Number of Lung Cancer Cases and Death from Such Cases

Table 11: Model Summary

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.783 ^a	.614	.607	.968

Table 12: Coefficients

Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.
		B	Std. Error	Beta		
1	(Constant)	.805	.372		.230	.042
	Lung Cancer Cases	.362	.037	.783	9.679	.000

Table 11 clearly shows a strong linear relationship exists between Lung Cancer and death from such disease, implying that increase in the number of lung cancer cases has very high positive effect on the occurrence of death ($r = 0.783$). Also, 61.4% of the variation in death occurrence is explained by lung cancer cases while 38.6% of the variation is due to other factors other than lung cancer. Table 12 shows that a unit increase in lung cancer cases results in an increase in the number of death occurrence ($b = 0.362$), implying that there is a direct relationship between the number of lung cancer cases and the number of death occurrence from the disease.

3.5 Calculation of Odds Ratio for Lung Cancer Cases.

184
185

Table 13: Age * State of Patient

Age		State of Patients		Total
		Death	Alive	
A	< 50	9	22	31
\bar{A}	≥ 50	14	16	30
Total		23	38	61

186
187
188
189
190
191
192
193
194
195
196

Table 14: Proportions; Age * State of Patient

Age		State of Patients		Total
		Death	Alive	
A	< 50	0.15	0.36	0.51
\bar{A}	≥ 50	0.23	0.26	0.49
Total		0.38	0.62	1

$$P(B/A) = 0.71$$

$$P(\bar{B}/A) = 0.55$$

$$O_A = 0.42$$

$$O_{\bar{A}} = 0.88$$

$$O = 0.47$$

$$RR = \frac{P(B/A)}{P(\bar{B}/A)} = 0.62$$

197
198
199
200
201
202
203
204

From the equations above, O_A is 5/12 implying that 5 out of every 12 lung cancer patients aged less than 50 years is expected to die. Similarly, $O_{\bar{A}}$ is 23/26 implying that 23 out of every 26 lung cancer patient aged more than 50 years is expected to die. Equation 5 revealed an odds ratio of 0.41 indicating that the odds of lung cancer patient aged less than 50 years dying is 51% lesser than those aged 50 years and above. Relative Risk of lung cancer patient dying is " $^{31}/_{50} \approx 0.62$ " times higher for patients aged 50 years and above when compared with those aged below 50 years of age.

4. Conclusion and Recommendation

205
206
207
208
209
210
211
212
213
214
215

Based on the findings so far, we hereby conclude that the prevalence of lung cancer is independent on sex, environment and educational level, this therefore implies that it depends on other factors not considered in the study, this may include; tobacco smoking, genetic factors and exposure to random gas, asbestos or other forms of air pollution. Also, lung cancer claims more life in Older patients (age ≥ 50 yrs) than in younger patients (age < 50 yrs). Therefore, the government should try as much as possible to eliminate tobacco smoking and the smoking of cessation. Policy interventions decreasing passive smoking in public areas such as restaurants and workplaces should be put in place. Also, the government to adhere to the World Health Organizations instructions to institute a total ban on tobacco advertising to prevent young people from taking up smoking.

216
217
218
219
220

References

221
222
223

- 224 1. Lung Carcinoma: Tumors of the Lungs. Merck Manual Professional Edition, Online edition.
225 Retrieved 15 August 2007. [http://www.merckmanuals.com/professional/pulmonary-](http://www.merckmanuals.com/professional/pulmonary-disorders/tumors-of-the-lungs/lung-carcinoma)
226 [disorders/tumors-of-the-lungs/lung-carcinoma](http://www.merckmanuals.com/professional/pulmonary-disorders/tumors-of-the-lungs/lung-carcinoma)
227
- 228 2. Falk, S; Williams, C (2010). Chapter 1. Lung Cancer—the facts (3rd ed.). Oxford University Press.
229 pp. 3–4. ISBN 978-0-19-956933-5. https://wikivisually.com/wiki/Lung_cancer
230
- 231 3. Horn, L; Lovly, CM; Johnson, DH (2015). Chapter 107: Neoplasms of the lung. In Kasper, DL;
232 Hauser, SL; Jameson, JL; Fauci, AS; Longo, DL; Loscalzo, J. Harrison's Principles of Internal
233 Medicine (19th ed.). McGraw-Hill. ISBN 978-0-07-180216-1.
234 [https://archive.org/stream/DennisKasperAnthonyFauciStephenHauserDanLongoJ.JamesonJosephLoscalzoHarrisonsPri/Dennis%20Kasper,%20Anthony%20Fauci,%20Stephen%20Hauser,%20Dan%20Longo,%20J.%20Jameson,%20Joseph%20Loscalzo-Harrison's%20Principles%20of%20Internal%20Medicine.%202%20vols.-McGraw-Hill%20\(2015\)_djvu.txt](https://archive.org/stream/DennisKasperAnthonyFauciStephenHauserDanLongoJ.JamesonJosephLoscalzoHarrisonsPri/Dennis%20Kasper,%20Anthony%20Fauci,%20Stephen%20Hauser,%20Dan%20Longo,%20J.%20Jameson,%20Joseph%20Loscalzo-Harrison's%20Principles%20of%20Internal%20Medicine.%202%20vols.-McGraw-Hill%20(2015)_djvu.txt)
235
236
237
238
239
- 240 4. Thun MJ, Hannan LM, Adams-Campbell LL, et al. (September 2008). Lung cancer occurrence in
241 never-smokers: an analysis of 13 cohorts and 22 cancer registry studies. PLoS Medicine 5 (9): e185.
242 doi:10.1371/journal.pmed.0050185. PMC 2531137. PMID 18788891.
243 <http://journals.plos.org/plosmedicine/article?id=10.1371/journal.pmed.0050185>
244
- 245 5. Obubu, M. Nwokolo P.C (2016). Prevalence of Breast Cancer in Delta State, Nigeria. World Journal
246 of Probability and Statistics. Vol. 2, No. 2, Pp 1-9
247
- 248 6. Osuji G.A., Obubu, M., Obiora-Ilouno H.O (2016). An investigation on the causes of Low birth
249 weight in Delta State, Nigeria. European Journal of Statistics and Probability. Vol. 4, No 1, pp. 1-6
250
- 251 7. Osuji G.A., Obubu, M., Obiora-Ilouno H.O (2016). Uterine Fibroid on Women's Fertility and
252 Pregnancy Outcome in Delta State, Nigeria. Journal of Natural Sciences Research, Vol. 6, No 2, pp.
253 27-33.
254
- 255 8. Alberg AJ, Samet JM (2010). "Chapter 46". Murray & Nadel's Textbook of Respiratory Medicine
256 (5th ed.). Saunders Elsevier. ISBN 978-1-4160-4710-0. [https://www.elsevier.com/books/murray-](https://www.elsevier.com/books/murray-and-nadels-textbook-of-respiratory-medicine/mason/978-1-4557-0873-4)
257 [and-nadels-textbook-of-respiratory-medicine/mason/978-1-4557-0873-4](https://www.elsevier.com/books/murray-and-nadels-textbook-of-respiratory-medicine/mason/978-1-4557-0873-4)
258
- 259 9. O'Reilly, KM; Mclaughlin AM; Beckett WS; Sime PJ (March 2007). Asbestos-related lung disease.
260 American Family Physician 75 (5): 683–688. PMID 17375514
261 <https://www.ncbi.nlm.nih.gov/pubmed/17375514>
262
- 263 10. Carmona, RH (27 June 2006). The Health Consequences of Involuntary Exposure to Tobacco
264 Smoke: A Report of the Surgeon General. U.S. Department of Health and Human Services.
265 Secondhand smoke exposure causes disease and premature death in children and adults who do not
266 smoke. Retrieved 2014-06-16. [https://www.healthypeople.gov/2020/tools-resources/evidence-based-](https://www.healthypeople.gov/2020/tools-resources/evidence-based-resource/the-health-consequences-of-involuntary-exposure-to)
267 [resource/the-health-consequences-of-involuntary-exposure-to](https://www.healthypeople.gov/2020/tools-resources/evidence-based-resource/the-health-consequences-of-involuntary-exposure-to)
268
- 269 11. Tobacco Smoke and Involuntary Smoking (PDF). IARC Monographs on the Evaluation of
270 Carcinogenic Risks to Humans (WHO International Agency for Research on Cancer) 83. 2004.
271 There is sufficient evidence that involuntary smoking (exposure to secondhand or 'environmental'
272 tobacco smoke) causes lung cancer in humans. ... Involuntary smoking (exposure to secondhand or
273 'environmental' tobacco smoke) is carcinogenic to humans (Group 1).
274
- 275 12. Lu C, Onn A, Vaporciyan AA, et al. (2010). 78: Cancer of the Lung. Holland-Frei Cancer Medicine
276 (8th ed.). People's Medical Publishing House. ISBN 978-1-60795-014-1.
277

- 278 13. Chapman, S; Robinson G; Stradling J; West S (2009). Chapter 31. Oxford Handbook of Respiratory
279 Medicine (2nd ed.). Oxford University Press. ISBN 978-0-19-954516-2.
280
- 281 14. Surveillance, Epidemiology and End Results Program. National Cancer Institute. Retrieved 15 July
282 2014.
283
- 284 15. World Cancer Report 2014. World Health Organization. 2014. pp. Chapter 1.1. ISBN 9283204298.
285
- 286 16. Honnorat, J; Antoine JC (May 2007). Paraneoplastic neurological syndromes. Orphanet Journal of
287 Rare Diseases (BioMed Central) **2** (1): 22. doi:10.1186/1750-1172-2-22. PMC 1868710.
288 PMID 17480225. [http://www.cancereffects.com/Lung-Cancer-\(Cancer-of-Lungs\)-Symptoms,-
289 Screening,-Rates,-Treatment.html](http://www.cancereffects.com/Lung-Cancer-(Cancer-of-Lungs)-Symptoms,-Screening,-Rates,-Treatment.html)
290
- 291 17. Greene, Frederick L. (2002). AJCC cancer staging manual. Berlin: Springer-Verlag. ISBN 0-387-
292 95271-3. <http://www.springer.com/gp/book/9780387884424>
293
- 294 18. Collins, LG; Haines C; Perkel R; Enck RE (January 2007). Lung cancer: diagnosis and management.
295 American Family Physician (American Academy of Family Physicians) **75** (1): 56–63.
296 PMID 17225705. <https://www.ncbi.nlm.nih.gov/pubmed/17225705>
297
- 298 19. Osuji G.A., Obubu, M., Obiora-Ilouno H.O., Okoro, C.N (2015). Post-Partum Hemorrhage in Delta
299 State, Nigeria; A Logistic Approach. International Journal of Sciences: Basic and Applied Research
300 (IJSBAR). Vol. 24, No 6, pp. 45-53
301
- 302 20. Osuji G.A., Obubu, M., Obiora-Ilouno H.O., Nwosu, D.F (2015). Perinatal Mortality and Associated
303 Obstetric Risk Factors in Urban Delta State, Nigeria; Rural-Urban Differences. International Journal
304 of Mathematics and Statistics Studies Vol. 3, No. 5, PP 32-46
305
- 306 21. Obubu, M., Okoye Valentine, Omoruyi Frederick, Ngonadi Lilian Oluebube (2017). Infant
307 Mortality; a continuing social problem in Northern Nigeria: Cox Regression Approach. American
308 Journal of Innovative Research and Applied Sciences.2017; 5(5):1-5
309
- 310 22. Maxwell O, Friday AI, Chukwudike NC, et al. A theoretical analysis of the odd generalized
311 exponentiated inverse Lomax distribution. Biom Biostat Int J. 2019; 8(1): 17-22. DOI:
312 10.15406/bbij.2019.08.00264.