

New Documentary of Acid-Stable Glutaminase Production by an Efficient Acidophilic *Aspergillus niger* CPGM 1439

ABSTRACT

Samples for isolation were taken from natural spoiled salted foods. All acidophilic isolates were subjected under comparison for finding the most potent glutaminase producing one. Optimization of the produced enzyme was done in the light of seven environmental and five nutritional factors including incubation time, temperature, pH, buffer system, static and different shaking speeds conditions, dark and light conditions, different inoculum size, additional carbon sources, additional nitrogen sources, heavy metals, vitamins, amino acids. Five fungal isolates have been selected and were tested for their ability to grow on L-glutamine containing medium at pH 3 and 2. Only 5 fungal isolates (F1.a, F2.a, F3.a, F4.a and F5.a) were able to grow efficiently on pH3, two of them were able to grow also on pH2. The activity of enzyme was measured extracellularly and intracellularly for each isolate and were 20.34, (2.68), 17.8, (0.423), 5.77, (5.77), 10.27, (1.39), 2.28 and (0.071) U/ml respectively. The highest enzyme activity producing isolate (F1.a) was selected and identified genetically. It was found to be closely related to *Aspergillus niger* strain CPGM 1439 with ~~similarity~~ 93% homology and accession number DQ196192.1. The yield was quite high after optimization reached 49.59 U/ml extracellularly which is essential advantage for such production method. Glutaminase enzyme was produced efficiently by the finally selected *Aspergillus niger* isolate and optimization lead to higher enzyme activity which was measured by U/ml in each step. The maximum yield was obtained at 35°C, using KCl- HCl buffer at pH1.8, 100 ppm Mn⁺², addition of iso- leucine 3.59% for 7 days incubation in dark condition and shaking speed 100 rpm.

Keywords: glutaminase, acidstable, enzyme activity, optimization, extracellular and intracellular.

1. INTRODUCTION

L- glutaminase (L-glutamine amidohydrolases E.C. 3.5.1.2) catalyses the hydrolysis of L-glutamine to glutamic acid and ammonia (Kumar *et al.*, 2012). This enzyme ~~is very economically important enzyme has wide range of economic importance~~ as it can be used in food industry as tool for hydrolysis of protein because of its action in conversion of L- glutamine to ammonia and glutamic acid which gave the special flavor to food products (Saptarshi and Lele, 2011) also, it is nowadays the core of many researches to be used as therapeutic agent for cancer by searching for ~~remarkable-potential and selective~~ inhibitors for this enzyme as it regard central enzyme for many conversions occur in the cell which lead to increase proliferation of cancer cells (Lukey *et al.*, 2013). Moreover, L-glutaminase is applied in the therapy of human immunodeficiency virus (HIV) (Roberts *et al.*, 1972) and could be used as efficient antiretroviral agent (Roberts and McGregor, 1991); L-glutaminase biosensors for monitoring L-glutamine level (Sabu *et al.*, 2000) and in food industries imparting the flavor and aroma to the foods (Loeliger, 2000; Durá *et al.*, 2004).

There is great trend to produce L-glutaminase enzyme with unique stability properties to withstanding harsh industrial conditions ~~so, it may by persist in industrial process~~ and ~~keep its- retain its~~ activity under such ~~tough harsh industrial processes/conditions~~. L- glutaminase can be found in different types of organisms (Binod *et al.*, 2017) including bacteria (Prakash *et al.*, 2009), fungi (Keerthi *et al.*, 1999), yeast

: [1P]Comment

Formatted: Font: Not Bold

(Fukushima and Motai, 1990), Actinomycetes (Suresh *et al.*, 2013) and can also be extracted from plants (Bülbül and Karakuş, 2013).

It was reported that, there are many factors that can affect the production of glutaminase enzyme from any microorganism which differ according to the nature of the producer microorganism as pH, temperature, incubation time, additional substances or vitamins of metals of natural supplements can greatly cause great influence on the enzyme yield, optimization of production is very important shotgun point (Durai *et al.*, 2014).

The aim of this study was to produce L- glutamines biologically with high yield and activity and tolerance or stable to high acidic medium ~~has the property of being still active under high acidic conditions.~~

2. MATERIAL AND METHODS

2.1. Isolation

Isolation was carried out on this medium: The medium was composed of (g/l) NaCl 10, MgSO₄.7H₂O 10, K₂HPO₄ 10, Glutamine 10 and Agar 20. The pH of the medium was adjusted at 2 and 3. Incubation was for five days at 30 °C, the isolates which were able to be subcultured with high growth were selected and purified.

2.2. Separation of the extracellular and intracellular crude enzyme

The extracellular crude enzyme was separated by filtration and the mycelium was washed with distilled H₂O. The intracellular contents of the cells were obtained by ultrasonic disruption of cells by ultrasonic processor (Cole Parmer Ultrasonic Homogenizer CPX 400) over three 15 s periods, and with an interval of 45 s between periods. The sonicated samples were centrifuged at 15,000 rpm for 30 min at 4°C to remove cell-debris using Sorvall cooling centrifuge. The supernatants were then used for assay of enzyme (Kalimuthu *et al.*, 2008).

2.3. Quantitative assay of glutaminase enzyme

The assay was done according to Imada *et al.* (1973). First the reaction mixture was prepared by adding 0.5 ml of the sample, 0.5 ml of 0.04M L-glutamine solution, 0.5 ml of distilled water and 0.5 ml of phosphate buffer 0.1M, pH 8.0. The mixture was incubated at 37°C for 15 min then, 0.5 ml of 1.5 M Trichloroacetic acid was added for stopping the reaction. After that, 0.1 ml of the mixture was added to 3.7 ml of distilled water and finally 0.2 ml of Nessler's reagent were added to complete the volume to 4 ml. the same steps were done for controls. The absorbance was measured at 450nm using a UV -Visible spectrophotometer. The absorbance reading was indication for ammonia liberated from the action of the enzyme so; standard curve using ammonium chloride was made using serial dilutions of ammonium chloride covering nearly all the range which had been obtained. The enzyme yield was expressed as Unit/ml (U/ml).

It's known clearly that, one international unit of L-glutaminase was defined as the amount of enzyme that liberates one micromole of ammonia under optimum conditions.

3. RESULTS AND DISCUSSION

3.1. Isolation of acidophilic glutaminase ~~producers~~ producing microorganisms

In the present investigation the starting basic point is searching for new sources for isolation of L-glutaminase producers from acidic environment to obtain novel isolates with special characters. In this respect 5 fungi were isolated at pH 3 (F1.a, F2.a, F3.a, F4.a and F5.a) from pickled lemon, pickels, lemon and salted cheese and only two isolates were able to grow at pH 2 (F1.a and F2.a). The five fungal isolates were assayed extracellularly and intracellularly for glutaminase production as recorded in table (1), Isolates F1.a followed by F2.a were the most producers extracellularly and recorded 20.34 and 17.80 U/ml, respectively. In addition, the intracellular enzymes produced by the five fungal isolates have moderate productivity compared with the extracellular glutaminase. Extracellular glutaminases are more advantageous over intracellular one as they could be produced economically industrially in the nutrient medium and purified easily; this was in full agreement with the results of **Renu and Chandrasekaran (1992)**.

Table 1. Quantitative assay of glutaminase for selected acidophilic isolates

| Isolate code | Production type | Released ammonia (M) | Enzyme Productivity (U/ml) |
|--------------|-----------------|----------------------|----------------------------|
| F1.a | Extracellular | 0.02034 | 20.34 |
| | Intracellular | 0.00268 | 2.68 |
| F2.a | Extracellular | 0.0178 | 17.80 |
| | Intracellular | 0.000423 | 0.423 |
| F3.A | Extracellular | 0.00577 | 5.77 |
| | Intracellular | 0.00577 | 5.77 |
| F4.A | Extracellular | 0.01027 | 10.27 |
| | Intracellular | 0.00139 | 1.39 |
| F5.A | Extracellular | 0.00228 | 2.28 |
| | Intracellular | 0.0000706 | 0.071 |

A variety of microorganisms, including bacteria, yeast, and filamentous fungi, from both soil and marine habitat have been reported to produce L-glutaminase. In this aspect, many researches have been reported isolation of glutaminase producers from soils as **El-Sayed (2009)** who isolated 20 fungi and tried to produce glutaminase. Also, **Sinha and Nigam (2016)** isolated a soil bacterium NRBT2228 on Zobell's agar medium which was used for screening and maintenance of the bacterial culture. In addition, several studies have been started using salt lakes or water in general as the source of the isolation as reported by **Durai et al. (2014)** who obtained his isolate from marine sediment of coastal environment growing on mineral salt glutamine medium as the medium for production (**Suresh, 1996**) while culture maintenance was on Bennet' agar medium (**Sabu, 1999**).

3.2. Optimization of glutaminase productivity for selected acidophilic isolate (F1.a)

seven environmental and five nutritional factors were carried out including incubation time, temperature, pH, buffer system, static and different shaking speeds conditions, dark and light conditions, different inoculum size, additional carbon sources, additional nitrogen sources, heavy metals, vitamins, amino acids.

Concerning the incubation time for the glutaminase production by isolate F1.a, the best production time was from 6 - 8 days with the best at 7th day, the reason for decreasing in the enzyme yield after 8th day might be attributed to the inactivation of glutaminase by the protease secreted by the fungus as reported by (Zhu *et al.*, 1994). While the optimum incubation temperature was 35°C and the best inoculum size used was 2.48×10^6 spore/ml and any further increase above this level resulted in a marginal decline in the enzyme production. These results were in full agreement with that of Hamed and Al-wasify (2016) who reported that the maximum yield of glutaminase by *Fusarium oxysporum* was achieved at 35°C, and after 7 days of incubation using inoculum of 100 µl / 30 ml medium. Also, El-Sayed (2009), reported the best production was at 30°C and 7 days of incubation and inoculum size was 3ml (ml/wt) for L-glutaminase produced by *Trichoderma koningii*.

In the present investigation, the effect of light and dark as well as speed of agitation on enzyme production were also studied. Dark was preferred by the selected fungal isolate and 100 rpm was also result in activation of the productivity of L-glutaminase (No literatures were available concerned with this this point of research with glutaminase production).




Fig.1. Effect of different incubation time, temperature, inoculum size, dark and light and agitation speed on glutaminase productivity by isolate (F1.a)

This is the first documentary to produce L-glutaminase by acidophilic microorganisms. Isolate F1a. can live and produce L-glutaminase at different acidic pH's ranged from 1.5 up to 5 by adjusting the pH using 1N HCL but it scored maximum productivity at 1.5 followed by 2.0 with 28.55 and 26.20 U/ml of enzyme productivity, respectively. Probably the isolate could be assigned as acidophilic with reference to enzyme production. By using different acidic buffer systems, Hydrochloric Acid-Potassium Chloride buffer was best buffer system used (25.93 U/ml) but it still less than that adjusted by the 1NHCl. In view of the findings of other investigators, few reviews were found concerned with the production of L-glutaminase with microorganisms at lower pH , only pH range from 4-6 was used in this respect as showed by Krishnan *et al.* (2011), pH 4.1 Shirazian *et al.* (2016), pH 5.0 by *Aspergillus oryzae* S 2; Ahmed *et al.* (2016), pH 4 by *Aspergillus* sp. ALAA-2000. The present isolate has the ability to produce L-glutaminase at pH 1.5, so the produced L-glutaminase must be acido-stable and can stand the acidity of the HCl in the stomach if used in food industry.

Table 2. Effect of different pH values and different buffer systems on glutaminase productivity for isolate F1a.

(a) different pH vales **(b) different buffer systems**

| pH | Final pH | Enzyme Productivity (U/ml) |
|-----|----------|----------------------------|
| 1 | 1.7 | 0.749 |
| 1.5 | 5.3 | 28.55 |
| 2 | 6.2 | 26.2 |
| 2.5 | 6.3 | 19.37 |
| 3 | 6.2 | 17.45 |
| 3.5 | 6.5 | 15.97 |
| 4 | 6.4 | 15.5 |
| 4.5 | 6.4 | 14.61 |
| 5 | 6.9 | 15.07 |

| Buffer system | pH | Enzyme Productivity (U/ml) |
|----------------|-----|----------------------------|
| HCl-KCl | 1 | 1.61 |
| | 1.2 | 1.85 |
| | 1.4 | 7.86 |
| | 1.6 | 19.37 |
| | 1.8 | 25.93 |
| | 2 | 13.04 |
| Phthalate-NaOH | 2.2 | 8.59 |
| | 2.4 | 13.49 |
| Glycine- HCl | 2.2 | 10.35 |
| | 2.4 | 7.69 |
| | | 9.24 |

Concerning the effect of different medium salts concentrations such as K_2HPO_4 , NaCl, $MgSO_4 \cdot 7H_2O$ on L-glutaminase productivity by the acidophilic isolate isolate F1a., a concentration of 1% of all the tested salts from the all concentrations used was the best it gave a productivity of 28.56 U/ml for all respectively as shown in Table 3. Incase of NaCl the isolate showed considerable level of L-glutaminase syntheses even in the low concentration of NaCl, and also at higher concentrations, up to 3 % (w/v), however, with further increase in sodium chloride concentration enzyme production decreased.

In case of medium composition, maximum enzyme production was achieved with addition of 2% glutamine and 2% soybean meal (Kiruthika and Saraswathy, 2013). Another study has shown enzyme activity 4.231U/ml using *Bacillus* for production of glutaminase enzyme on Zobell's agar media (ZM) and in the presence of 1g/L of L-glutamine (Sinha and Nigam, 2016). Also, maximum yield of enzyme recorded in medium supplemented with 1 % sucrose as carbon source and 0.025% glutamine concentration (Hamed and Al-wasify, 2016). Also minimal glutamine agar medium was used as the basic medium for production of enzyme (Sridevi *et al.*, 2016).

Table 3. Effect of different concentrations of K_2HPO_4 , NaCl and $MgSO_4 \cdot 7H_2O$ on L-glutaminase productivity by the most potent fungal isolate

| K_2HPO_4 (%) | Enzyme Productivity (U/ml) | NaCl (%) | Enzyme Productivity (U/ml) | $MgSO_4 \cdot 7H_2O$ (%) | Enzyme Productivity (U/ml) |
|-------------------|----------------------------------|-------------|----------------------------------|-----------------------------|----------------------------------|
| 0.5 | 13.59 | 0.5 | 1.25 | 0.5 | 19.31 |
| 1 | 28.56 | 1 | 28.56 | 1 | 28.56 |
| 2 | 22.72 | 2 | 20.86 | 2 | 23.69 |
| 3 | 23.19 | 3 | 22.77 | 3 | 0.52 |

In case of studying the effect of different glutamine concentration as a substrate for the production of L-glutaminase by the most potent acidophilic fungal isolate F1a., a concentration of 2% was exhibited the highest productivity, this result was in complete accordance with the results of **Kumar and Chandrasekaran (2003); El-Sayed (2009); Kiruthika and Nachimuth (2013)** who proved that a 2% of glutamine was enough to produce the highest L-glutaminase by *Pseudomonas* sp. BTMS-51; *Trichoderma koningii* and *Vibrio azureus* JK-79, respectively. On the other hand, **Sabu et al. (2000) and Iyer and Singhal (2010)** reported that only 0.25 and 0.5 % glutamine was enough in the production of L-glutaminase, respectively. While, **Wakayama et al. (2005)** found that 1% glutamine exhibited the highest L-glutaminase activity by *Stenotrophomonas maltophilia* NYW-81. Moreover, a concentration of 3% glutamine proved to produce the highest L-glutaminase activity by as reported by **Revanth and Raju (2013)** and maltose as reported by **Kiruthika and Nachimuth (2013)**.

Table 4. Effect of different glutamine concentrations on enzyme productivity for acidophilic isolate (F1.a)

| Glutamine (%) | Enzyme activity (U/ml) |
|---------------|------------------------|
| 0.25 | 7.44 |
| 0.5 | 15.12 |
| 1 | 25.25 |
| 2 | 39.25 |
| 3 | 28.95 |

In the present investigation, incorporation of additional carbon sources doesn't enhance any increase in the L-glutaminase productivity compared with control group, but on the other hand inhibit the productivity. In view of the findings of the other workers, many sources utilized found to be glutaminase inhibitors like fructose, sucrose (**Desai et al., 2016**) lactose and starch were found to have inhibitory effect on glutaminase (**Abdallah et al., 2012**) while enhancing has been also approved as on addition of glucose (**Sato et al., 1999; Sabu et al., 2000; Kumar and Chandrasekaran, 2003; El-Sayed, 2009; Iyer and Singhal, 2009; Krishnakumar et al., 2011**), sucrose (**Iyer and Singhal, 2010; Revanth and Raju, 2013**), maltose (**Pallem et al., 2010**), sorbitol (**Wakayama et al., 2005**), galactose (**Kumar et al., 2013**).

In the present study, all the nitrogen sources organic and inorganic tested along with L-glutamine resulted in a decrease in L-glutaminase yield compared to control. The results indicated very clearly that addition of additional nitrogen sources in did not enhance enzyme production, and contradictory these additional nitrogen sources inhibited L-glutaminase synthesis by the acidophilic fungus which was in compatibility with **Desai et al.(2016)** urea and peptone were recorded as inhibitors for glutaminase. Testing adding different nitrogen sources were concerning other researchers as **Sood et al. (2002)** who found that, addition of NH_4NO_3 or NH_4Cl enhanced the activity of glutaminase while KNO_3 decreased or has no effect under certain conditions typically with **Aly et al.(2017)** NH_4Cl also was found as glutaminase enhancer(**Rashmi et al., 2012**). Some researchers reported that, on supplementation with additional nitrogen compound the activity increased as yeast extract (**Kumar et al., 2013**), malt extract (**Pallem et al., 2010**) peptone and sodium nitrate (**Abdallah et al., 2012**).

This variability referred to the behavior of the producing microorganism for the present additional carbon or nitrogen source basically the presence of specific enzymes also, it was proved that carbon may

result in increasing N compound uptake and transport changing the behavior of microorganism for the production of glutaminase the core of the research (Fellbaum *et al.*, 2012)




Fig.2. Effect of additional different carbon and nitrogen sources on enzyme productivity for acidophilic isolate (F1.a)

Some enzymes may require the presence of specific metal for their activation or for increasing their activity as the microorganism can use this metal as electron donor or acceptor deriving the energy required for the microorganism thus, six different metal containing compounds were used at three different concentrations 50, 100 and 200 ppm of the metal testing their effect on glutaminase productivity. the acidophilic isolate f1.a showed maximum activity 39.93 U/ml at 100 ppm Mn^{+2} as manganous sulphate comparing to control 39.25 the other tested six metal caused inhibition of the enzyme mainly HgCl_2 and CdSO_4 inversely affected the glutaminase activity with decreasing badly. This may be due to the hazardous effect of Hg and Cd on the isolate growth in the first place and so its ability to survive and producing the enzyme owing to the fact that, not all microorganisms able to synthesizing the enzymes that can help them in detoxifying the hazardous toxic metals (Ehrlich, 1997). Kumar *et al.* (2013) paid out in research that, glutaminase reached maximum activity in the in the presence of MnSO_4 fitted with the results of this research. While in another research the isolate showed maximum glutaminase activity 32.5 U/ml in the presence of ZnCl_2 in concentration 1 w/v

(Nagaraju and Ram, 2018). Also, magnesium was reported as metal activator for the glutaminase by *Aspergillus wentii* MTCC 1901 as magnesium sulphate (Sameera and Raju, 2015) or as magnesium chloride (Ahmed *et al.*, 2016). Many other metals were incorporated and were found as glutaminase inhibitors as Li, Ca, Ba (Ahmed *et al.*, 2016).

Concerning the effect of vitamins on glutaminase, seven vitamins were used with different concentrations (50, 100 and 200 ppm). None of the used vitamins has enhancing impact on glutaminase productivity otherwise all tested vitamins caused inhibition of the enzyme. Vitamins role in microorganisms are cofactor may aiding in metabolism of amino acids or proteins and can increase the activity of the produced enzyme depending on the producing microorganism which interfering in its metabolic cycles (Rosenberg *et al.*, 2017).




Fig. 3. Effect of different (a) Vitamins and (b) heavy metals with different concentrations on glutaminase productivity by acidophilic

In case of supplementation with additional different amino acids together with the glutamine presence, Isoleucine played an important role as inducer for L-glutaminase synthesis for this acidophilic fungus under study followed by Alanine and Aspartic acid with glutaminase activity reached 49.59, 44.01 and 38.15, respectively while, the rest of amino acids were found inhibitory for L-glutaminase productivity by the acidophilic fungus. However, *Ahmed et al. (2016)* reported the production increased in the presence of cysteine. In addition, methionine caused increasing in glutaminase activity (*Sabu, 1999*). L-glutamic acid, l- aspartic and l-asparagine were recorded as glutaminase activity improvers (*Ohshima et al., 1976*) in contrast to *Chitan and Shete. (2012)* asparagine and other tested amino acids (l-tyrosine and l-lysine) were found to be glutaminase inhibitors.

Table 5. Effect of different amino acids on enzyme activity for acidophilic isolate (F1.a)

| Amino acid | Released NH ₄ ⁺ M | Enzyme activity (U/ml) |
|---------------|---|------------------------|
| Alanine | 0.04401 | 44.01 |
| Glycine | 0.03103 | 31.03 |
| Glutamic acid | 0.02948 | 29.48 |
| Asparagine | 0.03815 | 38.15 |
| Aspartic | 0.03492 | 34.92 |
| Iso-Leucine | 0.04959 | 49.59 |
| Cysteine | 0.03607 | 36.07 |
| Valine | 0.03518 | 35.18 |
| Tyrosine | 0.03581 | 35.81 |
| Proline | 0.00171 | 1.71 |

| | | |
|---------|---------|-------|
| Control | 0.03596 | 35.96 |
|---------|---------|-------|

The most potent acidophilic glutaminase producing isolate (F1.a) has been identified genetically based on 18S ribosomal RNA gene partial sequence, internal transcribed spacer 1, 5.8S ribosomal RNA gene, internal transcribed spacer 2 complete sequence and 28S ribosomal RNA gene partial sequence and has shown 93% similarity to *Aspergillus niger* strain CPGM 1439.

Among different sources of L-glutaminase producers, fungal strains possess an elevated edge over others due to their growth requirements and easy processing (Sathish *et al.*, 2008). Reports suggested that these enzymes produced by different microorganisms differed in some physiological, biochemical, catalytic and immunological properties. This led to the continuous screening program for isolation of novel microbial strains that could produce an effective enzyme with few limitations at use.




Fig.4. A dendrogram showing the sequence relationships between *Aspergillus niger* CPGM 1439 and several other strains based on 18S ribosomal RNA gene partial sequence, internal transcribed spacer 1, 5.8S ribosomal RNA gene, internal transcribed spacer 2 complete sequence and 28S ribosomal RNA gene partial sequence

Production and optimization for glutaminase enzyme by fungi was reported for many species as *Trichoderma reesei* (Bülbül and Karakuş, 2013), Genus *Aspergillus* was on the top of producers list as

Aspergillus flavus (Nathiya et al., 2011), *Aspergillus* sp. (Ahmed et al., 2016) *Aspergillus fumigatus* WL002 (Dutta et al., 2015), *Aspergillus wentii* MTCC 1901 (Sameera and Raju, 2015) *A. niger*, *A. fumigatus*, *A. phoenicis* and *Penicillium funiculosum* was also reported as glutaminase producer (Elshafei et al., 2012) .

4. CONCLUSION

Glutaminase enzyme is very important enzyme to be studied extensively and produced due to its applications in medical and industrial aspects. This ~~research~~ research aimed ~~to at~~ at isolating an efficient glutaminase producing microorganism. *Aspergillus niger* was ~~our obtained one isolated, identified~~ and produce glutaminase enzyme with great-high yield ~~reached nearly of about~~ reached nearly of about 50 U/ml after studying many factors that can affect production. Also, it is great advantage that it was produced at very low acidic pH 1.8 which suggests it has high prospect for industrial applications since stability in acidic pH is one of the properties of enzymes required for potential industrial applications. will lead to great chance for more important applications. We ~~It is therefore,~~ recommend that further ~~studying~~ studies carried out for modeling the production and production on large scale fermentors which is expected to will be completed in ~~coming~~ our ongoing studies. Also, it ~~can be useful~~ is suggested to try more natural or wastes as media for production.

REFERENCES

1. Abdallah NA, Amer SK, Habeeb MK. Screening Of L-Glutaminase Produced By Actinomycetes Isolated From Different Soils In Egypt. Int.J.ChemTech Res. 2012;4(4):1451-1460.
2. Ahmed MMA, Taha TM, Abo-Dahab NF, Hassan FSM. Process Optimization of L-Glutaminase Production; a Tumour Inhibitor from Marine Endophytic Isolate *Aspergillus* sp. ALAA-2000. J Microb Biochem Technol. 2016; 8: 382-389.
3. Aly MM, Kadi RH, Aldahlawi AM, Alkhatib MH, Wali AN. Production of the antitumor l-glutaminase enzyme from thermotolerant *Streptomyces* sp. D214 under submerged fermentation conditions. Journal of Experimental Biology and Agricultural Sciences. 2017;5(6):878-885.
4. Binod P, Sindhu R, Madhavan A, Abraham A, Mathew AK, Beevi US, Sukumaran RK, Singh, SP, Pandey A. Recent developments in l-glutaminase production and applications - An overview. Bioresour. Technol. 2017; 245(Pt B):1766-1774.
5. Bülbül D, Karakuş E. Production and optimization of l-glutaminase enzyme from *Hypocrea jecorina* pure culture. Preparative biochemistry and biotechnology J. 2012;43(4):385-397.
6. Chitanand MP, Shete HG. Condition optimization and production of extracellular lglutaminase from *Pseudomonas Fluorescens*. Int J Pharm Bio Sci. 2012;3(3):155 –162.
7. Desai SS, Chopra SJ, Hungung BS. Production, purification and characterization of L-glutaminase from *Streptomyces* sp. isolated from soil. Journal of applied pharmaceutical science. 2016;6(7):100-105.
8. Durá M.A, Flores M, Toldrá F. Effects of curing agents and the stability of a glutaminase from *Debaromyces spp*. Food Chemistry. 2004;86:385–389.

9. Durai S, Selvaraj B, Manikkam R, Ramasamy B. Production and optimization of l-glutaminase from *Vibrio* sp. m9 isolated from mahabalipuram marine sediments. World journal of pharmaceutical research. 2014;3(2), 2117-2126.
10. Dutta S, Ghosh S, Pramanik S. L-asparaginase and L-glutaminase from *Aspergillus fumigatus* WL002: Production and some physicochemical properties. Applied Biochemistry and Microbiology. 2015;51(4):425-431.
11. Ehrlich H. Microbes and metals. Applied Microbiology and Biotechnology. 1997;48(6):687-692.
12. El-Sayed ASA. L- glutaminase production by *trichoderma koningii* under solid state fermentation. Indian J. microbiol. 2009;49:243-250.
13. Elshafei AM, Hassan MM, Ali MA, Mahmoud DA, Elghonemy DH. Screening and optimization of l-asparaginase and l-glutaminase production by some filamentous fungi. Advances in Food Sciences. 2012;34(3): 150-158.
14. Fellbaum CR, Mensah JA, Pfeffer PE, Kiers ET, Bücking H. The role of carbon in fungal nutrient uptake and transport: Implications for resource exchange in the arbuscular mycorrhizal symbiosis. Plant Signaling and Behavior. 2012;7(11): 1509–1512.
15. Fukushima Y, Motai H. Continuous conversion of glutamine to glutamate by immobilized glutaminase producing Yeast. Journal of Fermentation and Bioengineering. 1990;69:189-191.
16. Hamed SR, Alwasify RS. Production and optimization of L-glutaminase from a terrestrial fungal *Fusarium oxysporum*. International journal of pharm tech research. 2016;9(4):233-241.
17. Imada A, Igarasi S, Nakahama K, Isono M. Asparaginase and Glutaminase activities of microorganisms. J. Gen. Microbiol. 1973;76:85-99.
18. Kalimuthu K, Babu SR, Venkataraman D, Bilal M, Gurunathan S.. Biosynthesis of silver nanocrystals by *Bacillus licheniformis*, Colloids and Surfaces B: Biointerfaces. 2008;65, 150-153.
19. Keerthi TR. L- glutaminase production by marine *Pseudomonas* sp., *PhD Thesis*, Cochin University of Science and Technology, Cochin. India. 1999.
20. Kiruthika, J., Saraswathy, N. 2013. Production of L-glutaminase and its optimization from a novel marine isolate *Vibrio azureus* JK-79. *African journal of biotechnology*. 12,(50):6944-6953.
21. Krishnakumar S, Rajan RA, Ravikumar S. Extracellular production of L- Glutaminase by marine alkalophilic *Streptomyces* sp.-SBU1 isolated from Cape Comorin coast. Indian Journal of Geo-Marine Sciences. 2011;40(5): 717-721.
22. Krishnan N, Nath DS, Jayaraman A, Muthusamy P. (): Optimised production of L-glutaminase: A tumour inhibitor from *Aspergillus flavus* cultured on agroindustrial residues. African Journal of Biotechnology. 2011;10(63):13887-13894.

23. Kumar L, Singh B, Adhikari DK, Mukherjee J, Ghosh D. (): A Temperature and Salt-Tolerant L- Glutaminase from Gangotri region of Uttarakhand Himalaya : Enzyme Purification and Characterization. *Appl Biochem Biotechnol*. 2012;166:1723-1735.
24. Kumar SR, Chandrasekaran M. Continuous production of L- glutaminase by an immobilized marine *Pseudomonas* sp. BTMS-51 in a packed bed reactor. *Process Biochemistry*, 2003;38: 1431–1436.
25. Kumar SS, Muthuvelayudham R, Viruthagiri T. Medium optimization for production of l- glutaminase (EC 3.5.1.2) by *Streptomyces griseus* under submerged fermentation. *International Journal of Science and Engineering Applications*. 2013;2(1).
26. Lyer P, Singhal R. Screening and Selection of Marine Isolate for L-Glutaminase Production and Media Optimization Using Response Surface Methodology. *Applied Biochemistry and Biotechnology*, 2009;159: 233–250.
27. Lyer P, Singhal R.. Glutaminase production using *Zygosaccharomyces rouxii* NRRL-Y 2547: Effect of aeration, agitation regimes and feeding strategies. *Chemical Engineering, Technology*, 2010;33: 52-62.
28. Loeliger J. Function and importance of glutamate for savory foods. *J. Nutr*. 2000;130:915–920.
29. Lukey MJ, Wilson KF, Cerione RA. Therapeutic strategies impacting cancer cell glutamine metabolism. *Future Med. Chem*. 2013;5(14), 1685–1700.
30. Nagaraju K, Ram MR. (: Studies on optimization of l-glutaminase from *Bacillus endophyticus* RN-7. *Int J Pharm Biol Sci*. 2018;8(3): 1146-1153.
31. Nathiya K, Nath SS, Angayarkanni J, Palaniswamy M. Optimised production of L-glutaminase: A tumour inhibitor from *Aspergillus flavus* cultured on agroindustrial residues. *African journal of biotechnology*. 2011;10(63): 13887-13894.
32. Ohshima M, Yamamoto T, Soda K. Further Characterization of Glutaminase Isozymes from *Pseudomonas aeruginosa*. *Agricultural and biological chemistry Journal*. 1976;40(11):2251-2256.
33. Pallem C, Manipati S, Somalanka SR. Process optimization of l-glutaminase production by *Trichoderma koningii* under solid state fermentation (ssf). *International Journal of Applied Biology and Pharmaceutical Technology*. 2010;1(3):1168-1174.
34. Prakash PJ, Poorani E, Anantharaman P, Balasubramaniam T. L-glutaminase production and the growth of marine bacteria. *Research journal of microbiology*, 2009;4(4), 168-172.
35. Rashmi AM, Gopinath SM, Kumar K, Murthy NTP. Optimization of submerged fermentation process for L- glutaminase produced by *Pseudomonas aeruginosa* BGNAS-5. *International journal of latest research in science and technology*. 2012;1(4):308-310.
36. Revanth B, Raju K. L-glutaminase production by *Aspergillus wentii* MTCC 1901 under solid state fermentation using mixed agro industrial residues, *International journal of chemical science*, 2013;11(1):277-290

37. Roberts J, Holcenberg JS, Dolowy WC. Isolation, crystallization and properties of Achromobacteraceae glutaminase-asparaginase with antitumor activity. J Biol Chem., 1972;247:84-90.
38. Roberts J, McGregor WG. Inhibition of mouse retroviral disease by bioactive glutaminase-asparaginase. Journal of General Virology. 1991;72, 299-305.
39. Rosenberg J, Ischebeck T, Commichau FM. Vitamin B6 metabolism in microbes and approaches for fermentative production. Biotechnol Adv. 2017;35(1): 31-40
40. Sabu, A. L- glutaminase production by marine fungi. PhD. thesis. Cochin university of science and technology. India. 1999.
41. Sabu A, Keerthi TR, Rajeev KS, Chandrasekaran M. L-Glutaminase production by marine *Beauveria* sp. under solid state fermentation. Process Biochemistry, 2000;35: 705-710.
42. Sathish T, Lakshmi GS, Rao CS, Brahmaiah P, Prakasham RS. Mixture design as first step for improved glutaminase production in solid-state fermentation by isolated *Bacillus* sp. RSP-GLU. Lett. Appl. Microbiol. 2008;47(4): 256–262.
43. Sato I, Kobayashi H, Hanya Y, Abe K, Murakami S, Scorzetti G. *Cryptococcus nodaensis* sp nov, a yeast isolated from soil in Japan that produces a salttolerant and thermostable glutaminase. Journal of Industrial Microbiology and Biotechnology. 1999;22,127–132.
44. Sameera V, Raju KJ. Optimization of process parameters for the production of l-glutaminase with mixed substrate by solid state fermentation using *Aspergillus wentii* MTCC 1901. International Journal of Research in Engineering and Technology. 2015;4(5): 328-333.
45. Saptarshi SD, Lele SS. Localization and disruption kinetics of L-asparaginase from *Erwinia carotovora* cells by high power ultrasound. Chem. Biochem. Eng. Q., 2011;25: 65–73.
46. Shirazia, P, Asad S, Amoozegar MA. The potential of halophilic and halotolerant bacteria for the production of antineoplastic enzymes: l-Asparaginase and l-glutaminase. EXCLI Journal. 2016;15:268-279.
47. Sood CR, Chanda SV, Singh Yd. (): Effect of different nitrogen sources and plant growth regulators on glutamine synthetase and glutamate synthase activities of radish cotyledons. Bulg. J. Plant Physiol. 2002;28(3-4):46–56.
48. Suresh. Chitinase production by marine fungi. PhD. Thesis. Cochin university of science and technology. Cochin, India. 1996.
49. Suresh KS, Muthuvelayudham R, Viruthagiri T. Medium optimization for production of L-glutaminase (EC 3.5.1.2) by *Streptomyces griseus* under submerged fermentation. International Journal of Science and Engineering Applications. 2013;2(1), 2319-7560.
50. Wakayama M, Yamagata T, Kamemura A, Bootim N, Yano S. Characterization of salt-tolerant glutaminase from *Stenotrophomonas maltophilia* NYW-81 and its application in Japanese soy sauce fermentation. The Journal of Industrial Microbiology and Biotechnology, 2005;32:383-390.

51. Zhu Y, Rizeman A, Tramper J, Bol J. Medium design based on stoichiometric analysis of microbial transglutaminase production by *Streptovercillium mobaraense*. *Biotech Bioeng.* 1996;50:291–298.