Original Research Article EVALUATION OF PATTERN OF ANTIHYPERTENSIVE PRESCRIPTIONS AND ADHERENCE TO JNC-7 GUIDELINE IN NATIONAL HOSPITAL ABUJA-NIGERIA

Abstract

Aim: To evaluate the pattern and acquisition costs of antihypertensive medications prescribed as well as adherence to JNC-7 guideline in management of hypertension.

Study Design: Retrospective cross-sectional.

Place and Duration of Study: Conducted at the Cardiology clinic of National Hospital Abuja-Nigeria using case notes of hypertensive patients that were seen from May, 2015 to April, 2016. *Methodology:* Patients variables were collected using standardized data collection form. The average drug acquisition costs (ADAC) were calculated for each antihypertensive drug class Adherence to the guideline was calculated as a percentage of the total number of patients' prescriptions that follow JNC-7 the guideline.

Results: The average age of the study population was 47.05 ± 8.59 years. The mean number of drugs per prescription in the study was 1.82 ± 0.64 . There were no significant differences in the demographic data, with the exception of body mass index (BMI and) Age group (*P*<0.05). 44.3% of the patients were on combination therapy, while 30.5% and 25.2% were on monotherapy and fixed dose combination respectively. The most commonly prescribed antihypertensive drug class was Calcium Channel Blockers (36.76%). There was significant difference in blood pressure outcome among patient on combination therapy and monotherapy as well as among patients whose prescription followed and did not follow the guideline (*P*<0.05). Adherence to the guideline occurred in 82.39% of the patients. The diuretics had the least acquisition cost in Naira (Cost per day: 5.89 ± 2.87 ; Cost per year: 2129.02 ± 1080.49) in relation to the other

antihypertensive prescribed. Similarly, monotherapy had the least acquisition cost in type of therapy.

Conclusion: Calcium channel blockers were the commonly prescribed antihypertensive medications. Alpha-2 adrenergic receptor agonist had the lowest cost utilized per year. There was better blood pressure control in patients whose prescriptions followed JNC-7 guideline.

Keywords

Antihypertensives; Prescriptions; Patterns; Monotherapy; Cost

Introduction

1.1 Background

Hypertension is defined as a systolic blood pressure (SBP) of 140 mm Hg and above and or diastolic blood pressure (DBP) of 90 mm Hg in an adult aged 18 years and above (1). The most common infectious diseases such as malaria, tuberculosis and HIV/AIDS together with childhood and maternal death are still the major causes of mortality within Nigeria and Africa as a whole (2). Nonetheless, worldwide, the burden of hypertension and other non-communicable diseases (NCDs) is swiftly rising, and the African continent may be highly affected compared to other regions of the world (3). The increasing number of cases of hypertension has been attributed to population growth, ageing, and modifiable risk factors such as excessive smoking and alcohol intake, sedentary life style and lack of physical exercise (4). The United Nations and foremost public health stakeholders have affirmed NCDs a cause for worldwide concern (3,5). Hypertension is the foremost modifiable risk factors for cardiovascular diseases (CVD), stroke and renal failure (6). It is the second leading cause of chronic kidney disease (CKD) (7). The proportion of the global burden of disease attributable to hypertension significantly increased

from about 4.5 percent (nearly1 billion adults) in 2000 (Kearney *et al.*, 2005), to 7 percent in 2010 (8). Furthermore, recent study reports have shown the prevalence of hypertension to be increasing (9). This makes hypertension the single most important cause of morbidity and mortality globally and highlights the urgent need of action to address the problem (9).

1.2 Statement of the Research Problem

Liu and Wang (2008), in a study conducted in Taiwan, demonstrated that in 6,536 newlydiagnosed cases of uncomplicated hypertension, CCBs and BBs were the most prescribed antihypertensive medications (10). Surprisingly, the prescription rate of thiazide diuretics which are the least expensive, and well-known first-line antihypertensive therapy was low (8.3% monotherapy and 19.9% overall). AlDrabah et al. (2012) observed that majority of patients in their study were prescribed monotherapy, followed by two drugs. A few others required at least three drugs. While ACEIs were the most commonly prescribed monotherapy, diuretics were the most commonly prescribed drugs in combination therapy. These researchers further observed that target BP control was not achieved in most patients which implied that monotherapy might not be sufficient for achieving adequate BP control in majority of the patients (11).

Similarly, the availability of hypertension management guidelines and evidence showing that hypertension is a major public health concern, nevertheless many physicians fail to assess blood pressure routinely, and in those with diagnosis of hypertension, they do not start treatment or titrate the dosage of antihypertensive medications effectively (12). Studies conducted in USA and Malaysia concluded that physicians are not adhering to evidence--based practice guidelines, which could be possible explanation for poor blood pressure control (13). A study conducted in USA revealed that only 50% of adults with hypertension have their blood pressure under control (14). Ardery et al. (2007) concluded that, guideline adherence among providers was nearly 54%,

and there was no significant improvement over time (15). The study was conducted in 6 community clinics in Iowa. Despite the fact that the JNC has been publishing evidence--based guidelines since 1977, there are still a number of providers that remain unaware or unfamiliar with the JNC guidelines. A significant number of providers are also using much higher blood pressure thresholds than the recommended 140/90 mm Hg to diagnose and treat hypertension (16,17). In one study, some providers didn't initiate therapy for their patients unless the SBP was greater than 160 mm Hg or the DBP was more than 95 mm Hg (16).

In Nigeria, the first hypertension management guideline was published in 1996, since then it has not been reviewed and updated (18). Also, the 2008 Standard Treatment Guidelines (STGs) has not been updated and the guidelines consider BBs as superior to ACEIs and ARBs in managing hypertension without compelling indication (FMOH, 2008). The lack of updated guidelines perhaps compelled Odili et al. (2008) to used international guidelines such as WHO/ISH, and ESH in the evaluation of the role of physicians in the overall management of hypertension and their adherence to treatment guidelines in a study conducted in University of Benin Teaching Hospital in southern part of the country (19). The study concluded that Physicians fairly complied with the stated guidelines. However, they did not appear to recommend lifestyle modification to their patients. Practice guidelines serve as useful tools for clinical decision making. They also help to reduce the variation in practice, guide appropriateness and measure quality of care.

The per capita income in Nigeria as at 2012 was \$2700.00 (two thousand seven hundred US Dollars). Also the minimum monthly wage is =N=18000.000 (eighteen thousand Naira equivalent of US\$120.00 (one hundred and twenty US Dollars at an exchange rate of =N=150 to the Dollar as at year 2012). These figures are low and if a big percentage of this monthly income

is spent on buying medicines for the management of hypertension, it leaves a very little amount for other expenses (20). The implication of this may be that patients may prioritize their spending and forego the purchase of the medications in order to buy food and other essential supplies for the family. The cost of prescription medications is thought to be a barrier for many patients to access the healthcare they need.

Jackson and co-workers observed that applying JNC-7 guidelines to clinical practice helped in achieving better BP control. Blood pressure control before the JNC-7 cohort was 40.8 % vs. 49.3 % after JNC-7 cohort (P<0.0001) (21). In another study by Abdulameer et al. (2012) 85.30 % of the prescriptions were in accordance to guidelines. It was observed that the treatment approach for cardiac complicated hypertension followed JNC-7 guidelines, except the lack of add-on therapy practice (ARBs, aldosterone antagonist) (13).

2.0 Material and Methods

2.1 Instruments

- 1. Standardized data collection form.
- 2. Patient case notes.
- 3. The seventh report of Joint National Committee (JNC-7) on prevention, detection, evaluation and treatment of high blood pressure guideline for hypertension.
- 4. National Hospital Drug Price List.

2.2 Study Design

A retrospective cross-sectional study was carried out at the cardiology clinic of NHA using data collection form. The case notes of hypertensive patients that were seen from May, 2015 to April, 2016 at the Cardiologic clinic were used

2.3 Study Population

All hypertensive patient case notes with diagnosis of only hypertension and hypertension with co-morbidities of diabetes and ischemic heart diseases that have attended the MOPD clinic of NHA within the study period.

2.4 Study Criteria

2.4.1 Inclusion criteria

1. Adult male and female patients diagnosed of hypertension aged 18 years and above.

2. Hypertensive patients with co-morbidity of diabetes mellitus and/or ischemic heart disease. Study findings in Nigeria, have shown the co-morbidity of type-2 diabetes mellitus and ischemic heart diseases to be very common in hypertension (20,22).

2.4.2 Exclusion criteria

1. Hypertensive patients with co-morbidities other than diabetes mellitus and/or ischemic heart disease.

- 2. Patients less than 18 years old.
- 3. Pregnant women.
- 4. Lactating mothers.

2.5 Sample Size and Sampling Method

2.5.1 Sample size calculation

The sample size was determined using the formula for estimating a single population proportion

(23), as follows:

 $n = Z^2 p(1-p)/w^2$

Where:

n=the required sample size

Z= the value of Z in the standard normal distribution which corresponds to a level of 0.05 with a value of 1.96.

p= prevalence of hypertension in Nigeria which was estimated at 25.1 to 32.8% (5). Hence taking the lower value, p=25.1%=0.251

w= the margin of error (precision)
$$=5\%=0.05$$

 $n=(1.96)^2 x 0.251 (0.749) / (0.05)^2 = 289$

Taking a 10% attrition rate, the required sample was = 289+29=318.

2.5.2 Sampling method

Selection of patient folders was done by the principal investigator. A systematic random sampling procedure was adopted among patient folders from May, 2015 to April, 2016. The sampling frame was defined as a function of cumulative total number of hypertensive patients seen over a 12 months period (N) divide by the sample size (n). Since the minimum sample size had been adjusted to 318, the sampling fraction therefore was determined as:

 $(N/n)= 600/318= 1.9 \approx 2$. The first folder from the month of May, 2015 was selected and subsequently every other folder was selected thereafter until the required sample size (318) was attained.

2.6 Data collection

Demographic data, diagnoses and drug prescribed were collected using data collection form (Appendix: I) from medical records of patients who attended the MOPD clinic during the study periods.

2.6.1 Prescribed antihypertensive medications

The antihypertensive drugs were classified into six therapeutic drug classes using JNC-7 guideline: (1) Beta-blockers (BBs), (2) Calcium-channel blockers (CCBs), (3) Diuretics (D), (4) Angiotensin-converting enzyme inhibitors (ACEIs), (5) Angiotensin-receptor blockers (ARBs) and (6) Alpha-2 Adrenergic receptor agonist (A_2AA).

Monotherapy was defined as the use of one antihypertensive medication. Patients who were prescribed more than one medication in one combination were defined as receiving fixed dose combination (FDC). While receiving more than one antihypertensive in different pills was defined as receiving combination therapy (24).

2.6.2 Drug cost

The individual drug cost was obtained from the Pharmacy Department (National Hospital Abuja) and entered into the data collection form against the prescribed antihypertensive medications. The average drug acquisition costs (ADAC, the cost of buying the drugs) was calculated for each drug class on a daily and annual bases and the percentage expenditure cost (%EC) also, was determined, using the following equations (13).

a. ADAC (daily) =
$$\frac{Total \ daily \ cost \ of \ drug \ class}{Number \ of \ prescriptions \ of \ the \ drug \ class}$$

b. ADAC (Annual) = ADAC (daily)

c.
$$%EC = \frac{EC \text{ of the drug class}}{Total EC \text{ of all prescriptions}} \times 100$$

Where EC is expenditure cost.

2.6.3 Adherence to JNC-7 guideline

Adherence to the guideline report by the panel appointed to the Seventh Joint National Committee (JNC-7) for the management of Hypertension was expressed as a percentage of the total number of patient (13). The percentage adherence was calculated as follows:

1. The first step was to specify the treatment guideline for each group of patients:

- a. Hypertension only (use of D or CCB, ACEI, ARB as a first-line therapy or added onto another drug therapy).
- b. Hypertension with Diabetes mellitus (use of an ACEI or ARB as a first-line therapy or added onto another antihypertensive drug therapy).
- c. Hypertension with Ischemic heart disease (use of a BB or CCB as a first-line therapy or added onto another antihypertensive drug therapy).
- d. Hypertension with Ischemic heart disease and Diabetes mellitus (use of an ACEI or ARB and BB or CCB as a first-line therapy or added onto another antihypertensive drug therapy).
- 2. In the second step, percentage of adherence (%A) to the specified guidelines was calculated:

$$\%A = \frac{Total number of cases that followed the guideline}{Total number of prescriptions} \times 100$$

2.7 Validation of Study Instrument

The designed data collection form was pretested in the hospital. The pretesting was undertaken with 10% of the sample size (later discarded) before the actual data collection took place. This was done in order to ensure that necessary data were collected.

2.8 Data Management and Analysis

The data were entered in excel and exported to statistical package for social sciences (SPSS) computer software version 20.0 and analyzed using Independent *t*-test, Paired t-test, Chi-square Anova and Tukey HSD post-hoc test. Significant difference was considered at p<0.05.

2.9 Ethical Approval

Ethical approval was obtained from the Research and Ethics Committee of National Hospital

(Appendix 1).

3.0 Results and Discussion

3.1 Results

3.1.1 Patient Demographic Characteristics

Out of 318 folders selected based on the selection criteria, 46.4% were male and 53.6% female. The difference in their ages was not statistically significant (*P*>0.05) (Table 1).

Variables	Male	Female	All patients
Gender (%)	148(46.4)	170(53.6)	318(100)
Age-group			
(years)			
29-43	46(31.1)*	$70(41.2)^{*}$	116(36.5)
44-58		73(42.9)*	166(52.2)
≥59	93(62.8)*	27(15.9)*	36(11.3)
	9(6.1)*		
Mean age in	47.3	46.9	47.1
years (±SD)	(±8.3)	(±8.9)	(±8.6)
Median Age	49	48	49
(years)			
BMI in kg/m ²	30.80	29.26	29.98
(±SD)	$(\pm 4.5)^{\ddagger}$	$(\pm 6.9)^{\ddagger}$	(±5.9)
Level of			
Education			
Some Levels of	22(6.9)	17(5.4)	39(12.3)
Education			
Secondary	18(5.7)	29(9.1)	47(14.8)
Tertiary	91(28.6)	108(33.9)	199(62.6)
Postgraduate	17(5.4)	16(5.0)	33(10.3)

Note:

*There is significant difference when Age group values are compared between genders using chi-square, P = 0.001.

 \ddagger =There is significant difference when mean BMI value of males and females are compared using independent t-test, P = 0.02.

BMI- Body Mass Index, SD- Standard deviation

3.1.2 Patients' Average Blood Pressure with Mean Number of Antihypertensives per Prescription

Both systolic blood pressure (SBP) and diastolic blood pressure (DBP) were not significantly

different at baseline (p>0.05). There was a significant reduction in systolic and diastolic blood

pressure in both sexes (p < 0.05) after initiation of treatment (Table 2).

Table 2: Patients' Average Blood Pressure with Mean Number of
Antihypertensive Medications Prescribed to Each of 318 Hypertensive
Studied Patients

Gender	PrTSBP mmHg (±SD)	PTSBP mmHg (±SD)	PrTDBP mmHg (±SD)	PTDBP mmHg (±SD)	Number of Antihypertensives per prescriptions
Male	161.2	130.8	96.1	81.9	1.77
	(±13.8)	$(\pm 14.7)^{*}$	(±8.5)	$(\pm 7.1)^{\dagger}$	(±0.6)
Female	160.3	131.7	96.8	82.9	1.86
	(±14.1)	$(\pm 16.0)^{\ddagger}$	(±8.4)	(±8.2) [↑]	(±0.7)
Mean	160.7	131.3	96.5	82.4	1.82
BP	(±13.9)	(±15.4)	(±8.5)	(±7.7)	(±0.6)

Note:

SD= Standard deviation

*P< 0.05, PrTSBP vs PTSBP in males (paired t-test).

 $\dagger P < 0.05$, PrTSBP vs PTSBP in females (paired t-test).

 $\ddagger P < 0.05$, PrTDBP vs PTDBP in males (paired t-test).

 \uparrow *P*< 0.05, PrTDBP vs PTDBP in females (paired t-test).

PrTSBP=Pretreatment systolic blood pressure

PrTDBP=Pretreatment diastolic blood pressure

PTSBP=Post-treatment systolic blood pressure

PTDBP=Post treatment diastolic blood pressure

3.1.3 Antihypertensive Medications Prescribed at the Cardiology Clinic

Calcium Channel Blockers (CCBs) were commonly prescribed and least prescribed were Alpha-2 adrenergic receptor agonists (A₂AA) (Table 3).

Types of antihypertensives	No. Prescriptions	%
prescribed		
BETA BLOCKERS		8.71
Atenolol	32	5.57
Metoprolol	7	1.22
Bisoprolol	5	0.87
Carvedilol	5	0.87
Propranolol	1	0.17
CCBs		36.76
Amlodipine	153	26.66
Nifedipine	58	10.10
DIURETICS		22.30
Hydrochlorthiazide	101	17.60
Spironolactone	7	1.22
Moduretic	20	3.48
ACEIs		14.46
Lisinopril	61	10.63
Ramipril	15	2.61
Enalapril	7	1.22
ARBs		13.59
Losartan	2	0.35
Valsartan	43	7.49
Telmisartan	24	4.18
Candesartan	9	1.57
A2AA		4.18
Methyldopa	24	4.18
Total	574	100.00

 Table 3: Antihypertensive Medications Prescribed Among the 318 Hypertensive Studied Patients

KEY TO ABBREVIATION

CCBs- Calcium channel blockers, ACEIs- Angiotensin converting enzyme inhibitors, ARBs- Angiotensin II receptor antagonists, A₂AA- Alpha-2 adrenergic receptor agonists

3.1.4 Gender Distribution of Blood Pressure Stages among the Studied Hypertensive Patients

Only 1.6% of the patients were pre-hypertensive at diagnosis and 80% of the patients were

female. Also, 28.9% of the studied patient had stage-1 hypertension, out of which 53.3% were

males. However, 55.7% of the stage-2 hypertensive patients were female (Table 4).

Patients				
BP Category	SBP/DBP (mmHg)	Male (%)	Female (%)	Total no of Patients (%)
Pre	120-139/80-89	1(20)	4(80)	5(1.6)
Hypertension Stage-1 Hypertension	140-159/90-99	49(53.3)	43(46.7)	92(28.9)
Stage-2 Hypertension	≥160/100	98(44.3)	123(55.7)	221(69.5)

 Table 4: Gender Distribution of Stages of Hypertension According

 to JNC-7 Treatment Guideline Among the 318 Studied Hypertensive

 Patients

Note:

BP= Blood Pressure

SBP= Systolic blood pressure

DBP= Diastolic blood pressure

3.1.5 Diagnosis and Prescriber Adherence to JNC-7 guideline with Blood Pressure Outcome in the 318 Studied Hypertensive Patients in the Cardiology Clinic, National Hospital Abuja

Patients' blood pressure after commencement of therapy based on prescribed antihypertensives using JNC-7 guideline is shown below (Table 5).

Diagnosis	No of Prescriptions following guideline	BP after commencement of therapy following guideline (mean ± SD)	No of Prescriptions not following guideline	BP after commencement of therapy not following guideline (mean ± SD)	Total Adherence (%)
HT	149	$129.8 \pm 6.8^*$	44	$141 \pm 5.5^{*}$	77.20
		$83.1 \pm 7.5^{\circ}$		$85.8 \pm 6.5^{\dagger}$	
HT + DM	70	$128.2 \pm 9.3*$	7	135.7±2.9*	90.91
		79.2 ± 6.1		82.9 ± 1.9	
HT + IHD	27	$124.4 \pm 8.5*$	2	$143 \pm 2*$	93.10
		$82.4\pm4.5^{\dagger}$		$87\pm3^{\dagger}$	
HT + DM	16	125.3 ± 8.4	3	133.7 ± 2.6	84.21
+ IHD		$80.1 \pm 5.7^{\dagger}$		$88\pm2^{\dagger}$	
Total	262		56		82.39

 Table 5: Diagnosis and Prescriber Adherence to JNC-7 Guideline with Blood Pressure

 Outcome in the 318 Studied Hypertensive Patients

Note:

BP-Blood pressure in mmHg (systolic and diastolic).

*= There is significant difference when mean systolic BP following treatment based on guideline and not based on guideline are compared using independent t-test among the hypertensive patient categories, P=0.0001 (HT); P=0.038 (HT+DM); P=0.005 (HT+IHD).

 †= There is significant difference when mean diastolic BP following treatment based on guideline and not based on guideline are compared using independent t-test among the hypertensive patient categories, P=0.03 (HT);
 P=0.024 (HT+IHD); P=0.033 (HT+DM+IHD).

H- Hypertension; DM- Type-2 diabetes mellitus; IHD- Ischemic heart disease.

3.1.6 Drug Acquisition (daily and annual) Costs of the Different Antihypertensive Drugs and Type of Therapy Prescribed in the Cardiology Clinic, National Hospital Abuja

The drug utilization and costs of different antihypertensive agents incurred per day as well as per year were represented in (Table 4.7). The diuretics had least acquisition cost (Cost per day: NGN6.4 ± 5.7; Cost per year: NGN2325.0 ± 2073.5) in relation to the other antihypertensive drug classes prescribed. The most expensive were the angiontensin II receptor blockers (P<0.05). Also, comparing the acquisition costs of modes of therapy, lower costs were incurred with monotherapy followed by fixed dose combination (FDC) while combination therapy was the most expensive (P<0.05). (Table 6).

Drug Class	No of Prescriptions	Average daily acquisition cost per prescription (mean ± SD)	Average annual acquisition cost per prescription (mean ± SD)
CCBs	211	34.5 ± 8.9	12592.0 ± 3259.5
Diuretics	128	6.4 ± 5.7	$2325.0 \pm 2073.5^{*}$
ACEIs	83	26.3 ± 2.2	$9610.5 \pm 806.7^{*}$
ARBs	78	127.4 ± 45.8	$46515.6 \pm 16698.8^{*}$
BBs	50	24.7 ± 8.9	$9015.5 \pm 3259.5^{*}$
A ₂ AA	24	15.0 ± 2	$5475.0 \pm 730^{*}$
Type of therapy			
Monotherapy	97	35.4 ± 10.1	$12928.3 \pm 3682.9^{\dagger}$
Combination therapy	141	134.8 ± 19.2	$49183.8 {\pm} 7004.4^{\dagger}$
Fixed Dose Combination	80	114.2 ± 19.7	$41675.7 \pm 7194.2^{\dagger}$
Note:			

Table 6: Drug Acquisition Costs (daily and annual) of Antihypertensive Dru	ıg
Class and Type of Therapy Prescribed Among the 318 Studied Hypertensive	e

Drug costs are in naira (NGN);

*=There is significant difference between average annual acquisition costs of drug class, when drug classes are compared using ANOVA test, P = 0.001.

 \dagger =There is significant difference between average annual acquisition costs of type of therapy, when modes of therapy are compared using ANOVA, P= 0.002.

3.1.7 Drug Utilization and Expenditure Cost

Alpha-2 adrenergic receptor blockers had the least percentage expenditure cost (0.14%) then diuretics (3.79%) with the corresponding percentage utilization of 4.18% and 22.3% respectively. ARBs had the highest percentage expenditure cost (46.27%) then CCBs (33.88%) (Figure 1).



Figure 1: The bar chart showed the utilization and expenditure patterns of antihypertensive medications in the cardiology clinic, National Hospital Abuja

Note:

*=There is significant difference when percentage drug utilization and percentage drug expenditure cost are compared with the antihypertensive drug class prescribed to the patients using chi-square, P = 0.001.

4.0 Discussion

The findings of our study revealed a mean age that was lower than the reported result from previous studies done in northern and southwestern Nigeria (25,26), but, however, significantly higher than reported findings of Egbi and co-workers in a study conducted in Bayelsa State, Niger-Delta region of the country (27). Most of the patients were below 59years of age (Table 1), which was in some way similar to the findings of Abdulameer and co-workers (13). This could be due to the fact that most of the patients in the study center were active civil or public servants that have not attained mandatory retirement age of 60years. The prevalence of hypertension was seen more in females compared to males (1.15:1), which corresponds to the findings of Camara

et al. (2015), and Musinguzi and Nuwaha (2013) (28,29). The studies were conducted in urban and rural settings of republic of Guinea and Uganda respectively. The difference in gender prevalence may be due to the fact that males do not regularly keep clinic appointments (30), or it may be a reflection of poorer awareness of BP status among men (31).

The mean number of antihypertensive drugs per prescription in this study (Table 2) was lower than reported values from studies conducted in other developing countries (2.2-3.8 drugs per prescription) such as Jordan, Malaysia (13,32). This may be due to the use of fixed dose combination in our study center. Most patients presented to the hospital with a mean blood pressure (BP) consistent with the stage 2 hypertension classification (Table 2). This is similar to the finding of studies done in Ilorin, Nigeria and Eastern Central Region of Portugal (33,34). In about 70% of the studied population the pretreatment BP classifications were in stage 2 category. This is disturbing because the risk of cardiovascular diseases is doubled in these patients (35). Therefore, blood pressure control is highly important in preventing the development cardiovascular diseases, stroke and renal failure. The patients in the pre-hypertension, who were placed on drugs, were so treated probably because of the presence of comorbidities. This is in keeping with the treatment guidelines which state that the presence of co-morbidities in prehypertension necessitates the use of drugs (19). Also, in this study, blood pressure at diagnosis and after treatment appeared to be the same in both sexes (Table 2). However, a study conducted in University of Benin Teaching Hospital revealed a significantly higher diastolic blood pressure at diagnosis (19).

The finding that CCBs were the most commonly prescribed drug in this study (Table 3), agrees with reports of studies conducted outside Nigeria (36,37). This was not the case in a study done in Lagos, Nigeria where thiazide was the most commonly prescribed antihypertensive (38). The

Lagos study result may probably have been influenced by ALLHAT study findings released in 2002. The ALLHAT study had emphasized the use of thiazide-like diuretic in its recommendations, and this has been reported to increase prescription pattern of the drug by physicians (39,40). The difference in the level of thiazide use in this study and the Lagos study, could be due to the influence of JNC-8 treatment guidelines, which suggest that diuretics or calcium channel blockers (CCBs), are suitable as initial drug treatment of hypertension in blacks, either as monotherapy or in combination therapy (1).

The ranking in terms of cost from the highest to the lowest was: ARB > CCB > ACEI >BBs >D $>A_2AA$ (Figure 1). Therefore, the antihypertensive acquisition cost analysis showed that ARBs had the highest drug cost compared to utilization followed by CCBs. The explanation of high cost is that, the use of innovator drugs adds more cost on drug expenditure per prescription that is available in National Hospital. Similarly, result of cost evaluation showed that diuretics had the lowest cost per day (CPD) -NGN (6.37 ± 5.68) and cost per year (CPY) -NGN (2325 ± 2073.5) followed by alpha₂ adrenergic agonists (Table 6). These results were comparable with other studies which evaluated the cost effectiveness of the available antihypertensives in India and Malaysia (13,41). The use of thiazides as a first-choice therapy would result in substantial cost savings due to their favorable price (42). The comparison of acquisition cost of therapy types showed that monotherapy was least costly compared to fixed dose combination (FDC) and combination therapy (Table 6). This finding was supported by results of other studies which observed that patients on three or four drug regimens had significantly higher treatment costs (41,43). Furthermore, there was a decrease in the treatment costs of hypertension with a coexisting disease in cases where the guideline was not followed (Table 5), indicating inadequate treatment. In contrast, in the treatment of essential hypertension, there was an increase in costs in

cases where the guidelines were not followed, which indicates that more expensive drugs were prescribed instead of the first-line treatment consisting of cheaper drugs such as diuretics. Similar findings had been reported somewhere (13).

4.0 Conclusions and Recommendation

Calcium channel blockers were the commonly prescribed antihypertensive medications. Alpha-2 adrenergic receptor agonist had the lowest cost utilized per year. There was better blood pressure control in patients whose prescriptions followed JNC-7 guideline. Consequently, prescribing antihypertensive medications using JNC-7 hypertension management guideline recommendations is associated with better blood pressure outcomes. Hence, it is good to follow the guideline while prescribing antihypertensive medications to hypertensive patient.

Appendix 1 (Ethical Approval)



NAL HOS

(Established by Act No 36 of 1999)

BOARD CHAIRMAN:

Ag. DIRECTOR OF ADMINISTRATION Alh. Y. O. Sadiq, Bsc (Soc), Msc (Soc), AHSAN, MNIPR

NHA/ADMIN/236/V.VII

CHIEF MEDICAL DIRECTOR / CEO Dr. J. A. F. Momoh, MBBS, MSC, FWACP(LM)

DIRECTOR OF CLINICAL SERVICES/CMAC Dr. Oluseyi Oniyangi, MBBS, FWACP, (Pead) FIPNA

20th July, 2016

RE: AN EVALUATION OF PATTERNS OF ANTIHYPERTENSIVES USE AND ADHERENCE TO GUIDELINES IN HYPERTENSIVE MANAGEMENT IN NATIONAL HOSPITAL ABUJA NIGERIA. NHA/EC/042/2016

Health Research Ethics Committee (HREC) Assigned number:

Name of Principal Investigator:

Address of Principal Investigator

Pharm. Abdullahi Yusuf

NHA/EC/042/2016

Department of Pharmacy & Pharmacy Practice Faculty of Pharmaceutical Science ABU, Zaria.

15th June, 2016

Date of Receipt of Valid Application:

Notice of Approval

This is to inform you that the research described in the submitted protocol, the consent forms, and other changes stated in the submitted research protocol addendum have been reviewed and given full approval by the Institutional Review Board (IRB) Committee, National Hospital Abuja.

This approval dates from 20th July, 2016 to 19th July, 2018. If there is delay in starting the research, please inform the HREC National Hospital Abuja so that the dates of approval can be adjusted accordingly. Note that no participant accrual or activity related to this research may be conducted outside of these dates. All informed consent forms used in this study must carry the HREC assigned number and duration of HREC approval of the study.

The National Code for Health Research Ethics requires you to comply with all institutional guidelines, rules and regulations and with the tenets of the Code including ensuring that all adverse events are reported promptly to the HREC. No changes are permitted in the research without prior approval by the HREC except in circumstances outlined in the Code. The HREC reserves the right to conduct compliance visit to your research site without previous notification.

Dr. Oluseyi Oniyangi (DCS/CMAC) For: Chairman, HREC, National Hospital

Plot 132 Central District (Phase II) P.M.B. 425, Garki - Abuja Nigeria Telephone: 0803-787-9543, 0809-751-9764, 0809-752-0012 E-mail: info@nationalhospitalabuja.net www.nationalhospitalabuja.net

References

- James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. Evidence-Based Guideline for the Management of High Blood Pressure in Adults. JAMA. 2014; 311(5):507. Available from: http://jama.jamanetwork.com/article.aspx?doi=10.1001/jama.2013.284427
- Akinlua JT, Meakin R, Umar AM, Freemantle N. Current Prevalence Pattern of Hypertension in Nigeria: A Systematic Review. Public Library of Science. 2015;10(10):e0140021. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26461923
- WHO. A global brief on hypertension: silent killer, global public health crises. 2013; Available from: http://apps.who.int/iris/bitstream/10665/79059/1/WHO_DCO_WHD_2013.2_eng.pdf
- Bernabé-Ortiz A, Carrillo-Larco RM, Gilman RH, Checkley W, Smeeth L, Miranda JJ, et al. Impact of urbanisation and altitude on the incidence of, and risk factors for, hypertension. BMJ Publishing Group Ltd and British Cardiovascular Society. 2017;103(11):827–33. Available from: http://www.ncbi.nlm.nih.gov/pubmed/28115473
- Adeloye D, Basquill C, Aderemi A V, Thompson JY, Obi FA. An estimate of the prevalence of hypertension in Nigeria: a systematic review and meta-analysis. Journal of hypertension 2015;33(2):230–42. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25380154
- 6. Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Blaha MJ, et al. Heart disease and stroke statistics-: a report from the American Heart Association. Circulation 2014;129(3):e28–292. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24352519
- 7. Gargiulo R, Suhail F, Lerma E V. Hypertension and chronic kidney disease. Elsevier. 2015;61(9):387–95. Available from: http://linkinghub.elsevier.com/retrieve/pii/S0011502915001017
- 8. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease. Lancet. 2010;380(9859):2224–60. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23245609
- 9. Forouzanfar MH, Liu P, Roth GA, Ng M, Biryukov S, Marczak L, et al. Global Burden of Hypertension and Systolic Blood Pressure of at Least 110 to 115 mm Hg, 1990-2015. JAMA. 2017; 317(2):165. Available from: http://jama.jamanetwork.com/article.aspx?doi=10.1001/jama.2016.19043
- Liu P-H, Wang J-D. Antihypertensive medication prescription patterns and time trends for newly-diagnosed uncomplicated hypertension patients in Taiwan. BMC health services research. 2008;8(133):1–11.

- 11. AlDrabah E, Irshaid Y, Yasein N, Zmeili S. Prescription pattern of antihypertensive drugs in Family Practice Clinics at Jordan University Hospital. Medicine Science International Medical Journal. 2012;2(1):1. Available from: http://www.scopemed.org/?mno=28051
- 12. Kotchen TA. The search for strategies to control hypertension. Circulation. 2010;122(12):1141–3. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20823382
- Abdulameer SA, Sahib MN, Aziz NA, Hassan Y, Alrazzaq HAA, Ismail O. Physician adherence to hypertension treatment guidelines and drug acquisition costs of antihypertensive drugs at the cardiac clinic: A pilot study. Patient Preference and Adherence. 2012;6:101–8.
- Nwankwo T, Yoon SS, Burt V, Gu Q. Hypertension among adults in the United States: National Health and Nutrition Examination Survey, 2011-2012. NCHS data brief. 2013;(133):1–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24171916
- 15. Ardery G, Carter BL, Milchak JL, Bergus GR, Dawson JD, James PA, et al. Explicit and Implicit Evaluation of Physician Adherence to Hypertension Guidelines. The Journal of Clinical Hypertension. 2007;9(2):113–9. Available from: http://doi.wiley.com/10.1111/j.1524-6175.2007.06112.x
- 16. Hyman DJ, Pavlik VN. Self-reported hypertension treatment practices among primary care physicians: blood pressure thresholds, drug choices, and the role of guidelines and evidence-based medicine. Archives of internal medicine. 2000;160(15):2281–6. Available from: http://www.ncbi.nlm.nih.gov/pubmed/10927724
- 17. Milchak JL, Carter BL, Ardery G, Dawson JD, Harmston M, Franciscus CL. Physician Adherence to Blood Pressure Guidelines and Its Effect on Seniors. Pharmacotherapy. Blackwell Publishing Ltd. 2008;28(7):843–51. Available from: http://doi.wiley.com/10.1592/phco.28.7.843
- 18. Akinkugbe OO, Salako L a., Mabadeje a. FB, Akinroye KK. Guidelines for the Management of Hypertension In Nigeria. A recommendation for health care providers prepared by the Nigerian Hypertension Society and adopted at the Consensus Meeting sponsored by The Nigerian Heart Foundation. 1996;(May):1–20. Available from: http://www.nigerianheart.org/downloads/management_of_hypertension.pdf
- Odili V, Oghagbon E, Ugwa N, Ochei U, Aghomo O. Adherence to International Guidelines in the Management of Hypertension in a Tertiary Hospital in Nigeria. Tropical Journal of Pharmaceutical Research. 2008;7(2):945–52. Available from: http://www.ajol.info/index.php/tjpr/article/view/14608
- 20. Osibogun A, Okwor TJ. Anti-Hypertensive Prescription and Cost Patterns in an Outpatient Department of a Teaching Hospital in Lagos State Nigeria. Open Journal of Preventive Medicine. 2012;4(4):156–63. Available from: http://www.scirp.org/journal/ojpm
- 21. Jackson JH, Sobolski J, Krienke R, Wong KS, Frech-Tamas F, Nightengale B. Blood pressure control and pharmacotherapy patterns in the United States before and after the

release of the Joint National Committee on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 7) guidelines. Journal of the American Board of Family Medicine. 2008;21(6):512–21. Available from: http://www.ncbi.nlm.nih.gov/pubmed/18988718

- 22. Peter OA, Joshua EF, Daughter OA, Nelson EC. Co-Morbidities of Patients with Hypertension Admitted to Amassoma General Hospital, Bayelsa State South-South of Nigeria. Glob J Pharmaceu Sci. 2017;2(4). Available from: https://juniperpublishers.com/gjpps/pdf/GJPPS.MS.ID.555594.pdf
- 23. Jung S-H. Stratified Fisher's exact test and its sample size calculation. Biometrical journal Biometrische Zeitschrift. 2014;56(1):129–40. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24395208
- 24. Sepehri G, Talebizadeh N, Mirzazadeh A, Mohsenbeigi M. The patterns of antihypertensive drug prescription by cardiologists in Kerman province of Iran, 2006. Pharmacoepidemiology and Drug Safety. 2008;17(2):180–5. Available from: http://doi.wiley.com/10.1002/pds.1521
- 25. Asekun-Olarinmoye E, Akinwusi P, Adebimpe W, Isawumi M, Hassan M, Olowe O, et al. Prevalence of hypertension in the rural adult population of Osun State, southwestern Nigeria. International journal of general medicine. 2013;6:317–22. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23641157
- 26. Saidu H, Karaye K, Okeahialam B. Cardiovascular disease risk factors among subjects with high-normal blood pressure in a Nigerian tertiary health institution. Sahel Medical Journal 2015;18(4):156. Available from: http://www.smjonline.org/text.asp?2015/18/4/156/176593
- 27. Egbi OG, Rotifa S, Jumbo J. Prevalence of hypertension and its correlates among employees of a tertiary hospital in Yenagoa, Nigeria. Annals of African medicine. 2015;14(1):8–17. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25567690
- 28. Camara A, Balde NM, Diakite M, Sylla D, Balde EH, Kengne AP, et al. High prevalence, low awareness, treatment and control rates of hypertension in Guinea: results from a population-based STEPS survey. J Hum Hypertens. 2015;237–44. Available from: http://www.nature.com/doifinder/10.1038/jhh.2015.92
- 29. Musinguzi G, Nuwaha F. Prevalence, awareness and control of hypertension in Uganda. Public Library of Science. 2013;8(4):e62236. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23614041
- 30. Geoghegan T. Why are men reluctant to seek medical help? BBC News. 2009. Available from: http://news.bbc.co.uk/1/hi/magazine/8154200.stm
- 31. Awuah RB, Anarfi JK, Agyemang C, Ogedegbe G, Aikins A de-G. Prevalence, awareness, treatment and control of hypertension in urban poor communities in Accra, Ghana. Journal of Hypertension. 2014;32(6):1203–10. Available from:

http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00004872-201406000-00010

- 32. Tandon VR, Sharma S, Mahajan S, Mahajan A, Khajuria V, Mahajan V, et al. Antihypertensive drug prescription patterns, rationality, and adherence to Joint National Committee-7 hypertension treatment guidelines among Indian postmenopausal women. Journal of mid-life health. 2014;5(2):78–83. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24970986
- 33. Morgado MP, Rolo SA, Pereira L, Castelo-Branco M, Silva A, Oliveira A, et al. Blood pressure control and antihypertensive pharmacotherapy patterns in a hypertensive population of Eastern Central Region of Portugal. BMC Health Services Research. 2010;10(1):349. Available from: http://bmchealthservres.biomedcentral.com/articles/10.1186/1472-6963-10-349
- 34. Oghagbon EK, Okesina AB. Pattern of some risk factors for cardiovascular disease in untreated Nigerian hypertensive patients. West African Journal of Medicine. 2007;25(3):190–4. Available from:http://www.ajol.info/index.php/wajm/article/view/28276
- 35. Chobanian A V, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, et al. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. Hypertension. 2003;42(6):1206–52. Available from: http://www.ncbi.nlm.nih.gov/pubmed/14656957
- 36. Kale A, Maniyar YA. Prescribing Patterns of Antihypertensive Drugs in A Tertiary Care Hospital. Scholars Academic Journal of Pharmacy (SAJP) Sch Acad J Pharm. 2013;2(5):416–8. Available from: www.saspublisher.com
- 37. Mohd, Mateti U V, Konuru V, Parmar MY, Kunduru BR. A study on prescribing patterns of antihypertensives in geriatric patients. Perspectives in Clinical Research. 2012;3(4):139.
- Ganiyu Kehinde A, Suleiman Ismail A. Assessment of antihypertensives utilization in a private teaching hospital in Nigeria. International Journal of Pharmacy and Pharmaceutical Sciences. 2012;4(SUPPL. 5):480–3.
- 39. Furmaga EM1, Cunningham FE, Cushman WC, Dong D, Jiang R, Basile J, Katz LA, Rutan GH, Berlowitz DR, Papademetriou V GP. National utilization of antihypertensive medications from 2000 to 2006 in the Veterans Health Administration: focus on thiazide diuretics. J Clin Hypertens. 2008;10:770–8. Available from: https://www.ncbi.nlm.nih.gov/pubmed/19090878
- 40. Muntner P1, Krousel-Wood M, Hyre AD, Stanley E, Cushman WC, Cutler JA, Piller LB, Goforth GA WP. Antihypertensive prescriptions for newly treated patients before and after the main antihypertensive and lipid-lowering treatment to prevent heart attack trial results and seventh report of the joint national committee on prevention, detection, evaluation. Hypertension. 2009;4:617–23. Available from: https://www.ncbi.nlm.nih.gov/pubmed/19221214

- 41. PR R. Anti Hypertensive Prescribing Patterns and Cost Analysis for Primary Hypertension: A Retrospective Study. Journal of Clinical and Diagnostic Research. 2014;8(9):HC19-22. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25386458
- 42. Ekwunife OI, Okafor CE, Ezenduka CC, Udeogaranya PO. Cost-utility analysis of antihypertensive medications in Nigeria: a decision analysis. Cost effectiveness and resource allocation. 2013;11(1):2. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3567962&tool=pmcentrez&re ndertype=abstract
- 43. Ilesanmi OS, Ige OK, Adebiyi AO. The managed hypertensive: the costs of blood pressure control in a Nigerian town. The Pan African medical journal. 2012;12:96. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23133696