

Original Research Article

LENGTH-WEIGHT RELATIONSHIP AND CONDITION FACTOR OF 26 FISH SPECIES CAUGHT BY CAST NET IN NEW CALABAR RIVER, NIGERIA

Comment [L1]: Modify as suggested in the comments

Abstract

The length-weight relationship and condition factor (K) of fish species caught by cast net were studied from three stations in the New Calabar River, Rivers State, Nigeria. A total of 1541 specimens of 26 fish species and representing 11 families were randomly collected using cast net with mesh sizes of 1.5 and 2.5cm. Total length (TL) was measured to 0.1 cm, while whole body weight (W) was taken to the nearest ± 0.1 g for each individual. Sample sizes of the species examined in this study ranged from 8.79 ± 0.25 (*Caranx hippos*) to 31.48 ± 4.93 cm (*Sphyraena barracuda*) in total length and 15.45 ± 0.40 (*Elops lacerta*) to 156.00 ± 39.30 g (*Pelmatolapia mariae*) in weight. The entire length-weight data in all the three stations were pooled together and the calculated correlation coefficient showed a high positive correlation between length and weight of all the fish species except in *Caranx hippos* (0.18) with low positive correlation. The b value obtained ranged from 0.61 for *Caranx hippos* to 3.53 for *Pelmatolapia mariae*. The mean condition factor ranged from 0.41 ± 0.03 to 4.23 ± 0.49 . The results of the present study will provide an effective tool for further studies of population dynamics and stock assessment studies.

Comment [L2]: Its always expected that different species have different total and standard lengths

Comment [L3]: Show figure

Formatted: Font: Not Italic

Comment [L4]: Conclude by letting us know the implications of the values derived and compare with previous data if any

Keywords: Cast net, Length-weight, Condition factor, New Calabar River.

Introduction

All fishing gears are species and size selective particularly in multispecies fisheries. The area of operation of a gear, the inconstant behavior of the fish relative to the gear, and size of the fish determine the part of a stock that can be caught by a gear. A generally important technical measure for fishing gears is the size selectivity which is defined as the probability of fish being retained in a fishing gear as a function of the length of the fish (Misund et al., 2001). In fisheries management, it is often desired that commercial fishing gear be highly selective for larger fish to minimize impact of fishing on the fish population and maximized yield (Gulland, 1983; Maclellan 1992).

Comment [L5]: Provide reference

Cast net is a falling gear, conical in shape with lead sunken or weights attracted at regular intervals on the lead rope forming the circumference of the cone. The cheapness and transportability make cast nets one of the most common gears in inland water fisheries (Jawad, 2000). This type of fishing gear is usually used in shallow waters -and cast from the shore or from a boat to catch -fish by falling and closing in on them. Cast nets are selective for lower size ranges, and larger, faster-moving fish can escape the falling net but may become entangled in the process (Welcome, 2001).

Knowing selectivity of the gear is very important since it affect population parameters such as length-weight relationship, gender ratio, estimate of population size through marking trails and growth and death ratios (Hamley, 1975). These relationships provide additional information about condition of fish in its habitat and also are vital in the biology of fisheries, assessing the fish's average weight in a given length using mathematical equations (Oscor *et al.* 2005). The parameters like general well-being of any fish species either in its natural habitat or cultivable environment, comparison of growth pattern, onset of maturity spawning, fecundity etc., can be assessed with the help of length—weight relation and condition factor (LeCren, 1951). Condition factor is important in understanding the life cycle of fish species and it contributes to adequate management of these species, hence, maintaining the equilibrium in the ecosystem (Imam *et al.*, 2010). The study was designed to provide basic scientific information on the length-weight relationship of some fish species in the New Calabar River, Niger Delta Nigeria.

Materials and Methods

The Study Area

The study area is the section of the New Calabar River as shown in Figure 1. The New Calabar River lies between longitude 006°53' 53.086"E and latitude 04°53' 19.020"N in Choba, Rivers

State, Nigeria. The entire river course is situated between longitude 7°60'E and latitude 5°45'N in the coastal area of the Niger Delta and empties into the Atlantic Ocean.

Three sampling stations (S1 – Aluu, S2 – Choba, and S3 – Iwofe) were established along the main course of the river. Fish species were collected monthly for 4 consecutive months (March to August, 2017) from the three sampling stations with the assistance of local artisanal fishers using different cast nets (1.5 and 2.5cm mesh sizes).

Comment [L6]: Any reasons in establishing these stations?




Figure 1: Map showing the sampling Stations

Formatted: Font: Not Italic

Fish Sampling

The fish were sampled on a monthly basis between the months of March to August 2017, from all with the assistance of local artisanal fishermen using cast nets of varying mesh sizes (1.5cm net area = 16.6 m² and 2.5cm). The local canoes used by the fishers were manned by an

Comment [L7]: Why not a year sampling?

Formatted: Font: Times New Roman

average of two men per boat. The specimens were immediately preserved in iced-packed cooler and transferred to the Fisheries Laboratory, Faculty of Agriculture, University of Port Harcourt, Choba where the identification were done, it was preserved in formalin in the laboratory, and immediately after, appropriate labelling and identification was made with the aid of relevant texts, Adesulu and Sydenham, (2007). Catch composition of cast nets were recorded by physical examination of the total catch, the Total Length (TL) and Standard Length (SL) were measured in centimeter (cm), and the Body Weights (BW) were measured in grams (g). The Total Length (TL) of each fish was taken from the tip of the snout (mouth closed) to the extended tip of the caudal fin using a meter rule.

Comment [L8]: Was this the determined as the catch-per-unit of effort?

Comment [L9]: Using which identification tool?

Comment [L10]: State the model of scale used and the error margin/deviation

The length–weight relationship is expressed by the equation $W = aL^b$, where W = body weight (g), and L = total length (cm), (Ricker, 1973). Parameters a and b were estimated by the logarithmic expression: $\log W = \log a + b \log L$ (Froese, 2006).

Formatted: Superscript

The condition factor which shows the degree of wellbeing of the fish in their habitat was determined by using the equation, $K = 100W/L^b$ (Gomiero and Braga, 2005). Where by K = condition factor W = the weight of the fish in gram (g) L = the total length of the fish in centimeters (cm) b = the value obtained from the length-weight equation.

Formatted: Superscript

Statistical evaluations of the variations observed in the different species were assessed using the SPSS (1999).

Comment [L11]: State the type of analysis

Results

A total of 1541 specimens of 26 fish species and representing 11 families. Sample sizes of the species examined in this study ranged from 8.79 ± 0.25 (*Caranx hippos*) to 31.48 ± 4.93 cm (*Sphyraena barracuda*) in total length and 15.45 ± 0.40 (*Elops lacerta*) to 156.00 ± 39.30 g (*Pelmatolapia mariae*) in weight (Table 1).

Comment [L12]: What are the dominant species?

Results of the LWR regressions are shown in Tables 2, 3 and 4. In station 1, apart from *Sardinella maderensis* with isometry growth pattern the remaining fish species showed negative allometry. The exponent b ranged from 0.33 (*Sphyraena barracuda*) to 3.35 (*Sardinella maderensis*). The coefficients of determination (r^2) of the LWR regressions ranged between 0.45 and 1.00. The mean condition factor ranged from 0.36 ± 0.03 recorded for *Sphyraena barracuda* to 3.80 ± 0.95 recorded for *Caranx hippos*.

Comment [L13]: Define all abbreviations to be used

Formatted: Font: Italic

Comment [L14]: Perfect correlation?

As far as the length weight relationship of station 2 was concerned, the growth pattern revealed allometry with the value of parameter b ranged from 0.99 for *Caranx hippos* to 4.23 for *Pelmatolapia mariae*. The coefficients of determination (r^2) of the LWR regressions ranged 0.32 to 0.98. The mean condition factor ranged between 0.95 ± 0.10 (*Mugil cephalus*) and 5.53 ± 0.85 (*Caranx hippos*).

Formatted: Font: Italic

In station 3, only *Coptodon zillii* revealed isometry growth pattern the remaining fish species showed allometry pattern. The exponent b ranged from 0.51 (*Caranx hippos*) to 3.12 (*Sardinella maderensis*). The coefficients of determination (r^2) of the LWR regressions ranged between 0.15 and 1.00. The mean condition factor ranged from 0.45 ± 0.05 to 3.08 ± 0.33 .

Comment [L15]: ???

The entire length-weight data in all the three stations were pooled together and the calculated correlation coefficient showed a high positive correlation between length and weight of all the fish species except in *Caranx hippos* (0.18) with low positive correlation. The b value obtained in this study ranged from 0.61 for *Caranx hippos* to 3.53 for *Pelmatolapia mariae*. The condition factor has been calculated for each species, the mean condition factor ranged from 0.41 ± 0.03 to 4.23 ± 0.49 .

Formatted: Font: Not Italic

Comment [L16]: Condition factor differs from species to species. It will be better to calculate condition factor for same species

UNDER PEER REVIEW

Discussion

In this study most of the samples consisted mainly juvenile with the sizes of fish species ranging from 8.79 ± 0.25 to 31.48 ± 4.93 cm in length and 15.45 ± 0.40 to 156.00 ± 39.30 g in weight. This can be ascribed to selectivity of the cast nets used in the study area and anthropogenic impacts, especially the fishing pressure and habitat destruction. The second most common indicator of unsustainable fishing is the observation of a decrease of large-sized fish, or a decrease in the mean size of the fish in the catch (e.g. Worm et al., 2009).

Length–weight relationships in fishes can be affected by multiple of factors including the number and length range of the sampled specimens (often affected by the type of fishing gear used), seasonality, habitat, gonad ripeness, sex, diet, stomach fullness, and growth phase (Froese, 2006; Karachle and Stergiou, 2008; Mir et al., 2013); however, these factors were not considered in the present study.

Hile (1936) and Martin (1949) opined that the value of 'b' may range between 2.5 and 4.0. In the literature, b values outside of this range are generally considered to be erroneous (Ricker 1975 Oscoz, 2005). LeCren (1951) pointed out that the variation in 'b' value is due to environmental factors, season, food availability, sex, life stage and other physiological factors. The b value obtained in this study ranged from 0.61 to 3.53 revealed that the studied species did not followed the cube law as all the species studied had allometric growth pattern.

According to LeCren (1951) and George et al. (1985) the relative condition factor (Kn) is an indicator of general well-being of the fish. (Kn) greater than one (1) is indicative of the general well-being of fish, whereas its value less than one (1) indicates that fish is not in a good condition. It was noticed that fish species in station 2 had highest condition factor values (0.95-5.53). This could be due to a difference in environmental conditions such as salinity.

The present work revealed that the pooled mean condition values factor ranged from 0.41 ± 0.03 to 4.23 ± 0.49 with only *Sphyraena barracuda* had less than one. This implies that the fish species are in good condition. However, the variations in the condition factor (K) observed in different fish species may be attributed to different factors, such as environmental condition, food availability and the gonadal maturity, as suggested by many workers (Le Cren, 1951).

Comment [L17]: At least the duration of the study could have been factored as you even claim

Comment [L18]: For which species?

Comment [L19]: Which one?

Comment [L20]: Any reason for this?

Conclusion

The study has provided baseline information to understand the length-weight relation and condition factor of different fish species caught using cast net in the New Calabar River. The study revealed that the catches are made up of relatively small sizes and allometric growth pattern in all the studied fish species. The condition factor indicated that almost all the species were thriving very well in the river. It is hoped that the results of the present study will provide an effective tool for further studies of population dynamics and stock assessment studies.

Comment [L21]: Suggesting species are what?? (growth or recruitment overfishing)

Table 1: Sizes Range of Fish Species Caught With Cast net

SPECIES	TOTAL LENGTH (Cm) Mean±SE	Range (Cm)	TOTAL WEIGHT (g) Mean±SE	Range (g)
CICHLIDAE				
<i>Coptodon guineensis</i>	16.35±0.25	8.7 - 33.7	121.80±6.73	13 - 697
<i>Coptodon zillii</i>	15.69±0.23	9.3 - 30.1	101.96±5.88	16 - 645
<i>Coptodon dageti</i>	16.16±0.52	9.5 - 26.3	96.15±9.04	20 - 311
<i>Sarotherodon galilaeus</i>	13.56±0.54	7.8 - 22.8	59.96±6.81	12 - 259
<i>Sarotherodon melanothron</i>	16.19±0.38	8.3 - 23.5	86.06±5.29	15 - 213
<i>Pelmatolapia mariae</i>	18.25±1.67	13.2 - 22	156.00±39.30	41 - 242
<i>Pelvicachromis taeniatus</i>	15.07±0.36	14.1 - 16.5	57.33±3.23	49 - 70
<i>Chromidotilapia guntheri</i>	14.25±0.39	13.2 - 15	55.75±2.78	48 - 61
<i>Hemichromis fasciatus</i>	14.33±0.45	13 - 14.9	57.25±3.57	47 - 63
MUGILIDAE				
<i>Liza falcipinnis</i>	20.96±0.41	9.1 - 37.1	97.39±5.86	13 - 370
<i>Liza grandisquamis</i>	9.80±0.70	9.1 - 10.5	13.50±1.50	12 - 15
<i>Mugil cephalus</i>	19.27±0.68	14.7 - 27.1	73.88±7.78	39 - 185
CLUPEIDAE				
<i>Ethmalosa fimbriata</i>	15.76±0.26	13.6 - 17.3	57.06±2.02	39 - 69
<i>Sardinella maderensis</i>	11.04±0.23	9.2 - 13.5	30.26±1.99	12 - 49
ALESTIDAE				
<i>Brycinus macrolepidotus</i>	15.23±0.67	9.6 - 22.5	51.78±4.65	16 - 94
<i>Brycinus nurse</i>	17.04±0.49	12.9 - 23.3	65.57±3.18	34 - 103
CLAROTEIDAE				
<i>Chrysichthys aluuensis</i>	13.45±1.47	9.9 - 22.4	39.50±4.32	27 - 69
<i>Chrysichthys nigrodigitatus</i>	15.11±1.62	9.8 - 22.3	50.90±16.84	16 - 195
LUTJANIDAE				
<i>Lutjanus agennes</i>	16.36±0.56	11.7 - 20.8	65.14±5.41	25 - 123
<i>Lutjanus dentatus</i>	16.24±0.42	12.8 - 20.6	64.09±4.25	32 - 120
CARANGIDAE				
<i>Caranx hippos</i>	8.79±0.25	7.3 - 10.5	25.72±1.04	19 - 33
<i>Trachinotus teraia</i>	10.83±0.92	7.5 - 14.1	27.10±2.27	19 - 38
ELOPIDEA				
<i>Elops lacerta</i>	12.76±0.30	7.6 - 15.7	15.45±0.40	9 - 22
HAEMULIDAE				
<i>Pomadasys jubelini</i>	10.89±0.30	8.8 - 14.3	21.78±2.43	11 - 69
MONODACTYLIDAE				
<i>Monodactylus sebae</i>	9.63±0.16	8.8 - 10.7	31.09±1.52	23 - 45
SPHYRAENIDAE				
<i>Sphyræna barracuda</i>	31.48±4.93	19.3 - 45	155.83±52.01	30 - 278

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Table 2: Growth Pattern and Conditional Factor for Station 1

Species	Condition factor (K)		A	b	r ²	Growth pattern
	Mean±SE	Range				
<i>Brycinus macrolepidotus</i>	1.44±0.05	0.83 - 1.94	-2.12	2.21	0.95	Negative allometry
<i>Brycinus nurse</i>	1.35±0.06	0.80 - 1.67	-0.72	1.72	0.92	Negative allometry
<i>Caranx hippos</i>	3.80±0.95	2.25 - 7.52	4.70	0.69	0.45	Negative allometry
<i>Chromidotilapia guntheri</i>	1.93±0.06	1.81 - 2.09	-0.95	1.87	0.99	Negative allometry
<i>Chrysichthys nigrodigitatus</i>	0.86±0.00	0.86 - 0.86	1.85	1.05	1.00	Negative allometry
<i>Coptodon dageti</i>	2.12±0.04	1.11 - 2.46	-3.33	2.81	0.94	Negative allometry
<i>Coptodon guineensis</i>	2.24±0.03	0.20 - 4.61	-3.15	2.76	0.91	Negative allometry
<i>Coptodon zilli</i>	2.22±0.02	1.48 - 2.98	-3.68	2.95	0.97	Negative allometry
<i>Elops lacerta</i>	0.86±0.08	0.49 - 2.05	0.28	0.96	0.85	Negative allometry
<i>Ethmalosa fimbriata</i>	1.46±0.03	1.29 - 1.62	-1.83	2.13	0.90	Negative allometry
<i>Hemichromis fasciatus</i>	1.95±0.07	1.84 - 2.14	-1.33	2.02	0.99	Negative allometry
<i>Liza falcipinnis</i>	0.88±0.02	0.44 - 1.73	-4.07	2.78	0.95	Negative allometry
<i>Lutjanus dentatus</i>	1.36±0.06	1.03 - 2.00	-2.21	2.24	0.84	Negative allometry
<i>Monodactylus sebae</i>	3.41±0.07	3.20 - 3.83	-2.61	2.66	0.94	Negative allometry
<i>Mugil cephalus</i>	1.08±0.09	0.75 - 1.24	-1.81	2.05	0.82	Negative allometry
<i>Pelmatolapia mariae</i>	2.33±0.03	2.27 - 2.37	1.87	1.17	0.96	Negative allometry
<i>Pomadasys jubelini</i>	1.64±0.00	1.64 - 1.64	1.84	0.44	1.00	Negative allometry
<i>Sardinella maderensis</i>	2.13±0.06	1.54 - 2.59	-4.70	3.35	0.88	Positive allometry
<i>Sarotherodon galilaeus</i>	2.16±0.24	1.06 - 3.49	-1.70	2.22	0.88	Negative allometry
<i>Sphyræna barracuda</i>	0.36±0.03	0.31 - 0.42	4.38	0.33	0.95	Negative allometry
<i>Trachinotus teraia</i>	2.63±0.44	1.32 - 4.50	1.02	0.96	0.96	Negative allometry

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Table 3: Growth Pattern and Conditional Factor for Station 2

Species	Condition factor (K)		A	b	r ²	Growth pattern
	Mean±SE	Range				
<i>Brycinus nurse</i>	1.37±0.07	0.83 - 1.58	-1.30	1.92	0.94	Negative allometry
<i>Caranx hippos</i>	5.53±0.85	2.17 - 7.45	5.38	0.99	0.40	Negative allometry
<i>Chrysichthys aluuensis</i>	2.15±0.31	0.60 - 3.11	1.23	0.94	0.84	Negative allometry
<i>Coptodon dageti</i>	1.17±0.04	1.08 - 1.24	0.16	1.32	0.98	Negative allometry
<i>Coptodon guineensis</i>	2.12±0.03	1.57 - 2.66	-3.61	2.91	0.98	Negative allometry
<i>Coptodon zillii</i>	2.18±0.02	1.49 - 2.54	-3.55	2.90	0.98	Negative allometry
<i>Liza falcipinnis</i>	1.00±0.08	0.83 - 1.20	-3.89	2.75	0.85	Negative allometry
<i>Monodactylus sebae</i>	3.37±0.09	3.20 - 3.48	-3.08	2.86	0.96	Negative allometry
<i>Mugil cephalus</i>	0.95±0.10	0.81 - 1.14	2.11	0.60	0.32	Negative allometry
<i>Pelmatolapia mariae</i>	2.02±0.26	1.63 - 2.51	-7.23	4.23	0.98	Positive allometry
<i>Sarotherodon galilaeus</i>	2.04±0.10	1.89 - 2.24	-1.47	2.08	0.84	Negative allometry
<i>Sarotherodon melanotheron</i>	2.38±0.26	1.87 - 2.72	0.89	1.26	1.00	Negative allometry

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Table 4: Growth Pattern and Conditional Factor for Station 3

Species	Condition factor (K)		a	b	r ²	Growth pattern
	Mean±SE	Range				
<i>Brycinus nurse</i>	1.34±0.12	0.79 - 1.74	-0.13	1.53	0.93	Negative allometry
<i>Caranx hippos</i>	3.08±0.33	2.17 - 4.33	2.05	0.51	0.15	Negative allometry
<i>Chrysichthys nigrodigitatus</i>	1.53±0.30	0.60 - 3.02	-1.42	1.90	0.68	Negative allometry
<i>Coptodon dageti</i>	2.45±0.32	1.15 - 4.55	-1.41	2.12	0.88	Negative allometry
<i>Coptodon guineensis</i>	2.21±0.03	1.15 - 4.61	-3.53	2.89	0.97	Negative allometry
<i>Coptodon zillii</i>	2.20±0.03	1.50 - 4.79	-3.77	2.98	0.97	Isometric
<i>Elops lacerta</i>	0.89±0.08	0.48 - 2.19	0.29	0.97	0.78	Negative allometry
<i>Liza falcipinnis</i>	1.08±0.05	0.72 - 2.70	-1.39	1.92	0.80	Negative allometry
<i>Liza grandisquamis</i>	1.45±0.15	1.30 - 1.59	0.38	0.78	1.00	Negative allometry
<i>Lutjanus agennes</i>	1.46±0.06	1.03 - 1.91	-2.36	2.32	0.84	Negative allometry
<i>Lutjanus dentatus</i>	1.66±0.08	1.33 - 2.22	-2.52	2.42	0.86	Negative allometry
<i>Monodactylus sebae</i>	3.47±0.06	2.94 - 3.69	-3.16	2.91	0.94	Negative allometry
<i>Mugil cephalus</i>	0.99±0.07	0.77 - 1.95	-2.19	2.18	0.79	Negative allometry
<i>Pelvicachromis taeniatus</i>	1.67±0.03	1.56 - 1.78	-2.08	2.26	0.96	Negative allometry
<i>Fantapenaeus notialis</i>	1.04±0.03	0.55 - 2.16	-1.04	1.66	0.79	Negative allometry
<i>Pomadasyus jubelini</i>	1.66±0.14	1.21 - 4.26	-2.26	2.21	0.53	Negative allometry
<i>Sardinella maderensis</i>	2.20±0.09	1.81 - 2.52	-4.12	3.12	0.89	Positive allometry
<i>Sarotherodon galilaeus</i>	2.04±0.07	1.05 - 4.64	-3.14	2.69	0.95	Negative allometry
<i>Sarotherodon melanotheron</i>	1.90±0.06	1.34 - 5.42	-2.90	2.61	0.92	Negative allometry

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

<i>Sphyraena barracuda</i>	0.45±0.05	0.39 - 0.55	-2.38	2.00	0.46	Negative allometry
----------------------------	-----------	-------------	-------	------	------	--------------------

Table 5: Pooled Growth Pattern and Conditional Factor

Species	Condition factor (K)		a	b	r ²	Growth pattern
	Mean±SE	Range				
<i>Brycinus macrolepidotus</i>	1.44±0.05	0.83±1.94	-2.12	2.21	0.95	Negative allometry
<i>Brycinus nurse</i>	1.35±0.04	0.79±1.74	-1.38	1.95	0.91	Negative allometry
<i>Caranx hippos</i>	4.23±0.49	2.17±7.52	4.55	0.61	0.18	Negative allometry
<i>Chromidotilapia guntheri</i>	1.93±0.06	1.81±2.09	-0.95	1.87	0.99	Negative allometry
<i>Chrysichthys aluuensis</i>	2.15±0.31	0.60±3.11	1.23	0.94	0.84	Negative allometry
<i>Chrysichthys nigrodigitatus</i>	1.47±0.28	0.60±3.02	-1.23	1.82	0.83	Negative allometry
<i>Coptodon dageti</i>	2.12±0.09	1.08±4.55	-2.43	2.47	0.83	Negative allometry
<i>Coptodon guineensis</i>	2.20±0.02	0.20±4.61	-3.42	2.85	0.95	Negative allometry
<i>Coptodon zilli</i>	2.20±0.02	1.48±4.79	-3.71	2.96	0.97	Negative allometry
<i>Elops lacerta</i>	0.87±0.06	0.48±2.19	0.28	0.96	0.81	Negative allometry
<i>Ethmalosa fimbriata</i>	1.46±0.03	1.29±1.62	-1.83	2.13	0.90	Negative allometry
<i>Hemichromis fasciatus</i>	1.95±0.07	1.84±2.14	-4.79	3.36	0.95	Positive allometry
<i>Liza falcipinnis</i>	0.98±0.03	0.44±2.70	-2.81	2.38	0.89	Negative allometry
<i>Liza grandisquamis</i>	1.45±0.15	1.30±1.59	0.38	0.78	1.00	Negative allometry
<i>Lutjanus agennes</i>	1.46±0.06	1.03±1.91	-2.36	2.32	0.84	Negative allometry
<i>Lutjanus dentatus</i>	1.47±0.05	1.03±2.22	-3.00	2.55	0.90	Negative allometry
<i>Monodactylus sebae</i>	3.43±0.04	2.94±3.83	-4.20	3.36	0.93	Positive allometry
<i>Mugil cephalus</i>	1.01±0.05	0.75±1.95	-2.34	2.22	0.81	Negative allometry
<i>Pelmatolapia mariae</i>	2.17±0.14	1.63±2.51	-5.33	3.53	0.98	Positive allometry
<i>Pelvicachromis taeniatus</i>	1.67±0.03	1.56±1.78	-2.08	2.26	0.96	Negative allometry
<i>Penaeus nitialis</i>	1.04±0.03	0.55±2.16	-1.08	1.68	0.80	Negative allometry
<i>Pomadasys jubelini</i>	1.66±0.13	1.21±4.26	-3.80	2.86	0.86	Negative allometry
<i>Sardinella maderensis</i>	2.16±0.05	1.54±2.59	-4.52	3.28	0.88	Positive allometry
<i>Sarotherodon galilaeus</i>	2.06±0.07	1.05±4.64	-3.07	2.68	0.94	Negative allometry
<i>Sarotherodon melanothron</i>	1.92±0.06	1.34±5.42	-2.87	2.60	0.93	Negative allometry
<i>Sphyraena barracuda</i>	0.41±0.03	0.31±0.55	-4.37	2.66	0.98	Negative allometry
<i>Trachinotus teraia</i>	2.63±0.44	1.32±4.50	1.02	0.96	0.96	Negative allometry

Formatted: Font: 12 pt


Formatted: Font: 12 pt, Not Italic

Formatted: Font: 12 pt


Formatted: Font: 12 pt, Not Italic

Formatted: Font: 12 pt


UNDER PEER REVIEW




Brycinus macrolepidotus




Brycinus nurse




Caranx hippos




Chromidotilapia guntheri




Chrysichthys aluuensis




Chrysichthys nigrodigatus




Coptodon dageti




Coptodon guineensis




Coptodon zilli




Elops lacerta




Ethimlosa fimbriata




Hemischromis fasciatus




Liza falcipinnis




Liza grandisquamis




Lutjanus agennes




Lutjanus dentatus




Monodactylus sebae




Mugil cephalus




Pelmatolapia mariae




Pelvicachromis taeniatus




Penaeus nitialis




Pomadasys jubelini




Sardinella maderensis




Sarotherodon galilaeus



Sarotherodon melanotheron



Sphyraena barracuda



Trachinotus teraia

Figure 2: Length-weight of fish species from New Calabar River

Formatted: Font: Not Italic

UNDER PEER REVIEW

References

- Adesulu EA and Sydenham DHJ. 2007. The fresh water and fisheries of Nigeria. *Macmillan Nigeria Publishers, Lagos*. 397 pp.
- Froese, R. 2006. Cube law, condition factor and Length-Weight relationships: history, meta-analysis and recommendations. *Appl. Ichthyol.*, 22, 241–253.
- George, J. P.; Sharma, A. K.; Venkateshvaran, K.; Sinha, P. S. R. K.; Venogopal, G. and Birader R. S. 1985. Length –weight relationship and relative condition factor in *Cirrhitina mirgala* and *Labeo rohita* from a sewage-fed tank. *The Annals of Zoology XXIII (2)*: 79-90
- Gulland, J.A. 1983. Fish stock assessment: A manual of basic methods- New York: Wiley.
- Gomiero, L. M., Braga, F. M. S. 2005. The condition factor of fishes from two river basins in Sao Paulo state, Southeast of Brazil. *Acta Scientiarum*. 27:73-78
- Hamley J.M. 1975 Review of Gillnet Selectivity. *Journal Fish Res., BD. Can* 32:1943-1969
- Hile, R. 1936. Age and growth of the Cisco, *Ambloplites rupestris* (Refinesequae) in Nebish Lake, Wisconsin. Trans. Wis. Acad. Sci. Arts. Lett. 33: 189-337
- Imam, T.S.; Bala, U.; Balarabe, M.L. & Oyeyi, T.I. 2010. Length-weight relationship and condition factor of four fish species from Wasai Reservoir in Kano, Nigeria. *African Journal of General Agriculture* 6(3):1595-6984.
- Jawad, L.A. 2006. fishing gear and methods of the Lower Mesopotamian Plain with reference to fishing management. *Marina Mesopotamica* 1:1:1-37
- Karachle, P. K.; Stergiou, K. I. 2008. Length-length and length weight relationships of several fish species from the North Aegean Sea (Greece). *J. Biol. Res. (Thessalon)* 10:149–157.
- Le Cren, E.D. 1951. The length–weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *J. Anim. Ecol.*, 20, 201–219
- MacLenna, D.N. 1992. Fishing gear selectivity. *Fish Res.*, 13:201-204.
- Martin, W. R. 1949. The mechanism of environmental control of body form in fishes. Univ. Toronto. Stu. Biol. 58: Ont. Fish. Res. Lab. 70: 1-91
- Mir, J. I.; Gusain, O.; Gusain, M. P.; Mir, F. A.; Sarkar, U. K.; Pandey, A. 2013. Length–weight and length–length relationships of the vulnerable dark mahseer, *Puntius chelunoides* (McClelland, 1839) from Garhwal Himalaya, India. *J. Appl. Ichthyol.* 30, 225–226.
- Misund, O.A., Kolding, J. and Fréon, P. 2002. Fish capture devices in industrial and artisanal fisheries and their influence on management. In: *Handbook of Fish Biology and Fisheries*, vol. II (eds. P.J.B.Hart and J.D. Reynolds). pp. 13-36. Blackwell Science, London

- Pillay, T. V. R. 1954. The biology of grey mullet, *Mugil tade* Forsskal with notes on its fishery in Bengal. *Proc. Nat. Inst. Sci. India*, 17: 187-217.
- Oscóz, J., Campos, F., Escala M.C. 2005. Weight-length relationships of some fish species of the Iberian Peninsula. *Journal of Applied Ichthyology*. 21:73-74.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. *Bulletin of the Fisheries Research Board of Canada* 191:1-382.
- Welcomme, R. L. 2001. Inland fisheries: ecology and management. *Fishing News Books*, Blackwell Science, Oxford, U.K., 358 pp.

UNDER PEER REVIEW