

## **Original Research Article**

### **THE EFFECT OF PARTIAL REPLACEMENT OF FISHMEAL WITH *Citrullus lanatus* AND *Moringa oleifera* SEED MEALS ON GROWTH PERFORMANCE OF *Clarias* *gariepinus* (Burchell, 1822) JUVENILES**

#### **ABSTRACT**

Four experimental diets were formulated to contain *Citrullus lanatus*, *Moringa oleifera* seed meal, *Citrullus lanatus* and *Moringa oleifera* seed meals as partial replacement for fishmeal (Treatment 2, 3 and 4 respectively) and Treatment 1 as control, which included fishmeal and soy bean meal as the protein sources were fed to one hundred and twenty (120) African catfish (*Clarias gariepinus*) of mean weight of  $25\pm 5.0$ g and mean length of  $14\pm 2.0$ cm in replicates for 10 weeks growth period. The results obtained showed that fish fed treatment 3 had highest mean weigh gain. More so, treatment 1 had the highest percentage weight gain and specific growth rate. Treatment 4 had the highest food conversion ratio and treatment 1 had the highest length increase. The result of the economic analysis of the four treatments showed the treatment 4 and 3 are the cheapest while treatment 1 is the most expensive. All the treatment diets showed that growth due to the different protein sources were not significant ( $p>0.05$ ). The result of this study therefore revealed that partial replacement of fishmeal with *Moringa oleifera* seed meal can be used in the diet of *Clarias gariepinus* juveniles.

**Keywords:** *Citrullus lanatus*, *Moringa oleifera*, *Clarias gariepinus*, diets.

#### **INTRODUCTION**

Aquaculture production is a major industry in many countries, and it will continue to grow as the demand for fisheries products increases and the supply from natural sources decreases. Nutrition plays a critical role in intensive aquaculture because it influences not only production costs but also fish growth, health and waste production [1]

Aquaculture production is becoming more and more intensive because production from capture fisheries has reached its maximum possible potential, as the catch is dwindling with each passing day [2]. [3] had stated that feed management determines the viability of aquaculture as it accounts for at least 60 percent of the cost of fish production [4], therefore the establishment of economically viable fish culture ventures requires the incorporation of agricultural wastes or by-products as feed ingredients or direct feed [5] to replace conventional feed stuffs whose dwindling supply has resulted into arbitrary hike in prices. Conventional ingredients used in fish feed are in high demand for human consumption and their yield are currently being affected by

climate change, hence out of concern for and the implications for food security as well as water and land use, there is urgent need to get local materials especially agricultural by-products of lower price to replace these costly feed materials. (Any references ??)

The importance of artificial feeding in aquaculture cannot be over-emphasized. It promotes faster growth, allows higher stocking density and shorter cultivation periods. However, the unavailability and affordability of adequate fish feed has significantly affected the development of aquaculture in Nigeria. (Any references ??) A well prepared and carefully formulated fish feed plays a significant role in fish culture [6]. The higher cost and competition imposed on some feed ingredients, such as soybean, groundnut cake, maize and sorghum used by human population as food, have necessitated the use of unconventional material for fish feed formulation. The price of finished feed in our part of the world continues to be on the rise, thereby removing the margin of profit accruing to the fish producers [7]. Thus, overcoming the burden of feed ingredients and reducing the cost of fish feed and fish products have been the burden of numerous researchers [8]. Fish farming, nutrition is critical because feed represents 40-50% of the production cost [9].

With the ever increasing cost of fishmeal in the tropics, [10] noted that several researchers have attempted to incorporate plant protein into formulated fish diet in order to reduce the quantity and cost of fishmeal without reducing the quality of the feed. According to [11] fishmeal contains some amino acids and fatty acids that are not present in tissues from terrestrial plants. The most limiting amino acids in plant protein are lysine and methionine [12, 13].

*Citrullus lanatus* seed meal is one of such agricultural by-product whose nutritive potential has not been effectively tapped in animal nutrition [14]. *Citrullus lanatus* seed is rich in minerals, protein, vitamins, carbohydrate and fibre [15, 16]. Watermelon seeds are rich in oil and protein [17, 18]. *Citrullus lanatus* seed oil proved to be good source of high quality edible oil characterized by low free fatty acid content [17]. The experience with *Citrullus lanatus* seed cake or meal in rations for animals showed that it is a good source of digestible protein comparable to other oil seed cakes like cottonseed, linseed etc [19] hence can be safely incorporated in animal feeds [20].

*Moringa oleifera*, which is one of the most widely cultivated species of the genus *Moringa* serves an example of nutritious vegetable tree with a variety of potential uses [21]. The immature green pods called “Drumstick” are probably the most valued and widely used part of the tree.

The seeds are sometimes removed from more mature pods and eaten like peas or roasted like nuts. The flowers are edible when cooked and are said to taste like mushrooms. The roots are shredded and used as condiment in the same way as horseradish.

In view of the increasing demand for fish and high cost of conventional feed ingredients, it is therefore necessary to investigate the effect of partial replacement of fish meal with *Citrullus lanatus* and *Moringa oleifera* seeds meals in the diets of *Clarias gariepinus* juveniles.

## **MATERIALS AND METHOD**

The juveniles of *Clarias gariepinus* for this research were obtained from Arazu's fish farm, Awka, Anambra State. (Where is the Anambra State?) The experiment which lasted for 70days was carried out also in 75 liters plastic tanks at Arazu's fish farm.

### **Procurement and Preparation of Seed Plants and Conventional Feed Ingredients for Feed Formulation**

Fishmeal, soybean, corn meal, wheat bran, vitamin and mineral premix, vegetable oil, salt and starch binder were procured from "Afor-Nnobi" market in Nnobi, Anambra state.

#### **Preparation of *Citrullus lanatus* seeds**

*Citrullus lanatus* seeds were gotten from large quantities of rotten or spoilt matured watermelon fruits, and were identified by a botanist. The seeds were soaked for twelve (12) hours to separate the seed nut from the fruit flesh [22]. The seed were sun dried, packaged in polythene bags and stored for processing. The dried seed were grounded into powder using a melon mill (Henry West GX 200). The milled watermelon seeds were then sieved with a hand sieved to separate the fine powder from the remaining fibre. The processed seeds were analyzed for proximate composition using the method described by the [23], to determine the crude protein, carbohydrate, fibre, fat, ash, dry matter, moisture content and Nitrogen free extract.

#### **Preparation of *Moringa oleifera* seed for feed formulation**

*Moringa oleifera* seeds were collected from Polycarp farm in Onitsha, Anambra state where it is commonly grown. The seeds were identified by a botanist. The seeds were shelled manually to obtain the cream coloured kernels. *Moringa* kernels were properly sorted out to remove stones and other dirt's. The clean kernels were grounded into powder using a melon mill (Henry West GX 200). The milled *Moringa* kernel seeds were also analyzed for proximate composition using

the method described by the [23], to determine the crude protein, carbohydrate, fibre, fat, ash, dry matter, moisture content and Nitrogen free extract.

### Formulation of Experimental Diets

The Experimental diets were formulated at Ezelagbo farms, Nise, Anambra State. Four diets were formulated. The percentage composition for each of the four diets is shown in Table 1. The experimental diets with crude protein levels of 40% were formulated using the Pearson square method. The individual ingredients were weighed using an electronic weighing balance (Citizen 5000MP) and then thoroughly mixed together in a bowl by hand. Water was added to the premixed ingredients and homogenized to a dough-like paste. The diets were then pelletised using a 3-mm electronic pelletizer (Model KL120B). The diets were sun-dried and stored in airtight plastic containers throughout the experimental period [2, 24].

**Table 1: Percentge Composition of the Experimental Diets**

Ingredients	Treatment 1 (Control)	Treatment 2 (F+C)	Treatment 3 (F+M)	Treatment 4 (F+C+M)
Fishmeal	40	30	30	20
Soybean	15	15	15	15
<i>Citrullus lanatus</i>	0	10	0	10
<i>Moringa oleifera</i> seed meal	0	0	10	10
Corn meal	20	20	20	20
Wheat flour	10	10	10	10
Wheat bran	10	10	10	10
Vit Min Premix	2	2	2	2
Vegetable oil	2	2	2	2
Salt	1	1	1	1
Total (%)	100	100	100	100

\*Keys: % (percentage), F (Fishmeal), C (*Citrullus lanatus*) and M (*Moringa oleifera* seed meal).

### Experimental Design

The experimental design used, was Completely Randomized Design (CRD). In this experiment, a total of 120 juvenile fish were used. After the acclimatization period which lasted for one week, fish were weighed using an electronic weighing balance (Citizen 5000MP) and homogenous sizes were randomly distributed in twelve (12) experimental plastic tanks with each tank containing ten juveniles of *Clarias gariepinus*. The fish were randomly assigned to four

treatment diets of T<sub>1</sub> – Fishmeal (control), T<sub>2</sub> - Fishmeal and *Citrullus lanatus* seed meal, T<sub>3</sub> - Fishmeal and *Moringa oleifera* seed meal and T<sub>4</sub> - Fishmeal, *Citrullus lanatus* and *Moringa oleifera* seed meal. Each treatment had three replicates. The experimental fish were fed twice daily (8.00AM and 4.00PM) at 5% body weight for ten weeks. The experimental tanks were thoroughly washed and water changed regularly at interval of three days throughout the experimental period. Water samples from the experimental tanks were collected and some physico-chemical parameters of the water analysed. These include; temperature, pH and dissolved oxygen (DO) using water sample kit (mercury in glass thermometer, pHep pocket-sized digital pH meter and Eutech Dissolve oxygen meter (model MW600) respectively). The juveniles were weighed every week to determine weight gain, and feed quantity given was adjusted accordingly.

#### **The Proximate Analysis of the ingredients and diets**

The feed used were analyzed for proximate composition at SMO Laboratory consult, Ibadan. Nigeria, using the methods of the Association of Official Analytical Chemists (AOAC, 2000) for crude protein (CP), moisture content, crude fibre, fat, ash and Carbohydrate.

#### **Indices of Growth and Feed Utilization Measured**

A Citizen 5000 MP electronic weighing balance was used to measure weights of juveniles per week till the end of the experiment (10 weeks), growth performance were estimated.

#### **Statistical Analysis**

The data obtained were subjected to Analysis of Variance (ANOVA) using Computer SPSS Version 22. Differences in means were evaluated at 5 percent (0.05) level of significance and direction of interaction between treatments were analyzed using Duncan HSD Post Hoc Test.

## **RESULTS**

**Table 2: Water Quality Parameters**

<b>Parameters</b>	<b>Mean values</b>
Temperature (°C)	30.18±0.21 <sup>a</sup>
Ph	6.03±0.13 <sup>a</sup>
Dissolve oxygen (mg/l)	4.42±0.10 <sup>a</sup>

*Mean values with the same superscript in the same column are not significantly different at (P>0.05).*

**Table 3: Proximate Composition of *Citrullus lanatus* and *Moringa oleifera* seeds used in formulation of experimental diets.**

Nutrients	<i>Citrullus lanatus</i> (%)	<i>Moringa oleifera</i> (%)
Crude protein	19.98	20.39
Crude fat	3.75	3.68
Crude fibre	3.89	3.82
Ash	6.63	6.65
Moisture	9.61	9.59
Dry matter	90.39	90.41
N.F.E	56.14	55.97

\*Key: N.F.E = Nitrogen Free Extract, % = percentage.

**Table 4: The Proximate compositions of the experimental diets of *Citrullus lanatus* and *Moringa oleifera* seeds in percentage (%) (Energy of diet groups should be given)**

Parameters	T <sub>1</sub> (Control)	T <sub>2</sub> (F+C)	T <sub>3</sub> (F+M)	T <sub>4</sub> (F+C+M)
Crude protein	20.18	19.89	20.26	20.58
Crude fat	3.59	3.65	3.61	3.57
Crude fibre	3.76	3.84	3.79	3.86
Ash	6.48	6.59	6.45	6.67
Moisture	9.46	9.55	9.41	9.63
Dry matter	90.54	90.45	90.59	90.37
N.F.E	56.53	56.48	56.47	55.69

\*Key: N.F.E = Nitrogen Free Extract, T<sub>1</sub> = (Fishmeal as control), T<sub>2</sub> = F+C (Fishmeal and *Citrullus lanatus* seed meal), T<sub>3</sub> = F+M (Fishmeal and *Moringa oleifera* seed meal) and T<sub>4</sub> = F+C+M (Fishmeal, *Citrullus lanatus* and *Moringa oleifera* seed meal).

**Table 5: Growth performance and feed utilization indices of *Clarias gariepinus* fed *Citrullus lanatus* and *Moringa oleifera* seed meal as partial replacement for fishmeal.**

**(Initial weights of treatments is much different. Experimental design is error. Please be controlled.)**

	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>
<b>MIW</b>	20.40±2.0 <sup>a</sup>	17.57±2.44 <sup>b</sup>	27.90±1.86 <sup>a</sup>	27.37±3.27 <sup>a</sup>
<b>MFW</b>	38.76±2.8 <sup>b</sup>	34.78±2.0 <sup>c</sup>	49.22±2.28 <sup>a</sup>	45.76±3.0 <sup>a</sup>
<b>MWG</b>	18.36±1.86 <sup>a</sup>	17.21±3.44 <sup>a</sup>	21.32±1.96 <sup>a</sup>	18.39±0.37 <sup>a</sup>
<b>PWG</b>	89.99±2.5 <sup>a</sup>	97.99±3.0 <sup>a</sup>	76.43±2.3 <sup>a</sup>	67.21±2.2 <sup>a</sup>

<b>SGR</b>	0.68±0.8 <sup>a</sup>	0.81±0.1 <sup>a</sup>	0.41±0.4 <sup>a</sup>	0.43±0.2 <sup>a</sup>
<b>FCR</b>	0.107±0.08 <sup>a</sup>	0.106±0.12 <sup>a</sup>	0.116±0.01 <sup>a</sup>	0.125±0.03 <sup>a</sup>
<b>MLI</b>	4.63±0.12 <sup>a</sup>	3.70±0.42 <sup>a</sup>	3.67±0.63 <sup>a</sup>	3.30±0.23 <sup>a</sup>
<b>SURVIVAL (%)</b>	100±0.00	100±0.00	100±0.00	100±0.00

Mean values with the same superscript in the same column are not significantly different at ( $P>0.05$ ).

**KEYS:**

**MIW:** Mean Initial Weight. **MFW:** Mean Final Weight. **MWG:** Mean Weight Gain.

**PWG:** Percentage Weight Gain. **SGR:** Specific Growth Rate. **FCR:** Food Conversion Ratio

**MLI:** Mean Length Increase. **%:** Percentage.

**RESULTS and DISCUSSION**

The results of the growth indices (Mean Weight Gain, Percentage Weight Gain, Specific Growth Rate, Food Conversion Ratio, and Mean Length Increase) of *Clarias gariepinus* juveniles fed on four dietary treatments showed that there was no significant difference ( $P>0.05$ ) in the fish growth.

The mean weight gain of the fish in the four treatments (T<sub>1</sub>-18.36±1.86, T<sub>2</sub>-17.21±3.34, T<sub>3</sub>-21.32±1.96 and T<sub>4</sub>-18.39±0.37) revealed that T<sub>3</sub> (21.32±1.96) had highest mean weigh gain though not significantly different ( $P>0.05$ ) from others. (Not necessary) The increased mean weight gain in T<sub>3</sub> (Fishmeal and *Moringa* seed) could be attributed to the high crude protein content of the feed type which is higher than T<sub>1</sub> (control diet) containing fishmeal. This agrees with the report of [25]. T<sub>2</sub> (Fishmeal and *Citrullus lanatus* seeds) had the lowest mean weight gain (17.21±3.34). [26], concluded that the inclusion level of 10% raw *Citrullus lanatus* seed meal in the diet of common carp is suggested to enhance growth performance and nutrient utilization. Although, [27] reported that *Citrullus lanatus* seed meal can be used as partial replacement for soybean in the practical diets of *Clarias gariepinus* up to 60%. He however suggested that the presence of anti-nutritional factors in plant ingredients could also be pointed at as a factor causing reduced growth rate and nutrient utilization by fish. Hence, the poor growth recorded in fish fed with *Citrullus lanatus* seed meal may be caused by the presence of these anti-nutritional factors [28]. (Which are the anti-nutritional factors in *Citrullus lanatus* seed meal?)

This result was in line with the findings of [29] who stated that growth is influenced through better food conversion and not just through stimulated feed intake. This indicates that fish which

consumed more food, utilizes it efficiently unlike the fish that consumed less food. Therefore, food consumption and not food conversion efficiency is a growth-limiting factor [30].

The mean Length Increase of the fish exposed to the four treatments ( $T_1$ - 4.63±0.12,  $T_2$ - 3.70±0.42,  $T_3$ - 3.67±0.63 and  $T_4$ - 3.30±0.23) revealed that  $T_2$  (3.70±0.42) had highest mean Length Increase while the least was recorded by those fed on  $T_4$  (3.30±0.23). The mean Length Increase was not significantly different ( $P>0.05$ ) among the four feed types. They all portray (another scientific word?) good length increase. (Are there any studies like for length increase ?)

The mean percentage mean weight gain of the fish in the four treatments ( $T_1$ -89.99±2.5,  $T_2$ - 97.99±3.0,  $T_3$ -76.43±2.3 and  $T_4$ -67.21±2.2) revealed that  $T_2$  (97.99±3.0) had highest mean percentage weigh gain while the least was recorded by those fed on  $T_4$  (67.21±2.2). The mean percentage weight gain was not significantly different ( $P>0.05$ ) among the four feed types. They all portray good weight increase. (Are there any similar studies ?)

The result of the specific growth rate (weekly) revealed that there was no significant difference ( $P>0.05$ ) in the specific growth rate.

The mean food conversion ratio of the fish exposed to the four treatments ( $T_1$ -0.107±0.8,  $T_2$ - 0.106±0.1,  $T_3$ -0.116±0.1 and  $T_4$ -0.125±0.03) revealed that  $T_4$ -0.125±0.03 had highest mean food conversion ratio while the least was recorded by those fed on  $T_2$  (0.106±0.1). The mean food conversion ratio was not significantly different ( $P>0.05$ ) among the four feed types. These showed they were able to convert their feed well into tissues and muscles which gave them good growth.

The results obtained from the water quality analysis showed that Temperature and Dissolve oxygen met the standard values for tropical fish culture as postulated by [31]. However, the mean value of pH was slightly lower than the standard value. No significant difference was recorded in the various parameters, suggesting that the fish can be reared in these water conditions comfortably.

The results obtained from the proximate analysis of *Citrullus lanatus* and *Moringa oleifera* seeds showed that the values of crude protein (CP) were 19.98% and 20.39% respectively. The crude protein value of *Citrullus lanatus* in this study was observed to be relatively lower when compared with the findings of [32] 24.4%, [33] 23.37%, [34] 21.46% and [35] 23.4% respectively. The differences could be in the environmental factors such as season, geographical



location and stage of maturity which play a minor role in determining nutritive value of *Citrullus lanatus* seeds meal [35]. The result obtained on crude protein value of *Citrullus lanatus* seed obtained from the study disagreed with the work of [36] who reported that *Citrullus lanatus* seeds, decorticated contained up to 34.5% crude protein content.

The result on crude protein value of *Moringa oleifera* seeds obtained from the study (20.39%) agrees with the findings of [37] who reported a range of 15.8 – 20.7% for protein content of Moringa (*Moringa oleifera*), pods. The value obtained was higher than the crude protein of raw *Moringa oleifera* seed flour (18.86%) and defatted *Moringa oleifera* seeds (17.13%) as reported by [38] and [39]. The value was however lower than raw *Moringa oleifera* seeds (35.97%), germinating and fermented *Moringa oleifera* seed flour (23.69% and 21.15%) as reported by [39]. The increased protein value of germinating and fermenting seed flour could be attributed to the biochemical activities of the germinating seeds and the activities of the microorganisms during fermentation [38]. [11] reported that Moringa seeds are good sources of fats, proteins, and crude fibers.

The results of the crude fibre contents in *Citrullus lanatus* and *Moringa oleifera* seeds conform to the range of 3-6% necessary for the growth of African catfish growth [40].

The results obtained from the proximate analysis of the four experimental diets types (T, T<sub>2</sub>, T<sub>3</sub> and T<sub>4</sub>) compared favourably with the findings of [25] who stated that usually fish growth will be directly proportional to the level of nutrient composition of its diet and if the level of crude protein ranges approximately within 20 to 40%. The results of the crude protein content of the four experimental diets were slightly lower than the recommendation of [41] who suggested that herbivorous and omnivorous fish require a diet with 25-35% crude protein. According to the report of [42], the nutrient requirement for growth, reproduction and normal physiological function of fish must be met, like other animals; but fish have much higher requirement for protein, so feed mixture with 25-45% of crude protein are mainly used.

The cost of a kilogram of the different feed types (T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub> and T<sub>4</sub>) in this study showed that feed type T<sub>4</sub> (Fishmeal, *Citrullus lanatus* and *Moringa oleifera* seed meal) was the cheapest (₦186.50), followed by T<sub>2</sub> (Fishmeal and *Citrullus lanatus* seed meal) (₦201.50), then T<sub>3</sub> (Fishmeal and *Moringa oleifera* seed meal) (₦221.50) and the costliest was feed type T<sub>1</sub> (Fishmeal as control) (₦226.50). (This prices per diets should be given as international value (for exp American dollars ) and Nigeria value. Otherwise, not mentioned in text. The difference in

cost of production of the four feed types was due to the different components/ingredients used in formulating the feeds.

## CONCLUSION

This research provides more insight in enhancing better growth performance of *Clarias gariepinus* juvenile fed with *Citrullus lanatus* and *Moringa oleifera* seeds as partial replacement for fishmeal. The results showed that their inclusion in the diet of *Clarias gariepinus* did not significantly increase the growth performance. However, partial replacement of fishmeal with *Moringa oleifera* seed meal gave a better growth performance and it's also cost effective when compared to fishmeal which is usually very expensive. Therefore, it can be included in the diet of *Clarias gariepinus* as this will boost the protein content and other nutrients present in the diet.

## REFERENCES

1. Gatlin, D.M. (2002). Nutrition and fish health. In: Fish Nutrition. J.E. Halver and R.W. Hardy (eds.), 3rd edition. London: *Academic Press*. pp. 671-702.
2. Gabriel U.U., Akinrotimi, O.A., Bekibele, D.O., Onunkwo, D.N., Anyanwu, P.E. (2007): Locally produced fish feed: potentials for Aquaculture development in subsaharan Africa. *African Journal of Agricultural Research*, **2**(7): 287-295.
3. Jamiu, D.M., Ayinla, O.A. (2003): Potential for the development of Aquaculture in African NAGA, *Worldfish Center Quarterly*, **26**(3): 9-13.
4. Akinrotimi O.A, Onunkwo D.N, Cliffe P.T, Anyanwu P.E, Orokotan O.O. (2007). The role of fish in nutrition and livelihoods of families in Niger Delta, Nigeria. *International Journal of Tropical Agriculture and Food Systems*. **1**(4): 344-351.
5. Shang, Y.C., Costa-Pierce, B.A. (1983): Integrated aquaculture-agriculture farming systems, some economic aspects. *Journal of World Mariculture Society*, **14**: 523-530.
6. Egbuonu, A.C.C. (2015). Comparative Investigation of the Proximate and Functional Properties of Watermelon (*Citrullus lanatus*) Rind and Seed. *Research Journal of Environmental Toxicology*. Volume **9** (3)160-167.
7. Amubode F.O. and Ogogo, A. O. (1994). Performance of snail *Archachatina marginata* fed levels of caloric protein supplementary diet. *Nigerian Journal of Forestry*, **24**: 34-43.
8. Ani, O. A. and Okeke, G. C. (2003). The substitution of pigeon pea (*Cajanus cajan*) seed meal for soybean in broiler finisher ration. *Proceeding 8<sup>th</sup> Annual conference of Animal Science Association Nigeria* Sept. 16-18<sup>th</sup> 2003 FUT. Minna: 10-12.

9. Steven, C. (2001). Understanding Fish Nutrition, Feeds and Feeding. 3<sup>rd</sup> ed. pp. 181-257. Academic Press, San Diego, CA, USA.
10. Halver, J. E. (2008). Protein and amino acids. Aquaculture development and coordination programme. Fish feed technology. F.A.O corporate document repository. Retrieved from <http://www.fao.org/docrep/x5738e/x573e04.htm> on 26/5/2016.
11. Foyd, N., Makkar, H. P. S., and Becker, K. (2001). The potential of *Moringa oleifera* for agricultural and industrial Uses. In J. Lowell and C. T. A. Fuglie (Eds.), The miracle tree: The multiple uses of (pp. 45–76). Wageningen: The Netherlands.
12. [Http://www.wikipedia.com](http://www.wikipedia.com) retrieved on June 16, 2016.
13. Eyo, A. A. (2003). Fundamentals of fish nutrition and diet development - An overview. National workshop. Fish feed development and feeding practices in Aquaculture. Organized by FISON/NIFER/FAO-NSPFS. Ed. Eyo A.A. 1-33pp.
14. Mohr, H.C. (1989): Water melon breeding. In: Bassett, M.J. (edition) Breeding Vegetable Crops. Avi Publication Company Incorporated, West Port, Connecticut USA., 37-66.
15. Duke, J.A., Ayensu, E.S. (1985): Medicinal plants of China. Reference publication vol. 1 p. 131.
16. Tarek, A.E., Khaled, M.T., (2001): Characteristic and composition of watermelon, pumpkin and paprika seed oils and flurs. *Journal of Agricultural Food Chemistry*, **49**(3): 1253-1259.
17. Mustafa, A.I., Badi, S.M. Salama, R.B., Elsayed, A.S., Hussain, A.A., (1972): Studies on watermelon seed oil. *Sudan Journal of Food Science Technology*, 4: 18-20.
18. Al-Khalifa, A.S. (1996): Physicochemical Characteristics, Fatty Acid Composition, an' Lipoxygenase Activity of Crude Pumpkin and Melon Seed Oils. *Journal of Agricultural and Food Chemistry*, 44: 964-966.
19. Kusel, J., (2001). Assessing well-being in forest dependent communities. *Journal of Sustainable Forestry*. **13** (12), 359–384.
20. Sastry, M.S., Singh, Y.P., Dutt, B. (1972). Studies on the toxicity of Bijada cake. *Indian Veterinary Journal*, 685.
21. Ramachandran, C.A, Peter, K.V, Gopalakrishnan, P.K. (1980). Drumstick (*Moringa oleifera*): a multipurpose Indian vegetable. *Economic Botany* **34** (3): 276-83.
22. Eyo, A. A. (2003). Fundamentals of fish nutrition and diet development - An overview. National workshop. Fish feed development and feeding practices in Aquaculture. Organized by FISON/NIFER/FAO-NSPFS. Ed. Eyo A.A. 1-33pp.
23. AOAC (Association of Official Analytical Chemists) (2000): Official methods of analysis AOAC international methods 934.01, 988.05, 920.39 and 942.05. Arlington, VA, USA: AOAC International.
24. Amisah, S., Oteng, M. A. Ofori, J. K. (2009). Growth performance of African Catfish, *Clarias gariepinus*, fed varying inclusions of *Leucaena leucocephala* leaf meal. *Journal of Applied Sciences and Environmental Management*, **13**(1): 21-26.
25. Erhirhie, E. O., and Ekene, N.E. (2013). Medicinal Values on *Citrullus lanatus* (Watermelon): Pharmacological Review. *International Journal of Research in Pharmaceutical and Biomedical Sciences*. Vol. 4 (4).
26. Lateef O. T., Victoria O. A., Victor T. O., and Saidu, U. (2014). Effect of Various Levels of Raw *Citrullus lanatus* Seed Meal Diets on Growth Performance of *Cyprinus carpio* Fingerlings. *Jordan Journal of Biological Sciences*. Volume 7, Number 4

27. Jimoh, W.A., Olawepo, K.D., Ayeloja, A.A., Ashraf, A.O and Shodamola, M.O. (2013). Evaluation of water melon seedmeal (*Citrullus lanatus*) as a replacer for Soybean seedmeal in the diet of African catfish (*Clarias gariepinus*). Proceedings of the 47th Annual Conference of the Agricultural Society of Nigeria Ibadan. Page 1228.
28. Sotolu, A.O., Faturoti, E.O. (2008): Digestibility and nutritional values of differently processed *Leucaena leucocephala* (Lam de Wit) seed meals in the diet of African Catfish (*Clarias gariepinus*). *Middle-East Journal of Scientific Research*, 3: 190-199.
29. Olivotto, I. K., Cardinali, M. M., Barbaresi, L. L, Maradonna, F. F. and Carnevali, O. O. (2003). Coral reef fish breeding: The secrets of each species. *Aquaculture*, **224**: 69 - 78.
30. Wang, N. N., Hayward, R. S., Noltie, D. B. (1998). Effect of feeding frequency on food consumption, growth, size variation, and feeding pattern of age-hybrid sunfish. *Aquaculture*, **165**: 261- 267.
31. Boyd, C.E and Gross, A (2000). Water use and conservation for inland aquaculture ponds. *Fisheries management and Ecology* **7** (1-2), 55-63.
32. Davies S.J. and Gouveia A. (2010). Response of common carp fry fed diets containing a pea seed meal (*Pisum sativum*) subjected to different thermal processing methods. *Aquaculture*, **305** (1-4):117-123
33. Fila, W.A., Itam, E.H., Johnson, J.T., Odey, M.O. Effiong, E.E., Dasofunjo, K., Ambo, E.E. (2013). Comparative Proximate Compositions of Watermelon *Citrullus lanatus*, Squash *Cucurbita Pepo*'l and Rambutan *Nephelium Lappaceum* *International Journal of Science and Technology*. Volume 2 No. 1.
34. Egbuonu, A.C.C. (2015). Comparative Investigation of the Proximate and Functional Properties of Watermelon (*Citrullus lanatus*) Rind and Seed. *Research Journal of Environmental Toxicology*. Volume **9** (3)160-167.
35. Erhirhie, E. O., and Ekene, N.E. (2013). Medicinal Values on *Citrullus lanatus* (Watermelon): Pharmacological Review. *International Journal of Research in Pharmaceutical and Biomedical Sciences*. Vol. 4 (4).
36. [Oyenuga, V. A., \(1968\). Nigeria's foods and foodstuffs. Ibadan, University Press](#)
37. Melesse, A.; Steingass, H. ; Boguhn, J. ; Schollenberger, M. ; Rodehutsord, M., (2012). Effects of elevation and season on nutrient composition of leaves and green pods of *Moringa stenopetala* and *Moringa oleifera*. *Agroforestry System.*, **86** (3): 505-518
38. Oluwole, S.I., Oluwole, A.A., and Oluwaseun, A. (2013). Comparative study on nutrient composition, phytochemical, and functional characteristics of raw, germinated, and fermented *Moringa oleifera* seed flour. *Food Science and Nutrition*. **1**(6): 452–463.
39. Peter, T. O., Philip, C. N. A. (2014). Proximate Analysis and Chemical Composition of Raw and Defatted *Moringa oleifera* Kernel. *Advances in Life Science and Technology*. Vol. 24.
40. Robinson *et al.*, (2001). A Practical Guide to Nutrition, Feeds, and Feeding of Catfish. *Mississippi Agricultural and Forestry Experiment Station*. Bulletin 1113, pp. 12.
41. Wilson R.P. and Halver J.E. (2005). Growth and Feed Utilization of *Oreochromis niloticus* fingerlings fed with diets containing soybean and cassava peelings: 3; 6-8
42. Davies S.J. and Gouveia A. (2010). Response of common carp fry fed diets containing a pea seed meal (*Pisum sativum*) subjected to different thermal processing methods. *Aquaculture*, **305** (1-4):117-123