

## Original Research Article

### Evaluation of Sharpshooter Lethal and Morphometric Indices Effects in *Clarias gariepinus*

#### ABSTRACT

##### Aim

In the present study, ~~the lethal and morphometrics indices effects were~~ evaluated ~~the lethal and morphometrics indices effects~~ in *Clarias gariepinus*.

##### Study Design

This study employs experimental design and statistical analysis of data and interpretation.

##### Place and duration of studies

This study was carried out in Applied Biology Special Laboratory Agbani, Enugu State University of Science and Technology Enugu State (ESUT), Enugu State Nigeria. It lasted thirty days.

##### Methodology

The effect of sharpshooter on the physicochemical parameters of the water used for the study was analysed using standard methods. The 96h LC<sub>50</sub> value estimated by Probit Analysis was 0.03 mg l<sup>-1</sup>. Based on the 96h LC<sub>50</sub>, the sublethal concentrations of sharpshooter (1/10<sup>th</sup> of 96h LC<sub>50</sub>, and 1/5<sup>th</sup> of 96 h LC<sub>50</sub> = 0.01 mg/L, 0.03 mg/L). The morphometric indices especially hepatosomatic index (HSI) and condition factor (K) were also estimated using standard methods.

##### Results

The physico-chemical parameters of the test water showed no significant difference ( $p > 0.05$ ) compared with the control. ~~Mortality caused by the pesticide increased with increase in concentration.~~ Mortality rate increased with increase in concentration with the highest recorded 0.05 mg l<sup>-1</sup> at 96 h (90% (27 fishes out of 30 fishes). The safe levels determined for the pesticide showed some variations. Whereas there was no significant difference ( $p > 0.05$ ) between 0.01 mg l<sup>-1</sup> treatments and control, 0.03 mg l<sup>-1</sup> caused a significant decrease ( $p < 0.05$ ) in HIS compared with control. Similarly, sharpshooter treatment caused a duration dependent significant increase ( $p < 0.05$ ) at day 15. The treatment with sharpshooter caused concentration and duration significant increase ( $p < 0.05$ ) in condition factor (K) compared with control.

##### Conclusion

This study has demonstrated that sharpshooter is toxic to *Clarias gariepinus* even at low concentrations. Therefore, the use of this pesticide in the environment especially farm lands and areas close to aquatic environment should be applied with caution to avoid the risk of contamination.

**Key words:** Sharpshooter; mortality; morphometric; *Clarias gariepinus*

Comment [FK1]: rewrite the sentence

## INTRODUCTION

The growing demand for increased food production to meet the need for the ever increasing global population has sophisticated agricultural technology in which pesticides especially insecticides play a crucial role. The aquatic living resources are very vulnerable to herbicides contamination as run-offs from farms and industries end up in water bodies [1]. Stability or variations in physicochemical parameters in water bodies' depends on human activities and the analyses of these parameters is useful for assessing the vulnerability of the water body and the organisms inhabiting there [2].

Sharpshooter is a broad spectrum pesticide consisting of both cypermethrin and profenofos in a formulation (profenofos 40 + cypermethrin 4 EC). *Profenofos* is a persistent and toxic organophosphorus insecticide widely used in agriculture for crop protection and pest control, thus marketed for these purpose [3-4]. Cypermethrin is a synthetic pyrethroid that has found wide acceptability. It is extensively used in agriculture and forestry because of its high activity against a broad spectrum of insect pest [5]. Nevertheless cypermethrin has been found to be highly toxic to fish [6] even in very low concentrations varying from 0.2 to 2.2 µg/L in 96hr [7]. Generally fishes exposed to toxicants have higher average concentration of bilirubin than ones not exposed [8]. [9] highlight the importance of evaluating growth response and oxidative stress in commercially important fish species.

African sharp tooth catfish *Clarias gariepinus* is a typical air-breathing catfish with scaleless bony elongated body with long dorsal and anal fins and a helmet like head. According to [10], it is probably the most widely distributed fish in Africa. They have an ubiquitous distribution in rivers, streams, ponds, dams, and lakes in Africa [11]. They are important commercial fish, widely consumed and cheap source of animal protein for low-income earners. The present study evaluated the effects of sharpshooter on the oxidative stress biomarker of *C. gariepinus*.

## MATERIALS AND METHODS

### Procurement of fish specimen and test chemical

A total of 90 juveniles of *C. gariepinus* were procured from Sacen Fish Farm, Enugu and transported in well aerated 500 litres capacity aquaria tanks to Applied Biology Special Laboratory Agbani, Enugu State University of Science and Technology Enugu State (ESUT), Enugu State Nigeria. The experimental fish were acclimatized for two weeks under laboratory condition, fed with top feed (a commercial feed) daily at 3 % body weight. Fecal matter and other waste materials were siphoned off and water was changed daily to reduce ammonia content in the water. Dead fishes were removed with forceps to avoid possible deterioration of the water quality. Ethical clearance was obtained from the Fishery Department, Ministry of Agriculture and Natural Resources committee on experimental animal care (MANR/FD/2017/EC101). Commercial formulation of profenofos 40 g and cypermethrin 4 g, with trade name "sharpshooter" supplied by West African cotton Ltd., Lagos Nigeria with CAS NO- 41198-08-7 and 52315-07-8 respectively were purchased in agrochemical shop in Ogbete Main Market Enugu.

### Determination of water quality parameter

Water quality parameters such as temperature, pH and dissolved oxygen were checked by direct reading methods using thermometer, pH meter and dissolved oxygen meter [12].

### Acute toxicity test

The test was conducted using a semi-static bioassay in 40 litres glass aquaria (60x30x30 cm). In the range finding test, the percentage mortalities of 0% and 100% lie between 0.01 mg/l and 0.05 mg/l. Therefore the definitive test was conducted consisted five concentrations of sharpshooter (0.01, 0.02, 0.03, 0.04, 0.05mg/l). During the exposures, each concentration ~~were-was~~ set in triplicate. Juveniles of *Clarias-C. gariepinus* were randomly exposed to different concentrations of sharpshooter. Another set of juvenile fish were simultaneously maintained in water without test chemical - control. Precaution was taken in the stocking of the fish by dropping them gently into the plastic aquaria. The experiment lasted for 96 hours (4 days). After 48 hrs of exposure, the test solution was changed so as to counter-balance the decreasing pesticide concentration.

The median lethal concentration (LC<sub>50</sub>) value was determined following the probit analysis method described by [13].

### Determination of safe levels

The Safe levels of the test pesticide were estimated by multiplying the 96 h LC<sub>50</sub> with different application factors (AF) and was based on [14-19].

### Determination of sublethal concentration

The 96 h LC<sub>50</sub> values of sharpshooter on *C. gariepinus* was 0.03 mg/l following the probit analysis method as described by [13]. Based on the 96h LC<sub>50</sub> value, the test concentration of sharpshooter was exposed to sublethal concentration (SL-1; 1/10<sup>th</sup> of 96h LC<sub>50</sub>, and 1/5<sup>th</sup> of 96h LC<sub>50</sub> = 0.01 mg/l, 0.03 mg/l). Ninety fishes were exposed to different sublethal concentrations and a control. Each treatment group ~~were-was~~ further randomized into three replicates of 10 fishes per replicate in 10 litres of water. The exposure lasted for 15 days during which the fish were fed with small quantity of food approximately 1% of total body weight about an hour before the test solution was renewed to avoid catabolism and subsequent mortality. On each sampling day (1, 5, 10 and 15), three fishes from each triplicate experiment including control were sacrificed.

### Determination of morphometric indices

The body weight and standard length of each fish were determined after each exposure interval. Thereafter, the liver dissected out, weighed so as to calculate the hepatosomatic index (HSI) and condition factor (K). The indices HSI and K were calculated according to according to [20].

$$HSI = \frac{\text{Liver weight}}{\text{body weight}} \times 100$$

$$K = \frac{\text{Body weight}}{\text{Total length (cm)}} \times 100$$

### Statistical analysis

The data obtained from the experiment were statistically analyzed using SPSS version 22. The data were subjected to two-way analysis of Variance (ANOVA) at significance difference of 5% probability level while Duncan multiple range test was used to determine the differences among treatment groups.

## RESULTS AND DISCUSSION

### Water quality parameters and percentage mortality rate of juveniles of *C. gariepinus* exposed to different concentrations of sharpshooter for 24, 48, 72 and 96 hours exposure period

The physico-chemical characteristics of the test water ~~is~~ are shown in Table 1. The pH of water varied within 8.45 - 8.60 in the treatments. The water temperature values ranged from 25.54°C – 25.81°C in the treatment. The dissolved oxygen varied within 5.0 - 5.5. Stability or variations in physicochemical parameters in water bodies' depends on human activities and the analyses of these parameters is useful for assessing the vulnerability of the water body and the organisms inhabiting there [2].

Table 2 showed the percentage mortality juveniles of *C. gariepinus* exposed to sharpshooter examined at different exposure periods (24, 48, 72 and 96 h) depending on different concentrations. The pesticide concentration of 0.05 mg/l at 96\_h exposure recorded highest mortality of 90% (27 fishes out of 30 fishes) while the least value at 0.01 mg/l at 96h exposure recorded the lowest mortality of 30% (9 fishes out of 30). The control recorded 0% mortality. Mortality rate increased with increase in concentration. The 96h LC<sub>50</sub> of 0.03 mg/l obtained for sharpshooter was lower than 0.38 and 1.25 mg/l reported for *O. niloticus* exposed to butachlor [21], respectively. Also, it is lower than 0.07 mg/l reported by [22] when *O. niloticus* was exposed to organophosphate commercial formulation pesticide. The 0.03 mg/l LC<sub>50</sub> in 96h obtained for sharpshooter in the present investigation indicates that the pesticide was very toxic to *C. gariepinus* juveniles. The toxicity of the pesticide was both exposure duration and concentration dependent, thus accounting for differences in LC<sub>10-90</sub> values obtained at different concentrations and durations of exposure.– The toxicity of compounds to organisms has however been known to be dependent on concentration, pH, temperature, developmental stages and exposure periods [23].

**Table 1. The Physico-Chemical Parameters of the Experimental Water Exposed to Different Concentration Levels of Sharpshooter**

S/N	Treatment (mg/L <sup>-1</sup> )	Temperature °C	DO (mg/l)	pH
1	Control	25.00±0.05	5.00±0.00	8.04±0.01
2	0.01	25.54±0.05	5.00±0.00	8.45±0.05
3	0.02	25.68±0.05	5.10±0.00	8.45±0.01
4	0.03	25.70±0.05	5.30±0.03	8.55±0.15
5	0.04	25.75±0.05	5.40±0.03	8.57±0.14
6	0.05	26.81±0.05	5.50±0.01	8.60±0.17

DO = Dissolved Oxygen

**Table 2. Percentage Mortality Rate of Juveniles of *Clarias gariepinus* Exposed to Different Concentrations of Sharpshooter for 24, 48, 72 and 96 Hours Exposure Period**

Conc. (µg/L)	Total Death	Survival / % mortality	Mortality			
			24 h	48 h	72 h	96 h
Control	0	100(0)	0	0	0	0
0.01	9	70(30)	0	3	3	3
0.02	11	63(37)	1	3	4	5
0.03	15	50(50)	2	3	5	5
0.04	20	33(67)	3	3	6	8
0.05	27	10(90)	3	6	9	9

Treatment size (n = 30)

#### Safe levels estimation

The safe levels were estimated following different methods (Table 3). In the present study, the safe levels determined for the pesticide showed some variations. However, due to large variation in safe levels as determined by different methods, the estimates of safe levels cannot be guaranteed. The estimated safe levels obtained for sharpshooter in *Clarias-C. gariepinus* in the present study, as calculated by multiplying the 96\_h LC<sub>50</sub> with application factor (AF) as recommended by different methods, varied from 3.00 x 10<sup>-3</sup> to 3.00 x 10<sup>-7</sup> mg/L. However, the large variation in safe levels determined by various methods has resulted in controversy

over its acceptability [24-25]. Dependence on LC<sub>50</sub> values could be a notable weakness in determining AF.

**Table 3. Estimated Safe Levels of Sharpshooter for *C. gariepinus* after 96 Hours**

Pesticides	96h LC <sub>50</sub> (mg/L <sup>+</sup> )	Method	Application Factor	Safe Level (mg/L <sup>+</sup> )
Sharpshooter	0.03	Hart et al (1948)	-	1.875 x 10 <sup>-03</sup>
		Sprague (1971)	0.1	3 x 10 <sup>-03</sup>
		CWQC (1973)	0.01	3 x 10 <sup>-04</sup>
		NAS/NAE (1973)	0.01 – 0.00001	3 x 10 <sup>-03</sup> – 3 x 10 <sup>-07</sup>
		CCREM (1991)	0.05	1.5 x 10 <sup>-03</sup>
		IJC (1977)	5% of 96h LC <sub>50</sub>	1.5 x 10 <sup>-03</sup>

Comment [FK2]: 03????

**Lethal concentration (LC<sub>50</sub>) of sharpshooter pesticide depending on exposure time for *C. gariepinus***

Table 4 showed the lethal concentration of sharpshooter in *C. gariepinus*. The LC<sub>50</sub> values with 95% confidence limits of different concentrations of sharpshooter were 0.05 (0.04-0.05), 0.04 (0.04-0.05), 0.03(0.03-0.04) and 0.03 (0.03-0.04) for 24\_h, 48\_h, 72\_h, and 9\_6h respectively. This showed that as the exposure time increases from 24h to 96h, the median lethal concentration decreases.

**Table 4. Lethal Concentration of Sharpshooter Depending on Exposure Time for Juvenile of *C. gariepinus***

Pesticides	Lethal concentration	Exposure Time (Hours)				
		24	48	72	96	
Sharpshooter	LC <sub>10</sub>	0.03 (0.02– 0.04) <sup>a</sup>	0.03 (0.02– 0.03) <sup>a</sup>	0.03 (0.02– 0.04) <sup>a</sup>	0.02 (0.01– 0.02) <sup>a</sup>	Formatted: Subscript
	LC <sub>20</sub>	0.04 (0.03– 0.04) <sup>a</sup>	0.03 (0.02– 0.04) <sup>a</sup>	0.04 (0.03– 0.04) <sup>a</sup>	0.02 (0.02– 0.03) <sup>b</sup>	Formatted: Subscript
	LC <sub>30</sub>	0.04 (0.03– 0.04) <sup>b</sup>	0.03 (0.03– 0.04) <sup>b</sup>	0.04 (0.03– 0.04) <sup>b</sup>	0.02 (0.02– 0.03) <sup>b</sup>	Formatted: Subscript
	LC <sub>40</sub>	0.04 (0.04– 0.05) <sup>b</sup>	0.04 (0.03– 0.04) <sup>a</sup>	0.04 (0.04– 0.05) <sup>b</sup>	0.03 (0.02– 0.03) <sup>a</sup>	Formatted: Subscript
	LC <sub>50</sub>	0.05 (0.04– 0.05) <sup>a</sup>	0.04 (0.04– 0.05) <sup>a</sup>	0.05 (0.04– 0.05) <sup>a</sup>	0.03 (0.03– 0.04) <sup>a</sup>	Formatted: Subscript
	LC <sub>670</sub>	0.05 (0.04– 0.06) <sup>a</sup>	0.05 (0.04– 0.06) <sup>a</sup>	0.05 (0.04– 0.06) <sup>b</sup>	0.04 (0.03– 0.05) <sup>a</sup>	Comment [FK3]: 670??? 60 or 70
	LC <sub>80</sub>	0.06 (0.05– 0.04) <sup>a</sup>	0.05 (0.05– 0.07) <sup>b</sup>	0.06 (0.05– 0.04) <sup>a</sup>	0.05 (0.04– 0.06) <sup>a</sup>	Formatted: Subscript
	LC <sub>90</sub>	0.07 (0.06– 0.12) <sup>a</sup>	0.06 (0.05– 0.07) <sup>b</sup>	0.07 (0.06– 0.12) <sup>a</sup>	0.06 (0.05– 0.08) <sup>a</sup>	Formatted: Subscript

#### Determination of morphometric indices

##### The hepatosomatic index (HSI) of juveniles of *Clarias-C. gariepinus* exposed to sharpshooter.

The hepatosomatic index (HSI) is shown at Figure 1. Whereas there was no significant difference ( $p > 0.05$ ) between 0.01  $\text{MgL}^{-1}$  treatments and control, 0.03  $\text{MgL}^{-1}$  caused a significant decrease ( $p < 0.05$ ) in HSI compared with control. Similarly, sharpshooter treatment caused a duration dependent significant increase ( $p < 0.05$ ) at day 15.

##### The condition factor (K) of juveniles of *Clarias-C. gariepinus* exposed to sharpshooter

The condition factor (K) in Figure 2. The treatment with sharpshooter caused concentration and duration significant increase ( $p < 0.05$ ) in condition factor (K) compared with control.

Morphometric indices serves as exposure index to environmental contaminants. The condition factor, a somatic biomarker is indicative of health and reflects feeding conditions as well as energy consumption and metabolism. Liver is the metabolic organ, it is a target for the metabolism in the fish body, the liver index (HSI) is a useful biomarker detect hazardous effects of the environmental stressors [26]. In sharpshooter, there were significant increase ( $p < 0.05$ ) in the HSI of the exposed fish as compared to the control. Increase in HSI have been reported in *Oreochromis-O. niloticus* exposed to paraquat herbicide® [27] and *Oreochromis O. mossambicus* exposed to azinphos-methyl® [28].

The Condition Factor (k) not only gives an indication of the fish health condition but can be used to elucidate the effects of contaminants in animals [29]. In the present study, fish exposed to all sharpshooter, showed significant decrease ( $p < 0.05$ ) indicating that the pesticide have effect on the condition factor of the exposed fish. Similar results have been reported in fish exposed to other toxicants [28]. This study thus, indicates that liver organ can be used as bio-indicator biomarker of pollutant effects of pesticide toxicity on fish and shows that fish are very sensitive to environmental changes.




Fig 1. Hepato-somatic indices (HSI) of *C. gariepinus* exposed to sharpshooter for 15-days. Letters indicated significant difference ( $p < 0.05$ ) in mean values among pesticide concentrations, and numerals indicated significant difference ( $p < 0.05$ ) in mean values among durations of exposure.






Fig 2. Condition Factor of *C. gariepinus* exposed to sharpshooter for 15-days. Letters indicated significant difference ( $p < 0.05$ ) in mean values among pesticide concentrations, and numerals indicated significant difference ( $p < 0.05$ ) in mean values among durations of exposure.

### Conclusion

From the research carried out on the juveniles of *C. gariepinus*, it was deduced that sharpshooter is toxic *C. gariepinus*. Also, short term exposure of juveniles of *C. gariepinus* to Sharpshooter at even low concentrations was sufficiently effective in disrupting physiological processes of *C. gariepinus*.

However, the use of this pesticide in the environment or near farm lands or in an area close to aquatic environment should be applied with caution to avoid the risk of pesticides contamination.

### Recommendations

We recommend that indiscriminate uses of insecticide should be monitored by government and non-~~Governmental~~ governmental organizations. Similar research to determine various effects of insecticide on fresh water and lakes should be carried out. Biological methods of controlling insects and pest should be adopted by farmers especially those around rivers and coastal regions.

### REFERENCES

1. Botelho R.G, Santos JB, Oliveira TA, Braga RR, Byrro ECM. Toxicidade aguda de herbicidas a tilapia (*Oreochromis niloticus*). Panta Daninha. 2009;27:621–626.
2. Nnamonu EI, Nkitnam EE, Ugwu FJ, Ejilibe OC, Ezenwosu SU, Ogbodo GU. Physicochemical Assessment of Vulnerability of the River Ebonyi in Eha-Amufu and Environs, Southeast Nigeria. Annual Research & Review in Biology. 2018;27(5):1-9.
3. Hassall KA. "The biochemistry and uses of insecticides". 2nd Ed., Macmillan press limited. Houndmills, Basingstoke, Taiwan. 1990;P. 81.
4. Chirions D, Geraud-povey F. "Effects de algunos insecticidas sobre rentomo fauna". Interciencia. 1996;21:31–36.
5. Casida JE, Gammon DW, Glickman AH, Lawrence LJ. Mechanism of Pyrethroid insecticides. Annual Review of Pharmacology and Toxicology. 1983;23:413–418.
6. US EPA. Pesticide fact sheet number 199. Cypermethrin. Office of pesticides and Toxic substances. Washington D.C. 1989;Pp 78.

Formatted: Font: Italic

7. David M, Mushigeni SB, Shivakumar R, Philip CH. Response of *Cyprinus ~~Carpio~~ carpio* (Linn) to sublethal concentration of cypermethrin. Alterations in Protein metabolic profiles. Chemosphere. 2004;56:347–352.
8. Nwamba HO, Acukanu C.E, Onyekwelu KC. (2006). Effect of crude oil and its product on bilirubin of African catfish *C. gariepinus*. Animal research international. 2006;3(3): 531–533.
9. Adeogun AO, Chukwuka AV, Ibor OR. Impact of abattoir and ~~Sawsaw~~-mill effluents on water quality of upper Ogun River (Abeokuta). American Journal of Environmental Science. 2011;7(6):525– 530.
10. Skelton P. A complete guide to the fresh water fishes of Southern Africa Shuk publishers, Cape town. 2001.
11. Association of Analytical Communities. Official methods of analysis. 16th ed. Association of Official Analytical Chemists. Arlington, Virginia, USA. 1995.
12. Adeyimi JA, Adewale OO, Oguma AY. Impact of pesticides on aquatic lives. Bulletin of Environmental Contamination and Toxicology. 2014;92(2):529-531.
13. Finney DK(ed). Probit analysis. Third edition, Cambridge: Cambridge University Press. 1971;pp.333.
14. Hart WB, Weston RF, Derman JC. An apparatus for oxygenating test solution which fish are used as test animals for evaluating toxicity. Transactions of American Fishery Society. 1948;75:288.
15. Sprague JB. Management of Pollutant Toxicity to Fish. Bioassay Methods for Acute Toxicity. Water Resources. 1971;3:793-821
16. Committee of Water Quality Criteria. A report of the Committee of Water Quality Research. Series, EPA-RS-73-003, US Environmental Protection Agency Report, CWQC: Cinlinnati, OLE. USA. 1972.
18. Canadian water quality guidelines, Canadian council of resources and Environmental Ministry, Inland Waters Directorate Environmenta Canada: Ottawa O.X. Canada. 1991.

19. International Joint Commission. New and Revised Great Lake Water Quality Objectives. Windsor: Ontario. 1977.
20. Khallaf EA, Authman M. A study of some reproduction character of *Bagrus Bayal*, Forskal, in Bahr Shabeen Canal. Journal of Egyptian German Society of Zoology, 1991;4:123– 38.
17. National Academy of Sciences/National Academy of engineering (NAS/NAE). Water Quality criteria. EPA-R3-033; US Government printing Office: Washington, DC, USA. 1973.
21. Nwani CD, Ivoke N, Ugwu DO, Atama CI, Onyeshii GC, Echi PC, Ogbonna SA. Investigation on acute toxicity and behavioural changes in a freshwater catfish, *Clarias gariepinus* (burchel 1822) expose to organophosphorous pesticides. Temifos. Paksitan Journal of Zoology. 2013;45(4):959–965.
22. Yaji AJ, Acute J, Onyiye SJ. (2011). Effect of cypermethrin on behaviour and biochemical indices of the freshwater fish *Oreochromis niloticus*. Electronic Journal of Environment, Agriculture and Food Chemistry. 2011;10:1927-1934.
23. Pandey AK, Nagpure NS, Trivedi SP, Kumar R, Kushwaha B, Lakra WS. Investigation on acute toxicity and behavioural change in chana punctatus (Bluch) due to organophosphate pesticide profenofos. Drug Chemical. Toxicology. 2011;34(4):424 – 428.
24. Buikerma JR, Naider-Lehner AL, Caims JR. Biological monitoring: part IV. Toxicology Testing. Environmental and Molecular Mutagenesis, 1982;33:239-262 .
25. Pandey S, Kumar R, Nagpure NS, Srivastava SK. Acute toxicity bioassays of mercuric chloride and malathion on air-breathing fish *Channa punctatus* (Bloch). *Ecotoxicology*. 2005;61:114-120.
26. Pait AS, Nelson JO. Vitellogenesis in male *Fundulus heteroclitus* (Killifish) induced by selected estrogenic compounds. *Aquatic Toxicology*. 2003;64:331-342.
27. Ada FB, Ekpenyong E, Bayim BPR. Heavy metal concentration in some fishes (*Chrysichthys nigrodigitatus*, *Clarias gariepinus* and *Oreochromis niloticus*) in the Great Kwa River, Cross River State, Nigeria. *Global Advanced Research Journal of Environmental Science and Toxicology*. 2012;1(7):183-189.
28. Jordaan MS, Reinecke SA, Reinecke AJ. Biomarkers responses and morphological effects in juvenile tilapia *Oreochromis mossanabicus* following subsequential exposure to the organophosphate to the organophosphate azinphosmethyl. *Aquatic Toxicology*. 2013;144

Formatted: Font: Italic

;145:138–140.

29. Kleinkauf A, Connor L, Swarbreck D (2004). General condition biomarkers in relation to contaminant burdens in European flounder (*Platichthys flesus*). *Ecotoxicology and Environmental Safety*. 2004;58:335–355.

Formatted: Font: Not Italic

UNDER PEER REVIEW