

**Original Research Article**  
**Assessment of total carbon stock in *Swietenia macrophylla* woodlot at Jhenaidah district in Bangladesh**

**ABSTRACT**

Woodlot plantations are supposed to have a massive outlook in carbon sequestration. Presence of large area woodlot plantations in Bangladesh would store a significant quantity of carbon. The purpose of this study was to determine the carbon stocks in woodlot plantations in Jhenaidah district, Bangladesh. Sixty sample plots of woodlot plantations were purposively selected from three upazilla (Kotchandpur, Kaligong and Moheshpur) of Jhenaidah district. The plot size was 10×10 m. Every individual tree present in the sampling plot was identified upto special level. Diameter at breast height (DBH) and total tree height were measured for all individual trees in each sample plot. Above ground biomass was calculated using the equation of Chave et al. (2005) and below ground biomass was estimated using the equation of Cairns et al. (1997) and converted into carbon using the equation of Chave et al. (2005). The estimated average tree density was  $1340 \pm 104.24$  stems  $\text{ha}^{-1}$  ranging between 800 and 2400 stems  $\text{ha}^{-1}$ . The estimated mean DBH and tree height were 19.52 cm and 12.57 m respectively. The total biomass in Mahogany (*Swietenia macrophylla*) woodlot plantations ranged between 52.48 and 824.44  $\text{Mg ha}^{-1}$  and the basal area in woodlot plantations ranged between 9.92  $\text{m}^2\text{ha}^{-1}$  and 86.21  $\text{m}^2\text{ha}^{-1}$ . The average total biomass was  $287.86 \pm 22.64$   $\text{Mg ha}^{-1}$  and average basal area was  $37.98 \pm 2.31$   $\text{m}^2\text{ha}^{-1}$ . In this study, the average carbon stock in Mahogany plantation was estimated to  $143.93 \pm 11.32$   $\text{Mg ha}^{-1}$  ranging between 26.24  $\text{Mg ha}^{-1}$  and 412.22  $\text{Mg ha}^{-1}$ . Three allometric models were developed and validated with equal strength ( $R^2$  0.97–0.98) using generalized linear regression. Woodlot plantations in Bangladesh can play a vital role in the UNFCCC's carbon mitigation and adaptation mechanism. So, the long-term sustainability of woodlot plantations must be addressed.

**Keywords:** Total carbon stock, *Swietenia macrophylla*, Woodlot, Climate change, Global warming etc.

**Comment [WBP1]:** Change the key words

**1. INTRODUCTION**

Global Warming, climate change and biodiversity conservation are the three concurrent issues in the present world. They are the consequence of the rise of greenhouse gases mainly  $\text{CO}_2$  [1, 2]. Human are mainly responsible of raising the amount of  $\text{CO}_2$  in the atmosphere through the combustion of fossil fuel and deforestation [3, 4]. According to UNFCCC (2007), the concentrations of atmospheric  $\text{CO}_2$  increased from a value of 278 parts per million to 379 parts per million in 2005, and the average global temperature rose by  $0.74^\circ\text{C}$ . The last 25 years this increased rate accelerated and recorded that 11 of the 12 months in a year were warmest [5]. Ecologists are worried about that the global warming will continue and the earth could warm by  $3^\circ\text{C}$  by 2100. If most of the countries try to reduce their greenhouse gas emissions, the Earth will continue to warm. It considers that the

average global temperature will go up to 4° C instead of 1.8° C by 2100 (UNFCCC, 2007). If this raising rate will continue the sea level up to 5m by melting the polar ice-cap [4] and as a result some South Asian countries will be the victim of sea level rising like Bangladesh.

Comment [WBP2]: Improve redaction

Bangladesh is one of the vulnerable countries of global warming in South-Asian. The total forest area in Bangladesh, according to Forest Department, is **estimated to be** 2.52 million ha **aproximately, corresponding** to 17.4% of **the its** land area **of the country**. This includes 1.52 Million ha **for** Forest Department controlled land, 0.73 million ha **of** Unclassified State Forests (USF) under the control of District Administration and 0.27 million ha **of** village forest land (mostly homesteads). In recent time, natural forests are decreasing very speedily due to anthropogenic activities such as housing, land conversion of other uses, deforestation, industrialization etc. The deforestation rate is faster than the previous time. But it is good that the amount of plantation forests is increasing during last two decades [6]. From 2005 to 2010, an average of 5 million ha of tree plantations have been established per year in Bangladesh. In 2010, the total amount of plantation forests is 264 million ha that covers up to 7% of the total forest area in the world (FAO, 2010). It assumed that, plantation forests provided about 35% supply of global wood in the year of 2000. It considers that this supply will go up to 44% by the year 2020 [3]. So, the importance of plantation forests is increasing undoubtedly.

Bangladesh, a developing country, is one of the lowest CO<sub>2</sub> emitting countries in the world. **Its is** estimated **that the** per capita CO<sub>2</sub> emission is 0.2 ton/year **in Bangladesh**, but the average for the developing countries is 1.6 ton/year. In developed countries, the per capita CO<sub>2</sub> emission is 20-50 times higher than the developing countries. In USA the per capita emission is 20 tons/year. Though Bangladesh is in low Green House Gas (GHG) emission category however provides no relief from the effects of Global Warming. If the sea level will rise in 1.5 meter, about 22,000 **sq.km<sup>2</sup> land of Bangladesh** will be inundated and 17 million people affected. Bangladesh will be the worst sufferer of global warming but **she** is not liable for this. Rather **she** is acting fundamental role in carbon sequestration. For minimizing the effects of global warming, Kyoto protocol, REDD+, carbon trading are well discussed issues **in the present world**. All the **treaties agreements** try to reduce the emission of CO<sub>2</sub> and enhancing the role of conservation, Sustainable Management of Forests and Enhancement of forest carbon stocks in Developing Countries [7].

Comment [WBP3]: Improve the redaction, can be the conuntry

Comment [WBP4]: What is that???? Improve the redaction of the whole paragraph

Bangladesh has already ratified the Kyoto protocol, REDD+ and involved in the carbon trading mechanisms. **In** carbon trading mechanisms, Bangladesh reduces the emission of carbon in the atmosphere within a certain limit and gets a certain amount of money in a certain period of time from the highest carbon emitting countries such as USA, Australia, and China etc. It encourages the use of green technology and helps in increasing forestation. It is an alternative way to recover the effects of global warming. Basically it is a reward to who are emitting less carbon dioxide and a charge on who are liable for global warming. It is not a day dream that one-day small holder farmers will get money in exchange of planting trees. So, it is very much necessary for Bangladesh to assess the biomass or carbon stock of the woodlot plantations for getting the carbon trading facilities. Considering the above background and justifications, this present study aims to assess carbon stocks of the woodlot plantations at Jhenaidah district in Bangladesh.

Comment [WBP5]: Set the numbers in this statement

## 2. MATERIAL AND METHODS

### 2.1 SITE DESCRIPTION

#### 2.1.1 GEOGRAPHIC AREA AND LOCATION

Jhenaidah is surrounded on the north by Kushtia and Rajbari districts, on the east by Magura district, on the south by Jessore district and on the west by Chuadanga district and India. The total area of the district is 1,964.77 km<sup>2</sup> (758.60 sq. miles). The district lies between 23°13' and 23°46' North latitudes and between 88°42' and 89°23' East longitudes.

Comment [WBP6]: Of ???

## 2.1.2 ANNUAL AVERAGE TEMPERATURE AND RAINFALL

About 90% of the total annual rainfall occurs during June- September and the annual rainfall is 1467 mm. The average daily temperature ranges from 12° C during December-January to about 31° C during May-August. The annual average temperature is 24° C, with the extreme lowest 8.1° C in January and the highest 40° C in May. Annual average maximum temperature is 37.1° C and minimum 11.2° C. Four main seasons namely the dry or winter season (December to February), the pre-monsoon hot season (March -May), the monsoon or rainy season (June-September) and the post monsoon or autumn season (October-November) are recognized.

Comment [WBP7]: Improve de the redaction, starting with temperature, the annual rainfall and so on.



Figure 1. Study Area (Jhenaidah district Map)

## 2.2 SAMPLE SIZE AND SAMPLING DESIGN

The most woodlot plantations of the study area are composed of *Switenia macrophylla* and the present study was conducted on woodlots of *Switenia macrophylla*. Purposive sampling was used for the data collection because of the uneven and discrete distribution of woodlots plantation in this region. During the reconnaissance survey, the study areas where woodlots plantation are dominant were selected through snow-ball methods. Twelve villages were selected from the study area (Kotchandpur, Maheshpur and Kaliganj Upazila). Five sample plots were selected from each village and total sixty sample plots from twelve villages were

Formatted: Font: Italic

Comment [WBP8]: It is redundant. Improve redaction and include the age of these woodlots

Comment [WBP9]: What kind of purposive sampling do you use?

selected. The size of each sampling plot was 10 m × 10 m (100 m<sup>2</sup>). The plantation plot whose individual's diameter is less than 8 cm was discarded purposively.

## 2.3 TOOLS

Diameter tape, measuring tape and Haga Altimeter were used for collecting primary data for the study area.

## 2.4 DATA COLLECTION PROCESS

### 2.4.1 PRIMARY DATA

Primary data were collected from the study area by using measuring tape, Spigel Relaskop and diameter tape for further analysis. The sample plot (10m\*10m) was alienated by using measuring tape. Diameters at breast height of individual species of the sample plot were measured by using diameter tape and heights of individuals were measured by using Haga Altimeter.

### 2.4.2 SECONDARY DATA

Secondary data and related information were collected from the following sources:

- Khulna university library
- Seminar library, Forestry and Wood Technology Discipline, Khulna University
- Published and Unpublished reports, journals, books.
- Newspapers.
- Regional center, BBS, Jhenaidah.
- Agriculture Information Services, Jhenaidah.
- Internet

## 2.5 DATA ANALYSIS

### 2.5.1 ALLOMETRIC COMPUTATIONS FOR ABOVEGROUND AND BELOWGROUND BIOMASS

Biomass equations relate DBH to biomass and biomass may differ among species as trees in a similar functional group can differ greatly in their growth form between geographic areas [8]. Considering these factors Chave *et al.* [9] developed allometric equation for tropical trees that was used for wide graphical and diameter range. By using Chave's allometric equation no (1), biomass of the Switenia macrophylla woodlots was estimated. The following Chave's universal allometric equation is:

$$AGB = \rho \times \exp(-1.499 + 2.148 \times \ln(DBH) + 0.207 \times (\ln(DBH))^2 - 0.0281(\ln(DBH))^3) \dots (1)$$

Herewhere,

AGB = Aboveground biomass

$\rho$  = Wood density (g cm<sup>-3</sup>), DBH = Diameter at breast height, ln = Natural logarithm

1.499 = Constant, 2.148 = Constant, 0.207 = Constant, 0.0281= Constant

**Comment [WBP10]:** Describe the use of this equation i.e. diameter range and how the variance inflation was controlled because it seems to be redundant?

Besides this Below Ground Biomass was calculated using the regression model suggested by Cairns *et al.* [10] as the most cost effective and practical method of determining root biomass.

$$\text{BGB} = \text{EXP} (-1.0587 + 0.8836 \times \text{LN}(\text{AGB})) \dots\dots\dots (2)$$

## 2.5.2 CONVERSION OF ABOVEGROUND BIOMASS TO CARBON

Estimated biomass was multiplied by the wood carbon content (50%). As almost all carbon measurement projects in the tropical forest assume that all tissues (i.e. wood, leaves and roots) consist of 50% carbon on a dry mass basis [9].

$$\begin{aligned} \text{Carbon (Mg)} &= \text{Biomass estimated by allometric equation} \times \text{Wood Carbon Content \%} \\ &= \text{Biomass estimated by allometric equation} \times 0.5 \end{aligned}$$

## 2.5.3 DENSITY AND BASAL AREA

Above ground carbon pools was computed using international standard common tree allometries combined with local tables of wood density by tree species. Statistical analysis was done in Microsoft Excel 2007 and SPSS-16 software. Basal area and density were measured to determine the correlation with the biomass of the woodlot.

The basal area  $\text{ha}^{-1}$  was calculated according to the formula (Equation no: 3)

$$\text{Ba/ha} = \frac{\epsilon\left(\frac{\pi}{4}\right)D^2}{\text{Sum of all quadrats area}} \dots\dots\dots (3)$$

Where,

Ba = Basal area in  $\text{m}^2 \text{ha}^{-1}$

D = Diameter at breast height in meter

$\pi = 3.14$

Following the formulas quantitative structure parameters of investigated trees were calculated

$$\text{Density (stem/ha)} = \frac{\text{Total no.of individuals of one species in all the plots}}{\text{Plot area} \times \text{Total no.of plots studied}} \dots\dots\dots (4)$$

$$\text{Basal area (m}^2/\text{ha)} = \frac{\text{Total basal area of individual species (m}^2\text{)}}{\text{Sample plot area (ha)} \times \text{Total no.of plots studied}} \dots\dots\dots (5)$$


## 3. RESULTS AND DISCUSSION

### 3.1 CARBON STOCKS OF THE WOODLOT

The total biomass (AGB and BGB) in [MahoganyMahogany](#) (*Switenia macrophylla*) woodlot plantations in Kotchandpur, Kaligong and Moheshpur upazilla were  $325.97 \text{ Mg ha}^{-1}$ ,  $199.63 \text{ Mg ha}^{-1}$  and  $337.99 \text{ Mg ha}^{-1}$  respectively (Table 1, 2 and 3, Fig. 2). In this study, the average carbon stocks in Mahogany (*Switenia Macrophylla*) plantations were estimated to  $162.98 \text{ Mg ha}^{-1}$ ,  $99.82 \text{ Mg ha}^{-1}$  and  $168.99 \text{ Mg ha}^{-1}$  in Kotchandpur, Kaligong and Moheshpur upazilla where the average density were  $1305 \text{ stem ha}^{-1}$ ,  $1390 \text{ stem ha}^{-1}$  and  $1325 \text{ stem ha}^{-1}$  respectively (Table no: 1, 2 and 3, Fig. 2). The average DBH were 19.52 cm, 16.09 cm and 19.56 cm; mean tree height was 12.57m, 11.62 m and 11.84 m; average basal area were  $42.01 \text{ m}^2\text{ha}^{-1}$ ,  $29.31 \text{ m}^2\text{ha}^{-1}$  and  $42.66 \text{ m}^2\text{ha}^{-1}$  in Kotchandpur, Kaligong and Moheshpur

upazilla respectively (Table 1, 2 and 3, Fig. 2). The biomass and carbon stock of Moheshpur ( $337.99 \text{ Mg ha}^{-1}$ ,  $168.99 \text{ Mg ha}^{-1}$ ) and Kotchandpur ( $325.97 \text{ Mg ha}^{-1}$ ,  $162.98 \text{ Mg ha}^{-1}$ ) upazilla was higher than the Kaligonj ( $199.63 \text{ Mg ha}^{-1}$ ,  $99.82 \text{ Mg ha}^{-1}$ ). Because most of the plantations in Kaligonj upazilla were new and the mean DBH was comparatively lower than the two upazilla. Stem density and tree height have lower effects on biomass and carbon stock in woodlot plantations. The stem density was higher in Kaligang upazilla ( $1390 \text{ stem ha}^{-1}$ ) then other two upazilla and mean tree height ( $11.62 \text{ m}$ ) was very close to Moheshpur ( $11.84 \text{ m}$ ) upazilla but biomass and carbon stock were lower than the two upazilla.

In Jhenaidah district, the estimated tree density was  $1340 \pm 104.24 \text{ stems ha}^{-1}$  ranging between 800 and 2400 stems  $\text{ha}^{-1}$  (Table 4). The estimated mean DBH and tree height were 19.52 cm and 12.57 m respectively (Table 4). The total biomass in Mahagony (*Swietenia macrophylla*) woodlot plantations ranged between 52.48 and 824.44  $\text{Mg ha}^{-1}$  and the basal area in woodlot plantations ranged between 9.92 and 86.21  $\text{m}^2 \text{ha}^{-1}$  (Table 4) in Jhenaidah district. The average total biomass was  $287.86 \pm 22.64 \text{ Mg ha}^{-1}$  and average basal area was  $37.98 \pm 2.31 \text{ m}^2 \text{ha}^{-1}$  (Table 4). In this study, the average carbon stock in Mahagony (*Swietenia Macrophylla*) plantation in Jhenaidah district was estimated  $143.93 \pm 11.32 \text{ Mg ha}^{-1}$  ranging between 26.24 and 412.22  $\text{Mg ha}^{-1}$  (Table 4) where 85% above ground and 15% below ground (Fig. 2).



**Figure 2. Carbon content of the woodlot plantations at three upazilla in Jhenaidah district**

### 3.2 BASAL AREA BASED ALLOMETRIC MODELS AND THEIR VALIDATION

Three types of models were developed for carbon assessment from the plot level mean basal area [Eqs. (1) to (3)]. We found a strong (mean  $R^2 = 0.97$ ; for Linear 0.97, Polynomial 0.97 and Power 0.98 models) and significant ( $P < 0.05$ ) relationship between mean biomass carbon and mean basal area for *Swietenia macrophylla* woodlot (Fig. 3). All three models were tested against 40 plots with Chave *et al.* [9] and Cairns *et al.* [10] for validation. Results of the GLRM revealed that all three models showed significant ( $P < 0.01$ ) and strong relationships ( $R^2 = 0.97$ ) with established models [9, 10] based on biomass carbon content (Fig. 3). We also tested our three models with Rahman *et al.* [11] and found that all three models showed significant ( $P < 0.01$ ) and strong relationships ( $R^2 = 0.99$ ) with three established models [11]. Given the high regression  $R^2$  (range: 0.97 to 0.98), our three models are equally strong in calculating the tree biomass carbon content. Therefore, our

**Comment [WBP11]:** Belowground?

**Comment [WBP12]:** Improve the redaction of this caption

**Comment [WBP13]:** If you have two or more variables do you have to report the adjusted R sq

basal area based allometric models are equally suitable for calculating biomass carbon content from the trees in woodlots.

$$\text{Biomass C} = 4.834 \times \text{BA} - 39.72 \quad (1)$$

$$\text{Biomass C} = -21.51 + 3.855 \times \text{BA} + 0.01 \times \text{BA}^2 \quad (2)$$

$$\text{Biomass C} = 1.256 \times \text{BA}^{1.291} \quad (3)$$




Figure 3. Basal area-based biomass models at the left side of this study validated with models established by Chave *et al.* [9] and Cairns *et al.* [10] at the right side.

### 3.3 RELATIONSHIP ASSESSMENT

In the plantation forest site, a significant positive correlation was observed among the following variables: DBH and total C, mean tree height and total C and basal area  $\text{m}^2\text{ha}^{-1}$  and total C. Two attributes basal area  $\text{ha}^{-1}$  ( $R^2=0.981$ ) (Fig. 6) and DBH ( $R^2=0.828$ ) (Fig. 4) were strongly related to aboveground carbon stocks, but stand mean height ( $R^2=0.572$ ) (Fig. 5) was weakly related because similar height of different plots having wide variation in number of trees and DBH.




Figure 4. Mean DBH and total carbon





Figure 5. Mean height and total carbon stock





**Figure 6. Mean Basal area and total carbon stock**

### 3.4 DISCUSSION

Managing the atmospheric carbon dioxide emission through minimizing deforestation, maximizing afforestation or reforestation, and checking global warming are the significant concern among scientists and policy makers [12, 13]. The United Nations Framework Convention on Climate Change (UNFCCC) is discussing policies and approaching to reduce CO<sub>2</sub> emissions from deforestation. Various mitigation and adaptation methods such as the Clean Development Mechanism (CDM), Reduced Emissions from Deforestation and Forest Degradation (REDD) and Reducing Emissions from Deforestation and Forest Degradation, and enhancing forest carbon stocks in developing countries (REDD+) are recognized by the UNFCCC. These are planned to connect multi-scale stakeholders in conservation and sustainable management of forest resources for mitigating global climate change [7].

Afforestation and reforestation are the fundamental parts of REDD+ as an effective mechanism for reducing global climate change [14, 15]. The accurate information on carbon stocks, biodiversity and the socioeconomic status of the communities in developing countries participating in the REDD+ financial mechanism is essential [13]. So, it is vital to obtain more truthful and precise biomass estimates for plantation forests or woodlots in order to develop about understanding of the role of plantation forests or woodlots in Bangladesh as well as in global carbon cycle.

The stem density (1340 tree ha<sup>-1</sup>) in our study was higher than recorded from other natural and protected forest in Bangladesh. For example, 381 trees ha<sup>-1</sup> in Chittagong Hill Tracts (South) Forest Division [16], 459 trees ha<sup>-1</sup> in Chunati Wildlife Sanctuary, Cox's Bazar [17], 464 trees ha<sup>-1</sup> in Dudpukuria-Dhopachori Wildlife Sanctuary of Chittagong South Forest Division [18], 257 tree ha<sup>-1</sup> in Ukhiya natural forests of Cox's Bazar Forest Division [19] and 369 stem ha<sup>-1</sup> in Ramu reserve forests of Cox's Bazar. The basal area (37.98 m<sup>2</sup> ha<sup>-1</sup>) in our study was higher than 16.88 m<sup>2</sup> ha<sup>-1</sup> in Chunati Wildlife Sanctuary, Cox's Bazar [17] and 27.07 m<sup>2</sup> ha<sup>-1</sup> in Dudpukuria-Dhopachori Wildlife Sanctuary of Chittagong South Forest Division [18]), but lower than 53.5 m<sup>2</sup> ha<sup>-1</sup> in Chittagong Hill Tracts (South) Forest Division [16]. This very high density of trees in woodlot plantations compared to many other natural and protected forest systems in Bangladesh may be because of maintaining a certain tree spacing (2m×2m) along with the higher survival rate from confirmed protection. Woodlot plantations in Bangladesh may therefore play an important role in producing more timber and revenue for the local community and more importantly sequester carbon compared to other natural and restored ecosystems in Bangladesh.

**Comment [WBP14]:** Discuss accordint to your results



Our established three basal area based allometric models can be useful for carbon calculation from the plot level mean basal area as all three models showed strong relationships in the GLRM analysis ( $R^2=0.98$ ). The mean biomass carbon of woodlot plantations ( $143.93 \text{ Mg ha}^{-1}$ ) in our study was also higher than the reported average tree biomass carbon content of  $83.72 \text{ Mg ha}^{-1}$  [14] and  $110.94\text{-ton ha}^{-1}$  [15] in hill forest of Bangladesh. Even our estimated carbon stock was higher than USA national average urban forest carbon storage ( $22.83 \text{ Mg ha}^{-1}$ ). Several measures of stand structure and productivity were also assessed to know the relationship with total carbon stock. Two attributes basal area  $\text{ha}^{-1}$  ( $R^2=0.981$ ) (Fig. 6) and DBH ( $R^2=0.828$ ) (Fig. 4) were strongly related to aboveground carbon stocks, but stand mean height ( $R^2=0.572$ ) (Figure 6) was weakly related because similar height of different plots having wide variation in number of trees and DBH. According to Kuyah *et al.* [20] DBH is very strongly ( $R^2=0.98$ ) related with aboveground biomass. Again Henry *et al.* [21] has shown tree volume is very strongly related with total aboveground biomass. The more basal area, stem density  $\text{ha}^{-1}$  and DBH indicate the more amount of biomass and more biomass means more carbon stock. But with similar height of different plots having wide variation in number of trees. For this reason, mean tree height is weakly related with aboveground carbon stock. By increasing woodlot plantation with long rotation with good diameter trees, we can increase the amount of total carbon stocks which can play a vital role in global carbon cycle. We estimated a much higher amount of carbon in woodlot plantations compared to other studies from tropical and subtropical regions due to the higher stem density and basal area (Table 4) in our study. This way woodlot plantation in Bangladesh can play an important role in atmospheric carbon sequestration.

#### 4. CONCLUSION

At present, natural forest is declining very hurriedly at shocking speed due to anthropogenic tricks (such as deforestation, industrialization, burning fossil fuels etc.) that boost global warming. According to FAO, the plantation forests and woodlots are increasing more than before. Bangladesh is probable to be one of the worst suffers of Global Warming due to extreme carbon emission and playing fundamental role in carbon sequestration especially through new plantation forest. Woodlot plantation is one of the major steps of plantation forest which has enormous potential in the lessening of global warming and adjustment to climate change. So, estimation of the forest carbon stocks is very essential which will facilitate us to appraise the quantity of carbon loss through deforestation or the amount of carbon that a forest can accumulate when such forests are regenerated. The woodlot plantations in Bangladesh could provide additional benefits to the local communities such as payment for environmental services for their participation in the UNFCCC's financial based carbon mitigation strategies. Further study on financial and biophysical aspect should be carried out in order to identify the effectiveness of woodlot plantation in the study area. One important limitation of this study is the application of universal biomass allometric equations to calculate total carbon stocks in woodlot plantation. Precision of these assessments could be improved by developing species specific local biomass allometric models. Unfortunately, such allometric equations are not available. Although there have been numerous studies carried out to estimate the forest biomass and the forest carbon stocks, but there is still a further need to develop a robust method to enumerate the estimates of biomass of all forest components, woodlots and carbon stocks more accurately.

**Comment [WBP15]:** Conclusions must be focused on the results of your research

## REFERENCES

1. Kumar BM. Species richness and aboveground carbon stock in the home gardens of central Kerala, India. *Agriculture, Ecosystems and Environment*. 2011;140:430-440.
2. Zhang Y, Duan B, Xian J, Korpelainen H, Li C. Links between plant diversity, carbon stocks and environmental factors along a successional gradient in a subalpine coniferous forest in Southwest China. *Forest Ecology and Management*. 2011;262:361-369.
3. Sharma MC, Baduni NP, Gairola S, Ghildiyal SK, Suyal S. Tree diversity and carbon stocks of some major forest types of Garhwal Himalaya, India. *Forest Ecology and Management*. 2010;260(12):2170-2179.
4. Detwiler RP, Hall CAS. Tropical forests and the global carbon cycle. *Science*. 1988;239:42-47.
5. IPCC, 2007. Fourth IPCC assessment report: climate change. Cambridge University Press, Cambridge, UK; 2007.
6. Community Risk-Assessment (CRA) and Risk-Reduction Action Plan (RRAP) Report; 2010.
7. Gardner TA, Burgess ND, Aguilar-Amuchastegui N, Barlow J, Berenguer E, Clements T, Danielsen F, Ferreira J, Foden W, Kapos V, Khan SM, Leesm AC, Parry L, Roman-Cuesta RM, Schmitt CB, Vieiram ICG. A framework for integrating biodiversity concerns into national REDD+ programmes. *Biological Conservation*. 2011;11:0-18.
8. Pearson T, Brown S, Birdsey RA. Measurement guideline for the sequestration of forest carbon. General Technical Report-NRS-18, USDA Forest Service, Northern Research Station, USA; 2007.
9. Chave J, Andalo C, Brown S, Cairns MA, Chambers JQ, Eamus D, Folster H, Fromard F, Higuchi N, Kira T, Lescure J P, Nelson BW, Ogawa H, Puig H, Yamakura T. Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Oecologia*. 2005;145:87-99.
10. Cairns MA, Brown S, Helmer EH, Baumgardner GA. Root biomass allocation in the world's upland forests. *Oecologia*. 1997;111(1):1-11.
11. Rahman MM, Kabir ME, Akon ASM. High carbon stocks in roadside plantations under participatory management in Bangladesh. *Global Ecology and Conservation*. 2015;3:412-423.
12. Kanowski P, McDermott C, Cashore B. Post Copenhagen strategies for the implementation of REDD+. In: Richardson K, Steffen W, Liverman D (Eds.), *Climate Change: Global Risks, Challenges and Decisions*. Cambridge University Press, NewYork USA. 2011;429-430.
13. Pandey SS, Cockfield G, Maraseni TN. Carbon stock dynamics in different vegetation dominated community forests under REDD+: a case from Nepal. *Forest Ecology Management*. 2014;327:40-47.
14. Bonan GB. Forests and climate change: forcings, feedbacks, and the climate benefits of forests. 2008;1444-1449.
15. Wang W, Lei X, Ma Z, Kneeshaw DD, Peng C. Positive relationship between aboveground carbon stocks and structural diversity in spruce dominated forest stands in New Brunswick, Canada *Forest Science*. 2011; 57(6):506-515.
16. Nath TK, Hossain MK, Alam MK. Diversity and composition of trees in Sitapahar forest reserve of Chittagong Hill Tracts (South) forest division Bangladesh. 1998;6 (1):1-9.

17. Rahman ML, Hossain MK. Status of fodder and non-fodder tree species in Chunati wildlife sanctuary of Chittagong forest division, Bangladesh. 2003;4(2):9-14.
18. Hossain MA, Hossain MK, Salam MA, Rahman S. Composition and diversity of tree species in duddpukuria-dhopachori wildlife sanctuary of Chittagong (South) forest division, Bangladesh. Res. J. Pharm. Biol. Chem. Sci. 2013;4(2):1447-1457.
19. Ahmed GU, Haque SMS. Percentage distribution of species and diameter class in natural forest of Bangladesh. 1993;17 (1):109-113.
20. Kuyah S, Dietz J, Muthuri C, Jamnadass R, Mwangi P, Coe R, Neufeldt H. Allometric equations for estimating biomass in agricultural landscape: Aboveground biomass. Agriculture, Ecosystem and Environment. 2012;158:216-224.
21. Henry M, Tiftonell P, Manlay RJ, Bernoux M, Albrecht A, Vanlauwe B. Biodiversity, carbon stocks and sequestration potential in aboveground biomass in smallholder farming systems of western Kenya. Agriculture, Ecosystem and Environment. 2008;129: 238-257.

## APPENDIX

**Table 1:** Average Height, Diameter at Breast Height, Basal Area, Stem Density, Biomass and Carbon stock in woodlot plantations at Kotchandpur Upazilla.

**Table 2:** Average Height, Diameter at Breast Height, Basal Area, Stem Density, Biomass and Carbon stock in woodlot plantations at Kaligong Upazilla.

**Table 3:** Average Height, Diameter at Breast Height, Basal Area, Stem Density, Biomass and Carbon stock in woodlot plantations at Moheshpur Upazilla.

Contents	Minimum	Maximum	Average	SD	SE
Mean Height (m)	6.54	15.11	12.57	2.24	0.50
Mean DBH (cm)	11.75	26.33	19.52	4.09	0.92
Basal Area (m <sup>2</sup> ha <sup>-1</sup> )	9.92	86.21	42.01	17.11	3.83
Stem density ha <sup>-1</sup>	900	2400	1305	466.19	104.24
Biomass (Mg ha <sup>-1</sup> )	52.48	747.79	325.97	157.24	35.16
Carbon stock (Mg ha <sup>-1</sup> )	26.24	373.89	162.98	78.62	17.58

Contents	Minimum	Maximum	Average	SD	SE
Mean Height (m)	6.77	15.48	11.84	1.99	0.44
Mean DBH (cm)	10.9	33.47	19.56	5.95	1.33
Basal Area (m <sup>2</sup> ha <sup>-1</sup> )	15.55	83.37	42.66	21.70	4.85
Stem density ha <sup>-1</sup>	800	2200	1325	355.22	79.43
Biomass (Mg ha <sup>-1</sup> )	80.84	824.44	337.99	220.85	49.38
Carbon stock (Mg ha <sup>-1</sup> )	40.42	412.22	168.99	110.43	24.69

**Table 4:** Average Height, Diameter at Breast Height, Basal Area, Stem Density, Biomass and Carbon stock in woodlot plantations in Jhenaidah District, Bangladesh.

Contents	Minimum	Maximum	Average	SD	SE
Mean Height (m)	6.54	15.55	12.01	2.25	0.29
Mean DBH (cm)	10.9	33.47	18.39	4.9	0.63
Basal Area (m <sup>2</sup> ha <sup>-1</sup> )	9.92	86.21	37.98	17.88	2.31
Stem density ha <sup>-1</sup>	800	2400	1340	378.31	48.84
Biomass (Mg ha <sup>-1</sup> )	52.48	824.44	287.86	175.35	22.64
Carbon stock (Mg ha <sup>-1</sup> )	26.24	412.22	143.93	87.67	11.32

Contents	Minimum	Maximum	Average	SD	SE
Mean Height (m)	7.46	15.55	11.62	2.51	0.56
Mean DBH (cm)	11.11	24.37	16.09	3.74	0.83
Basal Area (m <sup>2</sup> ha <sup>-1</sup> )	13.08	51.61	29.31	10.45	2.34
Stem density ha <sup>-1</sup>	800	2000	1390	311.03	69.55
Biomass (Mg ha <sup>-1</sup> )	68.52	450.86	199.63	98.05	21.92
Carbon stock (Mg ha <sup>-1</sup> )	34.26	225.43	99.82	49.02	10.96