

# Effect of transplanting method and gypsum rate on yield and yield contributing characters of boro rice in saline zone of Bangladesh

## Abstract

Salinity intrusion causes problems in the coastal areas of Bangladesh. Climate change creates hazards like cyclone, sea level rise, and storm surge have been increasing the salinity problem in many folds. The coastal region covers about 20% of the country, from where cultivable land more than 30%. Agricultural land uses in these areas are very poor, because of high content of salinity in Rabi season. Already, 830,000 million hectares of land already identified as affected by soil salinity. A field experiment was carried out at saline prone area at Binerpota, Satkhira under natural salinity condition during Rabi season 2017-2018. The experiment was carried out with two varieties namely, Binadhan-10 was evaluated under four transplanting methods i.e., M<sub>0</sub>: Control (No Slope/flat land), M<sub>1</sub>: Ridge and furrow (each furrow 30 cm wide accommodating 3 lines transplanting), M<sub>2</sub>: Ridge and furrow (each furrow 60 cm wide accommodating 5 lines transplanting) and four levels of gypsum with control G<sub>0</sub>: control, G<sub>1</sub>: 75 kg ha<sup>-1</sup>, G<sub>2</sub>: 150 kg ha<sup>-1</sup> as basal and G<sub>3</sub>: 150 kg ha<sup>-1</sup> (75 kg ha<sup>-1</sup> as basal +75 kg ha<sup>-1</sup> 42 DAT). The experiment was laid out in a split plot design with three replications. The unit plot size was 3 m x 4 m. The recommended fertilizer doses applied for the experiment were 80 kg N ha<sup>-1</sup>, 15 kg P ha<sup>-1</sup>, 50 kg K ha<sup>-1</sup>. Nitrogen, phosphorus, potassium, sulphur and zinc were supplied from urea, TSP, MoP, gypsum and zinc sulphate monohydrate respectively while urea was applied in three equal splits. Application of gypsum had significant effect on plant height, number of effective tiller m<sup>-2</sup>, length of panicle, total number of spikelets panicle<sup>-1</sup>, thousand grain weight, number of filled spikelets panicle<sup>-1</sup>, grain yield straw yield. It seems that the crop responded to the application of gypsum. Overall results suggest that an application of gypsum The highest grain yield (7.7 t ha<sup>-1</sup>) was produced in ridge and furrow method where gypsum rate was 150 kg ha<sup>-1</sup> (75 kg ha<sup>-1</sup> as basal + 75kg ha<sup>-1</sup> at weeks after transplanting followed by (7.4 t ha<sup>-1</sup>) basal application of 150 kg ha<sup>-1</sup> gypsum along with N, P, K, Zn and Boron might be necessary to ensure satisfactory yield of rice in saline prone area under natural salinity condition.

**Key words:** Salinity intrusion, zinc sulphate monohydrate

## Introduction

Bangladesh is a deltaic country with 14.4 million ha area [\(reference?\)](#). About 80% of the country's area consists of alluvial sediments deposited by the rivers Ganga, Brahmaputra, Tista, Jamuna, Meghna and their tributaries (Haque, 2006). However, a part of the cultivable area in coastal districts is affected with varying degree of soil salinity due to the intrusion of saline water during high tides. Soil salinity is also believed to be responsible for low cropping intensity in these districts (Rahman & Ahsan, 2001). The salinity affected area increased from about 0.83 million ha in 1973 to 1.02 million ha in 2000, and 1.05 million ha in 2009 (SRDI, 2010). Salinity is one of the major causes hindering agricultural productivity in the world. Globally nearly 7% of the world is afflicted by soil salinity. Salinity caused by anthropogenic factors (secondary [salinization](#) ~~Stalinization~~) is often related to large-scale development of irrigated agriculture without adequate drainage and clearing of natural deep rooted vegetation. Problems associated with the presence of excess salts in the soil have for long constrained agricultural productivity. More than 80 per cent of the total area of the Khulna, Bagerhat and Satkhira districts are already affected by different magnitudes of soil salinity of which about 35 per cent is in the grip of

46 strong salinity (Mainuddin *et al.* 2011). Soil degradation, which can be caused by salinity, is  
47 considered as an environmental impairment with severe adverse effects on agricultural  
48 productivity, particularly in arid and semiarid regions (Qadir et al., 2006). Effects of high level  
49 of soluble salts in soil mainly causes an increase in osmotic pressure that hindered to uptake  
50 water from soil. Soil salinity is considered the most critical environmental stress which can  
51 negatively affect rice growth and the metabolism process (Rodriguez-Navarro & Rubio, 2006).  
52 Several procedures and strategies that can be used to improve salt affected area. The chemical  
53 remediation is one of these reclamation strategies (Sharma & Minhas, 2005). The application of  
54  $\text{Ca}^{2+}$  amendments can improve different properties of soil and act as soil modifiers that can  
55 prevent development of sodicity which is directly related to plant growth, crop productivity and  
56 crop yields (Wong et al., 2009; Chintala et al., 2010). Specific chemical amendments gypsum  
57 ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ) can be used as direct source for  $\text{Ca}^{2+}$  cation; however gypsum is normally  
58 available. Gypsum plays a significant role in the reclamation of saline soils by providing a  $\text{Ca}^{2+}$   
59 cation to replace the exchangeable  $\text{Na}^+$  from the colloid's cation exchange positions and leaching  
60 it out from the root zone into groundwater (Sharma & Minhas, 2005). Fageria and Knupp (2014)  
61 reported that gypsum and lime application significantly improved growth and yield of rice. There  
62 are some agronomic management practices (ridge and furrow method) through which salinity  
63 level of a soil can be lowered and the stress effects can be mitigate. Application of gypsum and  
64 organic amendments and irrigation recast. Introduction of salinity tolerant rice varieties in  
65 combination with agronomic management practices (ridge and furrow method) for the  
66 amelioration of salinity stress effect is the key for improving crop productivity in coastal  
67 ecosystem of Bangladesh.

## 68 **Materials and methods**

### 69 **Experimental site**

70 The field experiment was conducted at saline prone area, Farmer's field Binarpota, Satkhira under  
71 natural salinity condition during 2017-2018. The experiment was carried out with Binadhan-10  
72 why Binadhan-10 not other varieties/hybrids?

73 (Is it rice variety...hybrid or??) was evaluated under four transplanting methods i.e.,  $M_0$ : Control  
74 (No Slope/flat land),  $M_1$ : Ridge and furrow (each furrow 30 cm wide accommodating 3 lines  
75 transplanting),  $M_2$ : Ridge and furrow (each furrow 60 cm wide accommodating 5 lines  
76 transplanting). four levels of gypsum with control  $G_0$ : control,  $G_1$ : 75 kg ha<sup>-1</sup>,  $G_2$ : 150 kg ha<sup>-1</sup> as  
77 basal and  $G_3$ : 150 kg ha<sup>-1</sup> (75 kg ha<sup>-1</sup> as basal +75 kg ha<sup>-1</sup> 42 DAT). The experiment was laid out  
78 in a split plot design with three replications.

### 79 **Seedlings raising**

80 Seedlings were raised in well prepared wet seed bed at the sub-station Satkhira farms. Before  
81 sowing, seeds were immersed in water for 24 hours and then they were taken out and kept in jute  
82 sacks in dark condition for 48 hours. Seedling nurseries for each variety were prepared by  
83 puddling the soil. The sprouted seeds were sown on a well prepared wet nursery bed in 1 January,  
84 2018. No manuring and fertilization was done but water and pest management practices were  
85 followed in order to raise healthy seedlings.

### 86 **Land preparation**

87 The land preparation was started one month prior to transplant of the seedlings. The land was  
88 thoroughly prepared with the help of a power tiller. Subsequently the land was sufficiently  
89 irrigated and ploughed and cross ploughed three times with country plough followed by  
90 laddering to have a good tilth. All kinds of stubble and residues of previous crop were removed  
91 from the field. After uniform leveling, the experimental plots were laid out according to the  
92 requirement of the treatment.

### 93 **Fertilization and manuring**

94 The plots of Boro rice were fertilized with N, P, K, Zn and Boron respectively according to the  
95 recommendation of BARC (2012). The whole amount of triple super phosphate, muriatic of  
96 potash, and zinc sulphate were applied to the soil at the time of final land preparation. Urea was  
97 applied in three equal splits. One split of urea was applied with other fertilizers as basal dose and  
98 the other two splits were applied 21 and 45 DAT. The seed bed was wet by application of water  
99 both in the morning and evening on the previous day before uprooting the seedling. Thirty days  
100 old seedlings were uprooted carefully from the seedling nursery for transplanting in the  
101 experimental plots. Only selected healthy seedlings were translated in the experimental plots in 1  
102 February 2018 in 20 cm apart line maintaining a distance of 15 cm from hill to hill with three  
103 seedlings hill<sup>-1</sup> proper care was taken during the growing period of the crop.

### 104 **Intercultural operation**

105 Intercultural operating were done in order to ensure and to maintain the normal growth of the  
106 plant as and when needed. After one week of transplanting dead seedling were replaced carefully  
107 by transplanting fresh seedlings from the same source. The experiment plots were infested with  
108 some common weeds which were removed twice by hand weeding. After transplanting six  
109 irrigation were needed to maintain 5-6 cm standing water in each plot. Finally the field was  
110 drained out 7 days before harvest. Observations were regularly made and the field looked nice  
111 with normal green plants.

### 112 **Harvesting and data collection**

113 The maturity of crops was determined when some 70% of the seeds became attain their  
114 characters color. Grain and straw yields plot were recorded after threshing by a pedal thresher  
115 winnowing and drying in the sun properly including the grains and straws of the sample plants.  
116 The weight of grains was adjusted to 12% moisture content. Grain and straw yield were them  
117 converted to t ha<sup>-1</sup>. From the 10 randomly harvested hills, the following data were recorded, plant  
118 height, number of total tillers hill<sup>-1</sup>, number of effective tillers hill<sup>-1</sup>, number of non-effective  
119 tillers hill<sup>-1</sup>, number of grain panicle<sup>-1</sup>, number of unfilled spikelet's panicle<sup>-1</sup>, 1000 grain weight,  
120 Grain yield (tha<sup>-1</sup>), Straw yield (t ha<sup>-1</sup>).

### 121 **Collection and preparation of soil samples**

122 The initial soil samples were collected from the plough depth level (0-15) cm. The samples were  
123 taken by means of an auger from different spots of the field and mixed thoroughly to make a  
124 composite sample. The composite sample was air dried ground and sieved through a 10-mesh  
125 (2mm) sieve and stored in a plastic bag for physical and chemical analysis. The initial soil  
126 sample was analyzed for physical and chemical properties in the Soil Science laboratory of  
127 BINA.

128

## 129 **Chemical analysis of soil sample**

130 Soil samples were analyzed for both physical and chemical characteristics. The soil samples  
131 were analyzed following the standard methods as follows.

132 Table 1. Chemical properties of the soil at the experimental field

| Chemical properties                                  |      |
|--|------|
| pH <sub>1:5</sub>                                    | 7.2  |
| EC <sub>1:5</sub> (dS m <sup>-1</sup> )              | 8.1  |
| Na <sup>+</sup> (meq L <sup>-1</sup> )               | 59   |
| K <sup>+</sup> (meq L <sup>-1</sup> )                | 0.29 |
| Ca <sup>2+</sup> (meq L <sup>-1</sup> )              | 6.0  |
| Mg <sup>2+</sup> (meq L <sup>-1</sup> )              | 10.3 |
| HCO <sub>3</sub> <sup>-</sup> (meq L <sup>-1</sup> ) | 7.1  |
| Cl <sup>-</sup> (meq L <sup>-1</sup> )               | 47.0 |
| SO <sub>4</sub> <sup>2-</sup> (meq L <sup>-1</sup> ) | 24.8 |
| SAR  | 18.5 |
| ESP (%)  | 31.2 |

133

## 134 **Data processing and analysis**

135 Data recorded for different parameters were subjected to analysis of variance (ANOVA) and the  
136 treatment means were compared using the least significant different test.

## 137 **Results and discussion**

138 Transplanting method had significant effect on most of the plant parameters. The highest grain  
139 yield (6.6 t ha<sup>-1</sup>) was produced in ridge and furrow (each furrow 30 cm wide accommodating 3  
140 lines transplanting) and lowest yield (5.8 t ha<sup>-1</sup>) in control (flat land) method. In case of gypsum  
141 rates, 150 kg ha<sup>-1</sup> (75 kg ha<sup>-1</sup> as basal + 75kg ha<sup>-1</sup> at 7 WAT (G<sub>3</sub>) gypsum produced the highest  
142 grain yield (7.3 t ha<sup>-1</sup>) followed by 150 kg ha<sup>-1</sup> gypsum application as basal (7.0 t ha<sup>-1</sup>). The plant  
143 height, number of total tillers hill<sup>-1</sup>, panicle length, number of filled grains panicle<sup>-1</sup> and thousand  
144 grain weight were higher in the 150 kg ha<sup>-1</sup> (75kg ha<sup>-1</sup> as basal +75kg ha<sup>-1</sup> at 7WAT (G<sub>3</sub>)  
145 application of gypsum than basal 150 kg ha<sup>-1</sup> application of gypsum. Interaction between  
146 transplanting method and gypsum application showed that the highest grain yield in ridge &  
147 furrow (each furrow 30 cm wide accommodating 3 lines transplanting) method with 150 kg ha<sup>-1</sup>  
148 gypsum (75kg ha<sup>-1</sup> as basal +75kg ha<sup>-1</sup> at 7WAT) (7.7 t ha<sup>-1</sup>) followed by ridge & furrow (each  
149 furrow 60 cm wide accommodating 5 lines transplanting) 150 kg ha<sup>-1</sup> gypsum application as  
150 basal (7.4 t ha<sup>-1</sup>) (Table 2).

151

152

153 Table 2. Effect of transplanting method, rates of gypsum on yield and yield contributing characters of rice


154

| Treatments                                    | Plant height (cm) | Total tillers hill <sup>-1</sup> (no) | Effective tillers hill <sup>-1</sup> (no) | Panicle length (cm) | Filled grains panicle <sup>-1</sup> (no.) | Unfilled grains Panicle <sup>-1</sup> (no.) | 1000 Seed wt. (g.) | Grain yield (t ha <sup>-1</sup> ) | Straw yield (t ha <sup>-1</sup> ) |
|---|-------------------|---------------------------------------|---|---------------------|---|---|--------------------|-----------------------------------|-----------------------------------|
| <b>Transplanting method</b>                   |                   |                                       |   |                     |   |   |                    |                                   |                                   |
| Control (Flat land) (M <sub>1</sub> )         | 102.3             | 12.8                                  | 12.0                                      | 26.0                | 132.8                                     | 5.3   | 24.2               | 5.8                               | 9.3                               |
| Ridge & Furrow 30cm (M <sub>2</sub> )         | 100.9             | 13.0                                  | 12.0                                      | 26.8                | 137.6                                     | 4.9   | 24.2               | 6.6                               | 9.8                               |
| Ridge & Furrow 60cm (M <sub>3</sub> )         | 101.3             | 12.6                                  | 11.9                                      | 25.9                | 134.9                                     | 6.1   | 24.4               | 6.4                               | 9.1                               |
| LSD <sub>0.05</sub>                           | NS                | NS                                    | NS  | 1.8                 | 17.0                                      | NS  | 1.4                | 0.4                               | 0.9                               |
| <b>Level of Gypsum</b>                        |                   |                                       |   |                     |   |   |                    |                                   |                                   |
| 0 kg ha <sup>-1</sup> (G <sub>0</sub> )       | 102.4             | 13.3                                  | 12.5                                      | 26.2                | 132.2                                     | 6.0   | 24.4               | 5.4                               | 8.5                               |
| 75kg ha <sup>-1</sup> (G <sub>1</sub> )       | 102.1             | 12.7                                  | 12.0                                      | 26.3                | 136.7                                     | 5.1   | 23.8               | 6.2                               | 9.2                               |
| 150 kg ha <sup>-1</sup> (G <sub>2</sub> )     | 100.7             | 12.4                                  | 11.4                                      | 26.2                | 138.8                                     | 5.3   | 24.5               | 7.0                               | 9.8                               |
| 150 kg ha <sup>-1</sup> (G <sub>3</sub> )     | 100.8             | 12.8                                  | 11.9                                      | 26.3                | 132.7                                     | 5.3   | 24.4               | 7.3                               | 10.2                              |
| LSD <sub>0.05</sub>                           | NS                | NS                                    | 1.0                                       | 0.9                 | 19.5                                      | NS  | 1.5                | 0.7                               | 1.0                               |
| <b>Transplanting Method × Rates of Gypsum</b> |                   |                                       |   |                     |   |   |                    |                                   |                                   |
| M <sub>1</sub> G <sub>0</sub>                 | 104.2             | 12.5                                  | 11.8                                      | 25.8                | 145.2                                     | 3.7   | 25.2               | 5.3                               | 8.0                               |
| M <sub>1</sub> G <sub>1</sub>                 | 102.2             | 11.9                                  | 11.4                                      | 25.9                | 132.1                                     | 5.4   | 24.9               | 6.3                               | 9.1                               |
| M <sub>1</sub> G <sub>2</sub>                 | 99.5              | 12.8                                  | 11.9                                      | 25.9                | 129.2                                     | 4.3   | 24.9               | 6.7                               | 9.1                               |
| M <sub>1</sub> G <sub>3</sub>                 | 103.5             | 13.9                                  | 13.0                                      | 26.6                | 124.8                                     | 7.9   | 24.9               | 6.8                               | 10.8                              |
| M <sub>2</sub> G <sub>0</sub>                 | 100.6             | 14.1                                  | 13.2                                      | 26.9                | 117.9                                     | 6.7   | 24.4               | 5.3                               | 8.3                               |
| M <sub>2</sub> G <sub>1</sub>                 | 102.1             | 12.2                                  | 11.3                                      | 27.8                | 138.1                                     | 3.9   | 24.3               | 6.0                               | 9.3                               |
| M <sub>2</sub> G <sub>2</sub>                 | 101.2             | 13.2                                  | 11.8                                      | 26.3                | 146.4                                     | 4.4   | 24.3               | 7.2                               | 10.8                              |
| M <sub>2</sub> G <sub>3</sub>                 | 99.8              | 12.5                                  | 11.6                                      | 26.1                | 148.1                                     | 4.4   | 24.2               | 7.7                               | 10.7                              |
| M <sub>3</sub> G <sub>0</sub>                 | 102.4             | 13.4                                  | 12.6                                      | 25.8                | 133.7                                     | 7.7   | 23.8               | 5.7                               | 9.0                               |
| M <sub>3</sub> G <sub>1</sub>                 | 102.0             | 13.9                                  | 13.2                                      | 25.4                | 139.9                                     | 5.9   | 23.6               | 6.3                               | 9.1                               |
| M <sub>3</sub> G <sub>2</sub>                 | 101.6             | 11.0                                  | 10.5                                      | 26.4                | 140.7                                     | 7.3   | 23.3               | 6.9                               | 9.3                               |
| M <sub>3</sub> G <sub>3</sub>                 | 99.0              | 12.0                                  | 11.2                                      | 26.1                | 125.1                                     | 3.5   | 23.3               | 7.4                               | 9.0                               |
| LSD <sub>0.05</sub>                           | NS                | 1.8                                   | NS  | 1.5                 | 33.7                                      | NS  | 2.5                | 1.2                               | 1.7                               |
| CV%   | 7.2               | 8.0                                   | 8.7                                       | 6.4                 | 14.6                                      | 11.5  | 6.1                | 10.6                              | 10.8                              |

155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170

### Water and soil salinity dynamics

Salinity causes unfavorable environment and hydrological situation that hinders the normal crop growth and development. The factors which contribute significantly to the development of saline soil are, tidal flooding during wet season (June to October), direct inundation by saline water, and lateral movement of saline ground water during dry season (November to May). The severity of salinity problem in Bangladesh increases with the desiccation of the soil. It affects crops depending on degree of salinity at the critical stages of growth, which reduces yield and in severe cases total yield is lost. Maximum salinity was observed during (March and April) at maximum tillering stage to flowering stage. Maximum salinity was also found at ridge and minimum in furrow.



**Figure 01. Water and Soil salinity status of experimental site**

171  
172  
173  
174

### Conclusions

176 Binadhan-10 was evaluated among three transplanting methods and three levels of gypsum at  
177 saline prone area. The highest grain yield (7.7 t ha<sup>-1</sup>) was produced in ridge and furrow method  
178 where gypsum rate was 150 kg ha<sup>-1</sup> (75 kg ha<sup>-1</sup> as basal + 75kg ha<sup>-1</sup> at ??(how many weeks)weeks  
179 after transplanting followed by (7.3 t ha<sup>-1</sup>) basal application of 150 kg ha<sup>-1</sup> gypsum along with N,  
180 P, K, Zn and Boron might be necessary to ensure satisfactory yield of rice in saline prone area  
181 under natural salinity condition.

182  
183

### References

184  
185

186 Chintala, R., McDonald, L. M., & Bryan, W. B. (2010). Grouping soils by taxonomy order to  
187 improve lime recommendations. *Commun. Soil Sci. Plant*, 41, 1594-1603.  
188 <http://dx.doi.org/10.1080/00103624.2010.485239>.

189 Fageria, N. K., & Knupp, A. M. (2014). Influence of Lime and Gypsum on Growth and Yield of  
190 Upland Rice and Changes in Soil Chemical Properties. *J. Plant Nut.*, 37(3), 1157-1170.  
191 <http://dx.doi.org/10.1080/01904167.2014.890219>.

192 Haque, S.A., 2006. Salinity problems and crop production in coastal regions of Bangladesh, Pak.  
193 *J. Bot.*, 38(5), pp.13591365.

194 Mainuddin, K., A. Rahman, N. Islam and S. Quasem. 2011. Planning and costing agriculture's  
195 adaptation to climate change in the salinity-prone cropping system of Bangladesh.  
196 International Institute for Environment and Development (IIED), London, UK.

197 Qadir, M., Noble, A. D., Schubert, S., Thomas, R. J., & Arslan, A. (2006). Sodicty-induced land  
198 degradation and its sustainable management: problems and prospects. *Land Deg. Devel.*,  
199 17, 661-676. <http://dx.doi.org/10.1002/ldr.751>.

200 Rahman, M.M. and M. Ahsan., 2001. Salinity constraints and agricultural productivity in coastal  
201 saline area of Bangladesh, *Soil Resources in Bangladesh: Assessment and Utilization*.

202 Rodriguez-Navarro, A., & Rubio, F. (2006). High-affinity potassium and sodium transport  
203 systems in plants. *J. Exp. Bot.*, 57, 1149-1160. <http://dx.doi.org/10.1093/jxb/erj068>  
204 Sharma, B. R., & Minhas, P. S. (2005). Strategies for managing saline/alkali waters for  
205 sustainable agricultural production in South Asia. *Agric. Water Manag.*, 78, 136-151.  
206 <http://dx.doi.org/10.1016/j.agwat.2005.04.019>.  
207 SRDI, 2010. Coastal Saline Soils of Bangladesh. Soil Resources Development Institute. Ministry  
208 of Agriculture, Dhaka, Bangladesh. pp.96.  
209 Wong, V. N. L., Dalal, R. C., & Greene, R. S. B. (2009). Carbon dynamics of sodic and saline  
210 soil following gypsum and organic material additions: A laboratory incubation. *Appl.*  
211 *Soil Ecol.*, 41, 29-40. <http://dx.doi.org/10.1016/j.apsoil.2008.08.006>.  
212  
213  
214  
215

UNDER PEER REVIEW