

Haematinic Effects of Aqueous Extract of Lophira lanceolata Leaves in Phenylhydrazine- induced Haemato- toxicity in Wistar Rats

By f f

Haematinic Effects of Aqueous Extract of *Lophira lanceolata* Leaves in Phenylhydrazine- induced Haemato-toxicity in Wistar Rats

Abstract

Aim: The haematinic activity of the aqueous extract of *Lophira lanceolata* leaves was investigated using rat model of phenylhydrazine- induced anaemia.

Methods and Results: Red Blood Cell (RBC) count, Haemoglobin (Hb) concentration and Packed Cell Volume (PCV) were analysed as indices of anaemia. Following phenylhydrazine administration to rats at a dose of 10mg/kg for 8 days, a significant decrease ($P<0.05$) the haematological parameters was observed indicating anaemia. However, treatment with grade doses (200, 400 and 800 mg/kg) of the aqueous extract of *Lophira lanceolata* leaves produced a significant ($P<0.05$) increase in the RBC count, Hb concentration and PCV time- and dose-dependently.

Conclusion: It was concluded that *Lophira lanceolata* leaves possess haematinic activity, making it useful in the management of anaemia.

Key words: Haematinic, *Lophira lanceolata*, Phenylhydrazine, Haemato- toxicity, Wistar rats

1.0 INTRODUCTION

Anaemia is the most prevalent nutritional deficiency disorder in the world. WHO defines anaemia as the condition in which the haemoglobin content of blood is lower than normal as a result of deficiency of one or more essential nutrients^[1]. According to WHO, more than 2 billion people worldwide suffer from anaemia with iron deficiency responsible for 50% of the cases^[2]. Anaemia occurs in women of reproductive age anaemia due to menorrhagia while in pregnancy the excess need of iron usually results to anaemia^[3]. Iron is important for formation of haemoglobin, myoglobin, cytochrome oxidase, peroxidase and catalase. The total quantity of iron in the body averages 4 to 5 grams, about 65 percent of which is in the form of haemoglobin^[4]. A man excretes about 1 mg of iron each day, mainly into the faeces. For a woman, the menstrual lost of blood brings the iron loss to a value of about 2mg/day^[5].

Through the ages man has learnt to take advantage of the many resources placed at his disposal by nature to meet his essential needs in all fields. As important reserves and sources of abundance, natural resources are indispensable for socio-economic development. According to Gbile^[6], the diversity of the flora in Africa partly explains the strength of traditional medicine.

38 Some Africans resort to orthodox methods of treatment due to mitigating circumstances such as high cost
39 of drugs, poverty and poor nutrition [7]. Many herbs have been used locally for the management of
40 anaemia and one of such plants is *Lophira lanceolata*. Therefore, this study was undertaken to
41 determine the haematinic potentials of the aqueous leaf extract of *Lophira lanceolata* in rats to
42 provide a scientific basis justifying the use of the plant in traditional medicine for the acclaimed
43 treatment of anaemia.

44 *Lophira lanceolata* is a tree of the tropical and sub-tropical regions. It is a common tree in
45 Cameroun, Nigeria and Sudan. It often grows gregariously on fallow land at the edge of forests.
46 It is a tree of 8 to 10 m tall, straight or twisted, with leaves alternate, clustered at the end of short
47 straight branches, glabrous, bright and blade oblong-lanceolate. The bark surface is corky grey
48 [8]. *Lophira lanceolata* is used in traditional medicine to treat several illnesses. The decoction of
49 the fresh leaves is administered orally against headaches, dysentery, diarrhoea, cough, abdominal
50 pains and cardiovascular diseases. It is also used on skin to cure wounds [8].

51

52

53 2.0 MATERIALS AND METHODS

54 2.1 Materials

55 2.1.1 Chemicals and drugs

56 All chemicals/drugs used were purchased locally.

57 2.1.2 Animals

58 Adult Wistar rats of either sex weighing 180–220g were used for this study. They were kept in
59 stainless steel cages under standard laboratory conditions. They were maintained on clean water
60 and standard rodent feed.

61 2.2 Methods

62 2.2.1 Plant Collection and Identification

63 The leaves of *Lophira lanceolata* were collected from a natural habitat in Okpella Area of Edo
64 State, Nigeria. The plants were identified at the herbarium unit of the Department of Biology,
65 University of Benin, Benin-City, Nigeria and voucher specimens were deposited for future
66 references.

67 2.2.2 Preparation of Extracts

68 The leaves of *Lophira lanceolata* were shade- dried for seven (7) days and pulverized using an
69 electric blender. Two thousand (2000) gram of the pulverized leaves was soaked in distilled
70 water for 72- hours. The resulting mixture was filtered using Whatmann filter paper (Size No1)
71 and the extract (referred to as LLAE henceforth) was concentrated using a free- dryer.

72 2.2.3 Acute Toxicity Study

73 The oral median lethal dose (LD₅₀) of the extract was determined in rats according to the method of
74 Lorke [9].

75 2.2.4 Experimental Design

76 A total of 25 adult wistar rats were weighed and divided into 5 groups of 5 animals each and
77 treated as follows:

78 Group 1: received distilled water (1 ml) daily (normal control).

79 Group 2: Anaemic and received distilled water (1 ml) daily (anaemic control).

80 Group 3: Anaemic and received LLAE at 200 mg/kg body weight/day

81 Group 4: Anaemic and received LLAE at 400 mg/kg body weight/day

82 Group 5: Anaemic and received LLAE at 800 mg/kg body weight/day

83 Rats were made anaemic by oral intubations of phenylhydrazine (10 mg/kg body weight) daily
84 for 8 days. Rats that developed anaemia with haemoglobin concentration lower than 13 g/dl were
85 selected for the study.

86 2.2.4.1 Haematological investigation

87 Blood was collected by ocular puncture after overnight fast. The blood was collected before
88 induction of anaemia, after induction of anaemia with PHZ and during 1, 2, and 3 weeks of
89 treatments. The haemoglobin concentration (Hb), red blood cell count and pack cell volume
90 (PCV) were assessed.

91 2.2.5 Statistical Analysis

92 Statistical analysis was carried out using SPSS version 20.0. All the data were expressed as mean
93 \pm SEM and the statistical differences between the means were determined by one way analysis of

variance (ANOVA) which was followed by Duncan test and difference between means at $P > 0.05$ were considered significant.

3.0 RESULTS

3.1 Acute Toxicity

The results of acute toxicity studies showed no mortality or physical changes in skin and fur, eyes and mucus membrane, respiratory rate, circulatory signs, autonomic and central nervous system effects up to a dose of 5000 mg/kg of aqueous extract of *Lophira lanceolata*. The oral LD_{50} of the extract was then taken to be > 5000 mg/kg.

3.2 Effect of the Administration of Aqueous Extract of *Lophira lanceolata* Leaves on Haematological Parameters of Wistar rats

Tables 1, 2, 3 and 4 shows the changes in haematological parameters of rats treated with phenylhydrazine and the extract. Following the administration of phenylhydrazine (PHZ) for 8 days, the RBC, Hb, and PCV of rats decreased significantly ($P < 0.05$) (Table 1) giving rise to macrocytic anaemia. After one week of treatment of the anaemic rats in groups 3, 4, and 5 with *Lophira lanceolata* extract, there was no significant ($P > 0.05$) changes in RBC, Hb, and PVC compared to the anaemic control (group 2) (Table 2). However, at 2 week- post treatment, The Hb, RBC and PCV significantly ($P < 0.05$) and dose- dependently increased with 800mg/ kg of the extract taking the values to near normal range (Table 3) Similarly, at 3 week- post treatment, the Hb, RBC and PCV significantly ($P < 0.05$) and dose- dependently increased but with 800mg/ kg of the extract this time taking the values to the normal range (Table 4). Figures 1-3 shows the changes in Hb, PCV and RBC per group during the 3- week treatment regimen. The Hb of the treated rats did not show significant changes within the first week of the experiment but there was a steady increase up to 3 week- post treatment (Figure 1). Similar observations were made for RBC and PCV (Figures 2 and 3 respectively).

5

124 **Table 1: Effect of Phenylhydrazine on some Haematological Parameters after 8- days of**
 125 **Administration**
 126

Treatment	1 Group 1	Group 2	Group 3	Group 4	Group 5
Hb (g/ dl)	16.31±1.11 ^b	11.23±1.11 ^a	12.06±1.13 ^a	11.17±1.23 ^a	11.33±1.25 ^a
RBC (x10 ⁶ µg /l)	6.29±0.80 ^b	3.29±0.81 ^a	3.07±0.33 ^a	3.05±0.44 ^a	3.07±0.53 ^a
PCV (%)	51.11±2.33 ^b	43.23±3.17 ^a	42.11±1.13 ^a	42.13±2.26 ^a	43.12±3.01 ^a

6

127 Data are presented as mean ± SD. Data was analysed by one- way ANOVA followed by
 128 Duncan post- hoc test for multiple comparisons, (n=5). Mean values having different lower
 129 ca 30 alphabets as superscripts are considered significant (p< 0.05) within the rows. Group
 130 1: Control received 1ml distilled water. Groups 2-5: received 10mg/ kg phenylhydrazine for
 131 8 days

132

133

2

134 **Table 2: Effect of the Aqueous Extract of *Lophira lanceolata* on the Haematological**
 135 **Parameters of Wistar rats one week- post treatment**
 136

Treatment	1 Group 1	Group 2	Group 3	Group 4	Group 5
Hb (g/ dl)	16.58±2.01 ^b	11.73±1.01 ^a	12.16±1.13 ^a	12.22±1.78 ^a	12.05±1.33 ^a
RBC (x10 ⁶ µg /l)	6.67±1.02 ^b	3.03±0.72 ^a	3.67±0.42 ^a	3.53±0.79 ^a	3.77±0.30 ^a
PCV (%)	51.55±2.13 ^b	43.43±2.42 ^a	44.21±1.32 ^a	45.23±3.21 ^a	45.22±2.99 ^a

6

137 Data are presented as mean ± SD. Data was analysed by one- way ANOVA followed by
 138 Duncan post- hoc test for multiple comparisons, (n=5). Mean values having differen 3 lower
 139 case alphabets as superscripts are considered significant (p< 0.05) within the rows. Group
 140 1: Normal control and received 1ml distilled water. Group 2: Anaemic control and received
 141 1ml distilled water. Group 3: Anaemic and received 200 mg/ kg LLAE, Group 4: Anaemic
 142 and received 400 mg/ kg LLAE, Group 5: Anaemic and received 800 mg/ kg LLAE

143

144

145

146 Table 3: Effect of the Aqueous Extract of *Lophira lanceolata* on the Haematological
147 Parameters of Wistar rats two week- post treatment

Treatment	Group 1	Group 2	Group 3	Group 4	Group 5
Hb (g/ dl)	16.98±1.45 ^b	11.43±1.25 ^a	12.12±1.32 ^a	14.28±1.23 ^{ab}	16.12±2.21 ^b
RBC (x10 ⁶ µg /l)	6.29±0.80 ^b	3.13±0.52 ^a	3.68±0.48 ^a	4.31±0.99 ^{ab}	5.43±0.98 ^{ab}
PCV (%)	51.11±2.33 ^b	42.08±2.44 ^a	44.23±2.32 ^a	48.15±2.73 ^{ab}	50.45±2.91 ^b

148 Data are presented as mean ± SD. Data was analysed by one- way ANOVA followed by
149 Duncan post- hoc test for multiple comparisons, (n=5). Mean values having different
150 case alphabets as superscripts are considered significant (p< 0.05) within the rows. Group
151 1: Normal control and received 1ml distilled water. Group 2: Anaemic control and received
152 1ml distilled water. Group 3: Anaemic and received 200 mg/ kg LLAE, Group 4: Anaemic
153 and received 400 mg/ kg LLAE, Group 5: Anaemic and received 800 mg/ kg LLAE

154
28
155 Table 4: Effect of the Aqueous Extract of *Lophira lanceolata* on the Haematological
156 Parameters of Wistar rats three week- post treatment
157

Treatment	Group 1	Group 2	Group 3	Group 4	Group 5
Hb (g/ dl)	17.26±1.05 ^b	10.86±1.42 ^a	14.12±1.18 ^{ab}	16.99±1.42 ^b	17.01±2.06 ^b
RBC (x10 ⁶ µg /l)	6.97±1.01 ^b	3.25±0.63 ^a	5.31±0.41 ^{ab}	5.81±0.78 ^{ab}	6.93±1.01 ^b
PCV (%)	51.85±3.26 ^b	41.19±1.32 ^a	46.25±2.43 ^{ab}	47.31±1.23 ^{ab}	50.96±2.08 ^b

158 Data are presented as mean ± SD. Data was analysed by one- way ANOVA followed by
159 Duncan post- hoc test for multiple comparisons, (n=5). Mean values having different lower
160 case alphabets as superscripts are considered significant (p< 0.05) within the rows. Group
161 1: Normal control and received 1ml distilled water. Group 2: Anaemic control and received
162 1ml distilled water. Group 3: Anaemic and received 200 mg/ kg LLAE, Group 4: Anaemic
163 and received 400 mg/ kg LLAE, Group 5: Anaemic and received 800 mg/ kg LLAE

164
165
166
167
168




Figure 1: Changes in Haemoglobin concentration (g/dl) across the groups within the 3-week treatment period. Group 1: Normal control and received 1ml distilled water. Group 2: Anaemic control and received 1ml distilled water. Group 3: Anaemic and received 200 mg/ kg LLAE, Group 4: Anaemic and received 400 mg/ kg LLAE, Group 5: Anaemic and received 800 mg/ kg LLAE




Figure 2: Changes in RBC concentration ($\times 10^6 \mu\text{g/l}$) across the groups within the 3- week treatment period. Group 1: Normal control and received 1ml distilled water. Group 2: Anaemic control and received 1ml distilled water. Group 3: Anaemic and received 200 mg/ kg LLAE, Group 4: Anaemic and received 400 mg/ kg LLAE, Group 5: Anaemic and received 800 mg/ kg LLAE




Figure 3: Changes in PVC (%) across the groups within the 3- week treatment period. Group 1: Normal control and received 1ml distilled water. Group 2: Anaemic control and received 1ml distilled water. Group 3: Anaemic and received 200 mg/ kg LLAE, Group 4: Anaemic and received 400 mg/ kg LLAE, Group 5: Anaemic and received 800 mg/ kg LLAE

4

231 4.0 DISCUSSION

232 Phenylhydrazine produces both aryl and hydroxyl radicals when incubated with rat liver
 233 microsomes ^[10] and oxidized by hydrogen peroxide at pH 7.4 and 37°C ^[11]. The radicals induce
 234 oxidative stress in the red cell membrane resulting in haemolysis by lipid peroxidation ^{[12], [13], [14],}
 235 ^[15]. Sub-chronic intoxication of rats with PHZ (10 mg/kg/day for 8 days) results in marked
 236 haemolytic anaemia characterised by decreased RBC, Hb and PCV ^[16]. Similar results were
 237 obtained in our study when experimental rats were administered PHZ (Table 1).

1

238 The main function of the RBC is the transportation of oxygen to tissues in the body. As such, any
 239 pathological or physiological condition that affects the RBC alters its function and this may be
 240 detrimental to the body. In this study PHZ altered the function of RBC by haemolysis
 241 characterised by decreased levels of RBC, Hb and PCV. However, our results indicated that the
 242 aqueous leaf extract of *Lophira lanceolata* markedly increased the concentration of
 243 haemoglobin, red blood cell count and the packed cell volume beginning from 2 week after
 244 treatment. It was also observed that the recovery of the treated groups was dose related with the
 245 highest dose of 800 mg/kg effecting the highest change. At the third week of the experiment,
 246 treatment of anaemic rats with *Lophira lanceolata* increased the RBC, Hb and PCV to near
 247 normal values. *Lophira lanceolata* is the well known source of minerals, Sterols, Proteins and
 248 other vitamins. These chemical constituents of the plant might be responsible for the haematinic
 249 activity.

250 5.0 Conclusion

2

251 The oral administration of the aqueous leaf- extract of *Lophira lanceolata* significantly improved
 252 the haematological parameters affected by phenylhydrazine administration. From this study, it
 253 was inferred that *Lophira lanceolata* leaves possess haematinic potential, therefore it could be
 254 useful in the management of anaemia.

255

256

257

258

259

260

REFERENCES

- [1]. Kawaljit kaur, B.D. (2014). Arya girls college,jalandar cantt,Punjab, *European journal of zoological research*,3(1):32-36.
- [2]. Eb 29:n mehdi and hany sady (2014). Prevalence and risk factors of iron deficiency anaemia, *American journal of health research*,2(5):319-326
- [3]. Ramesh chellan, lopamudra paul (2010); prevalence of iron deficiency anaemia in india, *journal of population and social studies*,19.
- [4]. Sembulingam, K. and Prema Sembulingam, Iron and Haemoglobin metabolism, *Essentials Of Medical Physiology*, fourth edition 2007, p.no.72.
- [5]. Arthur. C. (2014). Guyton and John.E.Hall, *Blood, Iron Metabolism, Text book of medical physiology*, 9th edition, p 430.
- [6]. Gbile, R. (1986). Haematology of conventionally maintained Lac P Wistar rats during the first year of life. *Lab. Anim.* 16: 198-200.
- [7]. Brown D. (2001). The Herb Society of America New Encyclopedia of Herbs and their uses.New York: (D.K), 31
- [8]. Arbonier M. (2000) Arbres, arbustes et lianes des zones sèches d'Afrique de l'ouest. *CIRAD, MNHN, UICN*, 425-427.
- [9]. Lorke, D. 1983. "A new Approach to Practical Acute Toxicity Testing." *Archives of Toxicology* 54: 275-287.
- [10]. Gannett PM, Lawson TS, Kolar C, Toth B (1997). Aryl radical formation during the metabolism of arylhydrazines by microsomes. *Chem. Res. Toxicol.* 10 (12): 1372-1377.
- [12]. McMillan DC, Jensen CB, Jollow DJ (1998). Role of lipid peroxidation in dapsone induced haemolytic anaemia. *J. Pharmacol. Exp. Ther.* 287 (3): 868-876.
- [13]. Cighetti G, Debiasis S, Paroni R, Allevi (1999). Free and total malondialdehyde assessment in biological matrices by GC/MS. What is needed for an accurate detection? *Analytical Chemistry*. 266 (2): 222-229.
- [14]. Zimmermann L, Antebi H, Morvan BC, Alcindor G (1997). In vitro peroxidation of plasma and erythrocyte lipids during WR 1339 induced hyperlipidemia in Wistar rats. *Ann. Pharm. Fr.* 55 (6): 246- 253.
- [15]. Nelson C, Erikson K, Pinero DJ, Beard JL (1997). In Vivo dopamine metabolism is altered in iron deficient anaemic rats. *J. Nutr.* 127(12): 2282-2288.
- [16]. Unami A, Nishina N, Terai T, Sato S, Tamura T, Noda K, Mine Y (1996). Effect of cisplatin on erythropoietin production in rats. *J. Toxicol. Sci.* 21(3): 157-65.

Haematinic Effects of Aqueous Extract of *Lophira lanceolata* Leaves in Phenylhydrazine- induced Haemato-toxicity in Wistar Rats

ORIGINALITY REPORT

43%

SIMILARITY INDEX

PRIMARY SOURCES

1	academicjournals.org Internet	527 words — 16%
2	www.ajol.info Internet	97 words — 3%
3	Wang, J.. "Structural features and hypoglycaemic effects of <i>Cynomorium songaricum</i> polysaccharides on STZ-induced rats", <i>Food Chemistry</i> , 20100515 Crossref	87 words — 3%
4	ayurvedika.ru Internet	73 words — 2%
5	Medinat. Y. Abbas, Jane. I. Ejiofor, Musa. I. Yakubu. "Acute and chronic toxicity profiles of the methanol leaf extract of <i>Acacia ataxacantha</i> D.C (Leguminosae) in Wistar rats", <i>Bulletin of Faculty of Pharmacy, Cairo University</i> , 2018 Crossref	60 words — 2%
6	www.scielo.br Internet	52 words — 2%
7	www.cibtech.org Internet	31 words — 1%
8	ijod.uaeu.ac.ae Internet	28 words — 1%
9	www.imedpub.com	

	Internet	26 words — 1%
10	www.tjpr.org Internet	26 words — 1%
11	whatshelikes.in Internet	26 words — 1%
12	www.ijpcsonline.com Internet	25 words — 1%
13	Olubunmi Abosede Wintola, Anthony Jide Afolayan. "Toxicological Evaluation of Aqueous Extract of Lauridia tetragona (L.F.) R.H. Archer Leaf in Wistar Rats", International Journal of Pharmacology, 2018 Crossref	23 words — 1%
14	svict.org Internet	22 words — 1%
15	www.cerbafaso.org Internet	18 words — 1%
16	www.herbsociety.org Internet	17 words — 1%
17	P. Berander. "Using students as subjects in requirements prioritization", Proceedings. 2004 International Symposium on Empirical Software Engineering, 2004. ISESE '04., 2004 Crossref	16 words — < 1%
18	www.ijbs.org Internet	16 words — < 1%
19	jahm.in Internet	15 words — < 1%
20	eprints.usm.my Internet	

15 words — < 1%

- 21 Adebiyi, Adedeji O., Ayodele A. Oyedeji, Elijah E. Chikwendu, and Olubunmi A. Fatoke. 14 words — < 1%

"Phytochemical Screening of Two Tropical Moss Plants: *Thidium gratum* P. Beauv and *Barbula indica* Brid Grown in Southwestern Ecological Zone of Nigeria", American Journal of Analytical Chemistry, 2012.

Crossref

- 22 Ismaila Alhaji Umar, Uche Samuel Ndidi, Abdulmalik Salman Abdullahi, Amos Ohilebo, Emmanuel Taiye Oluwafunmini. " antitrypanosomal and antioxidant activities of methanol extracts of selected Nigerian plants ", Journal of Herbs, Spices & Medicinal Plants, 2017 13 words — < 1%

Crossref

- 23 utq.edu.iq 12 words — < 1%
Internet

- 24 irjponline.com 12 words — < 1%
Internet

- 25 etheses.saurashtrauniversity.edu 11 words — < 1%
Internet

- 26 Howlader, Islam. "Antinociceptive and neuropharmacological activities of ethanolic extract of the fruits of *Ficus hispida* Linn.", African Journal of Pharmacy and Pharmacology, 2012. 11 words — < 1%

Crossref

- 27 www.caunesp.unesp.br 10 words — < 1%
Internet

- 28 www.bioline.org.br 10 words — < 1%
Internet

- 29 ejournal.undip.ac.id 9 words — < 1%
Internet

30 Gakunga, Ndukui, Godfrey Kateregga, Larry Sembajwe, and John Kateregga. "Antidiarrheal activity and Phytochemical profile of the ethanolic leaf extract of *Leonotis nepetifolia* (Lion's ear) in Wistar albino rats", *Journal of Intercultural Ethnopharmacology*, 2013.

Crossref

31 Adzu, B.. "Studies on the use of *Cassia singueana* in malaria ethnopharmacy", *Journal of Ethnopharmacology*, 200310

Crossref

32 Badole, Sachin L., Shweta N. Shah, Naimesh M. Patel, Prasad A. Thakurdesai, and Subhash L. Bodhankar. "Hypoglycemic Activity of Aqueous Extract of *Pleurotus pulmonarius*. in Alloxan-Induced Diabetic Mice", *Pharmaceutical Biology*, 2006.

Crossref

33 www.ajpcr.com

Internet

34 Vongtau, H.O.. "Anti-nociceptive and anti-inflammatory activities of the methanolic extract of *Parinari polyandra* stem bark in rats and mice", *Journal of Ethnopharmacology*, 200401

Crossref

35 Stanley O. Aniagu. "Short-term toxicity studies of *Ficus thonningii* Blume (Moraceae) leaf extract in rats", *International Journal of Food Science & Technology*, 3/2008

Crossref

36 Ladeji, O.. "Hypoglycemic properties of aqueous bark extract of *Ceiba pentandra* in streptozotocin-induced diabetic rats", *Journal of Ethnopharmacology*, 200302

Crossref

37 www.unu.edu

Internet

38 repository.up.ac.za

Internet

8 words — < 1%

39 Ojewole, J.A.O.. "Evaluation of the anti-inflammatory properties of *Sclerocarya birrea* (A. Rich.) Hochst. (family: Anacardiaceae) stem-bark extracts in rats", *Journal of Ethnopharmacology*, 200304 7 words — < 1%
Crossref

40 Udoamaka F. Ezuruike, Jose M. Prieto. "The use of plants in the traditional management of diabetes in Nigeria: Pharmacological and toxicological considerations", *Journal of Ethnopharmacology*, 2014 7 words — < 1%
Crossref

41 McGregor, Douglas. "Methyl tertiary-Butyl Ether: Studies for Potential Human Health Hazards", *Critical Reviews in Toxicology*, 2006. 6 words — < 1%
Crossref

42 J.J. Klawe. "IRON | Biosynthesis and Significance of Heme (Haem)", Elsevier BV, 2003 6 words — < 1%
Crossref

EXCLUDE QUOTES OFF
EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE MATCHES OFF