

MICROSCOPIC ILLUSTRATION OF *PELARGONIUM X HORTORUM* (GERANIACEAE)

ABSTRACT: For identification of a crude drug, there are several parameters which standardize it for sure. Microscopic features describe a crude drug very well. Chances of adulteration are very common due to morphological similarities in different species of drugs and to avoid such confusions, standardization via microscopy helps to create a valuable profile of a given crude drug. Involving different parts of plant drug in microscopy viz leaves, stem, roots and flower etc. helps a lot in identifying the original drug. Transverse section of different parts of plant, powder microscopy and determination of leaf constants like stomatal number, stomatal index, vein islet number, vein termination number and palisade ratio of *Pelargonium x hortorum* describes the basic features of the drug and authenticate it as the original one.

KEYWORDS: *Pelargonium*, Identification, Microscopic evaluation.

INTRODUCTION:

Pelargonium x hortorum L.H. Bailey is a species of the genus *Pelargonium* commonly used as an ornamental plant. *Pelargonium x hortorum* species is probably a hybrid species between *Pelargonium-P. zonale* and *Pelargonium-P. inquinans* belonging to family Geraniaceae. Plant can be propagated by stem cuttings and requires peaty or loamy soil and flourishes in sunny conditions^[1]. The plant varies from height of 45 to 50



Fig.1a: General view of the leaves of *P. x hortorum*

Formatted: Font: Italic

cm with fragrant green colored decorative leaves up to 5 to 7.5 cm in length with reticulate venations and crenate margin. Flowers appears in many colors like red, pink, orange or white having five petals positioned around the center as ball shaped clusters. The inflorescence is long rigid peduncle. Generally tap root system is present in the plant.



Fig.1b: Flowers of *P.x hortorum*

Comment [SS2]: Fig 1a and Fig 1b don't go in sentence.

Comment [SS1]: what is the citation of this sentence ??

Different species of *Pelargonium* are available abundantly in nature with immense pharmacological potential and exhibit antifungal ^[2], mosquito repellent ^[3], anxiolytic, antidepressant ^[4] and pediculicidal activities ^[5]. *Pelargonium* derived essential oils (citronellol, geraniol, p-menthone and α -pinene etc.) are extensively used in perfumery, cosmetics, soaps, creams and aromatherapy products ^[6-7]. The aim of this study is to standardize the particular species of *Geranium* i.e *G.hortorum* and as such no microscopic profile has been available of this species.

Comment [SS3]: Where did this species come from? This species is no the world.

Comment [SS4]: The aim of this study, is insufficient.

MATERIAL AND METHODS:

Collection of plant:

The plant is collected in the month of January 2019 from Govt. College of Pharmacy Rohru, Distt. Shimla, Himachal Pradesh, India and the collected samples were subjected to microscopic examination.

Microscopy: Anatomical sections of the fresh leaf, petiole, stem and roots were prepared for the microscopic studies and examined under Trinocular microscope Olympus-CH-20i model and compound microscope.

For determination of leaf constants like stomatal number, stomatal index, vein islet number, vein termination number and palisade parenchyma ratio camera lucida was used Camera lucida-

Stomatal number and index determination: The fragment of leaf was cleared by boiling with chloral hydrate solution. Epidermal layer was then peeled out using forcep. A square of 1mm was drawn on a drawing paper using Camera lucida and stomata were counted and stomatal index was calculated using formula-

$$\text{Stomatal index (S.I.)} = \frac{S}{E+S} \times 100$$

Where S= Number of stomata,

E= Number of epidermal cells

42

43 *Determination of Vein-islet and termination number:* Fragments of leaf was cut in 2 mm x 2 mm
44 rectangular shape and boiled in chloral hydrate solution followed by dilute hydrochloric acid for
45 few minutes. A square of 1mm was drawn on a drawing paper using Camera lucida and vein
46 islets and terminations were counted.

47 *Determination of palisade ratio:* Fragments of leaf was cut in 2 mm x 2 mm rectangular shape
48 and boiled in chloral hydrate solution followed by dilute hydrochloric acid for few minutes. A
49 square of 1mm was drawn on a drawing paper using Camera lucida and palisade cells were
50 focused underlying four epidermal cells ^[8-9].

51

52

53

54

55

56 **RESULTS:**

57 The microscopic examination of the plant consists of its transverse section of leaf, petiole, stem
58 and root. The results of the T.S, powder characteristics and leaf constants are given in the Figure
59 2(a-h).

60




Figure: 2 a: Trichomes- CTR: Unicellular, uniseriate covering trichome, GTR: Unicellular, uniseriate Glandular trichome




Figure: 2 b: Wavy epidermal cells and stomata-EP: Epidermal cells, ST: Stomata




Figure: 2 c: Transverse section of Petiole-VB: Vascular bundles, CR: Cortex, TR: Trichome




Figure: 2 d: Transverse section of leaf




Figure: 2 e: Transverse section showing mesophyll layer (palisade cells) in leaf

Comment [SS5]: Spongy parenchyma, upper and lower epidermis layers should be shown over the Figure 2e.




Figure: 2 f: Transverse section of root-TR: Trichome, CR: Cortex, PH: Pith, XV: Xylem vessels




Figure: 2 g: Transverse section of stem-ED: Endodermis, VB: Vascular bundle, CR: Cortex, PT: Pith




Figure: 2 h: Powder microscopy of leaf-CR: Crystals, TR: Trichomes

Comment [SS6]: What is the Crystal type ???

Conclusion:

The macroscopic study reveals the physical characteristics of plant whereas the microscopic studies give us vital information about the histological arrangement of different plant parts. Transverse section of leaf showed the presence of unicellular covering, glandular trichomes, anomocytic stomata, mesophyll type is dorsiventral, bilayered ~~rectangular~~ cylindrical arrangement of palisade parenchyma cells, wavy walled epidermal cells, five to six layered parenchymatous tissue, few calcium oxalate crystals and vascular bundles. In petiole five to ten celled layered cortex are present with circularly arranged vascular bundles in parenchymatous tissue. Stem ~~portion~~ cross-section showed presence of endodermis with 15-20 layered cortex, circularly arranged vascular bundles and pith. Roots cross - section showed presence of covering trichomes with 15-20 layered cortex and xylem vessels with lignified phloem. Powder characteristic of leaf showed presence of prismatic calcium oxalate crystals, abundantly scattered unicellular covering trichomes and ~~fragments~~ fragments of parenchymatous cells. Evaluation of different leaf constants like stomatal number (262-280), stomatal index (12-16.6), vein islet number (4-8), vein termination number (12-18), palisade ratio (2-6) helps in framing the microscopic illustration of Pelargonium-P.x hortorum, family Geraniaceae.

Comment [SS7]: How many layers of endodermis ?

Comment [SS8]: The roots do not contain covering trichomes. This is scientifically a big mistake. Comment the root section well.

REFERENCES:

1. https://en.wikipedia.org/wiki/Pelargonium_%C3%97_hortorum (Accessed on dated 10 Jan. 2019).
2. Dianez F, Santos M, Parra C, Navarro MJ, Blanco R, Gea FJ. Screening of antifungal activity of 12 essential oils against eight pathogenic fungi of vegetables and mushroom. Lett Appl Microbiol. 2018; 67(4):400-410.

- 84 3. Ralambondrainy M, Belarbi E, Viranaicken W, Baranauskiene R Venskutonis
85 PR, Despres P, Roques P, El Kalamouni C, Sélambarom J. In vitro comparison of three
86 common essential oils mosquito repellents as inhibitors of the Ross River virus. PLoS
87 One. 2018; 13(5):e0196757.
- 88 4. Abouhosseini Tabari M, Hajizadeh Moghaddam A, Maggi F, Benelli G. Anxiolytic and
89 antidepressant activities of *Pelargonium roseum* essential oil on Swiss albino mice:
90 Possible involvement of serotonergic transmission. Phytother Res. 2018; 32(6):1014-
91 1022.
- 92 5. Limoncu ME, Balcioglu C, Oyur T, Zeybek G, Zeybek. In vitro Investigation of the
93 Pediculicidal Activities of the Volatile Oil Components of Some Medical Plants Raised
94 in Turkey. Turkiye Parazitol Derg. 2017; 41(4):208-213.
- 95 6. Farukh S. Sharopov, Hanjing Zhang and William N. Setzer. Composition of geranium
96 (*Pelargonium graveolens*) essential oil from Tajikistan. American Journal of Essential
97 Oils and Natural Products 2014; 2 (2): 13-16.
- 98 7. Saraswathi J., Venkatesh K., Nirmala Baburao, Majid Hameed Hilal and A. Roja Rani.
99 Phytopharmacological importance of *Pelargonium* species. Journal of Medicinal Plants
100 Research Vol. 2011; 5(13): 2587-2598.
- 101 8. Khandelwal KR. Practical Pharmacognosy Techniques and Experiments. 9th ed. Nirali
102 Prakashan, Pune, 2003.
- 103 9. Evans WC. Trease and Evans Pharmacognosy. WB Saunders Ltd., London; 2002.
- 104
105
106