

**PREVALENCE OF INTESTINAL PARASITIC
INFECTIONS AMONG PATIENTS ATTENDING
USMANU DANFODIYO UNIVERSITY TEACHING
HOSPITAL, SOKOTO, NIGERIA**

ABSTRACT

Background: Intestinal parasitic infection is one of the major health issues in developing countries particularly in Sub-Saharan Africa. It has been estimated to affect about 3.5 billion people globally and 450 million people are thought to be ill as a result of such infections, the majority being children.

Aims: The study aimed to determine the prevalence and associated risk factors of intestinal parasitic infections among patients attending Usmanu Danfodiyo University Teaching Hospital, Sokoto, Nigeria

Study Design: This was a cross-sectional, descriptive study

Place and Duration of Study: This study was conducted among patients attending Usmanu Danfodiyo University, Teaching Hospital, Sokoto, Sokoto state, between May to November 2017.

Methodology: A total of 243 participants were enrolled in the study. Standard parasitological examination was carried out on stool samples using microscopy followed by formal ether concentration methods

Results: Finding revealed that 29 (12%) were positive for intestinal parasitic infections. Males recorded higher prevalence than the females with 19 (11.9%) and 10 (11.8%), respectively. ~~Intestinal parasites continue to remain a serious public health problem in North-western Nigeria.~~

Conclusion: ~~Intestinal parasites continue to remain a serious public health problem in North-western Nigeria.~~ Low level of education, occupational status, poor water supply were among the significant risk factors for these infections. Creating awareness, level of sanitation, water supply and deworming programme among school children will reduce prevalence and intensity of parasitic infections among the study community.

Keywords: Prevalence study, Intestinal parasitic infection, UDUTH, Sokoto State, Nigeria.

34

35 1.0 INTRODUCTION

36 Intestinal parasitic infection is one of the major public health burden in developing countries
37 particularly in Sub-Saharan Africa. It has been found to affect about 3.5 billion people globally and
38 450 million people are thought to be ill as a result of such infections, the majority being children [1]. In
39 Nigeria, intestinal helminthes infections have continued to prevail because of poor standards of living,
40 poor environmental sanitation and ignorance of simple health promoting behaviours [2,3]. Intestinal
41 helminthes infections are most common in school age children and they tend to occur in high intensity
42 in this age group [4,5,6].

43 These infections have been associated with an increased risk for nutritional anaemias, protein energy
44 malnutrition, growth deficits in children, physical weakness and low educational performance of
45 school children [7,8] and [9] and also causing high morbidity and mortality rate.

46 Parasitic infections are governed by behavioural factors biological, environmental, socioeconomic and
47 health systems factors. Local conditions such as quality of domestic and village infrastructure;
48 economic factors such as monthly income, employment and occupation and social factors such as
49 education influence the risk of infection, disease transmission and associated morbidity and mortality
50 [10,11]. These infections are more prevalent among the poor segments of the population. They are
51 closely associated with low household income, poor personal and environmental sanitation, and
52 overcrowding, limited access to clean water, tropical climate and low altitude. Intestinal parasitic
53 infections such as amoebiasis, ascariasis, hookworm infection and trichiuriasis are among the ten
54 most common infections in the world [12].


55 In Nigeria there is dearth of information on the magnitude of intestinal parasitic infections and
56 predictors among patients attending Usmanu Danfodiyo University Teaching Hospital, Sokoto,
57 Nigeria. Information generated could be used for planning public health control programmes which is
58 aim as an important step for initiation of treatment and prevention strategies as well as reducing
59 morbidity and mortality due to parasitic infections in the area.

60 3.0 MATERIALS AND METHODS

61 3.1 STUDY AREA

62 The study area is Usmanu Danfodiyo University Teaching Hospital, a tertiary health facility located in
63 Sokoto metropolis, the Sokoto State Capital. It serves as a referral centre for more than 10 million
64 people of the Nigerian States of Sokoto, Zamfara and Kebbi; and neighbouring Niger and Benin
65 Republic in the West African sub-region [13].

66 Sokoto State is located at the extreme part of North-Western Nigeria between longitude 3° and 7°
67 east and between latitude 10° and 14° north of the equator. It shares borders with Niger-Republic to
68 the North, Kebbi State to the South-West and Zamfara State to the East [13]. The state covers a total
69 land area of about 32,000 square kilometres and a population of 4,602,298 million based on 2013
70 projection [14]. Sokoto State has semi-arid climate and vegetation is largely Sudan Savannah with an
71 annual rainfall between 500 – 1300 mm and temperature ranges between 15°C and over 40°C
72 during warm days [13].



73

74 Figure 11. Map of Sokoto State Showing Study Area [13].

75

76 **3.2 STUDY DESIGN**

77 This is a cross-sectional descriptive study that was carried out on 243 samples collected from UDUTH
78 Service laboratory, from May to November 2017.

79 **3.3 SUBJECT AND SELECTION**

80 The subjects **were** selected or recruited in UDUTH Service laboratory using systematic sampling
81 method to recruit all patients that meet the inclusion criteria.

82 **3.3.1 Inclusion criteria:**

- 83 1. All patients with uncontaminated stool sample (formed, semi formed and unformed) were
84 recruited for study and
- 85 2. Patients who give their consent to participate in the study.

86 **3.3.2 Non inclusion criteria:**

- 87 1. Patients with stool sample contaminated with urine or mixed with soil were excluded from the
88 study and
- 89 2. Patients who refuse to give consent in the study.

90 **3.4 SAMPLE SIZE**

91 The sample size was calculated using the formula outlined below

92 $n = \frac{(z-a)^2 (p) (1-p)}{d^2}$ [15].

93 d^2

94 And with the prevalence (p) estimated according to the study that was carried out in Sokoto that
95 reveals that the intestinal parasitic infection has a prevalence of 17.5%. Hence, it was used to
96 calculate the sample size. [16].

97 Where; n= Sample size

98 Z= standard normal deviate at 95% (1.96)²

99 p = prevalence 17.5%, 17.5/100 = 0.175

100 ~~q = complement of p (1-p)~~

101 d = precision 5% (0.05)

102 $n = \frac{(1.96)^2 \times 0.175 \times (1-0.175)}{(0.05)^2}$

103 $n = 221$

104 Using an attrition rate of 10%. Therefore; 221 + (0.1 x 221)

105 Actual sample size = 243 patients

106 **3.5 SAMPLING METHOD**

107 A systematic sampling method was used to recruit all patients that meet the inclusion criteria.

108 The laboratory register had about six hundred patients (600) that submitted their stool for evaluation
109 in the previous year (January 2016 to December 2017). This was used to determine the sampling
110 frame.

111 $K=N/n: 600/243 = 2.5 \sim 3$

112 A sampling interval of 3 was achieved.

113 Using simple random sampling; the first patient was chosen between number 1 and 3 for the first
114 week of study.

115 For any randomly chosen numbered patient; thereafter a sampling interval of 3 would be used for the
116 subsequent patients that present themselves at the facility until the sample size is achieved.

117 **3.6 SAMPLE COLLECTION**

118 An approximate amount of 100g faeces was collected into clean, dry and screw cap, leak proof
119 containers.

120 **3.7 STUDY TOOL**

121 **The questionnaire was design as a structured questionnaire before it was administered** to obtained
122 patient information. It was structured into the following subheadings; demographic information, socio-
123 economic data, clinical history and laboratory investigation. The questionnaire was pretested and
124 validated at a similar site to the study area in the state specialist hospital, Sokoto; corrections was
125 made thereafter where necessary.

126 **3.8 SAMPLE PROCESSING**

127 **3.8.1 Macroscopy**

128 The procedure for macroscopy was done as outlined below;

129 1. Presence of worms: The presence of adult helminthes or segments example: *Ascaris*, *Taenia*
130 *species*, *Enterobius vermicularis* and gravid *Taenia species* would be determined.

131 2. Consistency (degree of moisture): This varies with diet but certain clinical conditions associated
132 with parasite presence may be suggested by particular consistencies. It were described as hard,
133 formed, semi-formed and diarrhoeic (watery).

134 3. Colour: Any abnormal colour example, pale yellowish passed in steatorrhoeac conditions such as
135 Giardiasis, dark or black-stools occur when iron or bismuth is taken or when there is intestinal
136 haemorrhage were determined.

137 4. Pathologic odour: This can be offensive or non-offensive and it were determined.

138 5. Abnormal features seen (composition): This can be mucus, blood or fat globules. They were
139 determined and reported as appropriate.

140 **3.8.2 DIRECT MICROSCOPIC EXAMINATION USING NORMAL SALINE AND IODINE** 141 **PREPARATION**

142 The procedure is outlined below;

143 About 1-2mg of stool were emulsified in 1-2 drops of normal saline (0.9%) or Lugol's iodine solution.
144 Then a cover-slip were placed and then the slide were scanned under the x10 and x40 objectives

145 lenses of a light microscope. Saline direct smear is used mainly for the detection of intestinal protozoa
146 trophozoites motility. Iodine direct smear allows the examination of the characteristics features of the
147 protozoa and the identification of the *Entamoeba histolytica/dispar* (~~*Entamoeba histolytica/dispar*~~) cyst
148 from the commensal *Entamoeba coli*. [17]

Formatted: Font: Italic

149 **3.8.3 SEDIMENTATION METHOD**

150 Formalin ethyl ether sedimentation method were used.

151 **3.8.3.1 PRINCIPLE**

152 Is a qualitative method that sediment various types of parasites by gravity or by centrifugal force [18].

153 **3.8.3.2 Formal ether concentration technique**

154 **Procedure**

155 A 0.5g of faecal sample was added to a glass container containing 10mls of 10% formalin and then
156 mixed thoroughly. A Funnel was placed on a gauge and strain into a 15mls centrifuge tube and
157 centrifuge for 2minutes at 1500 rpm. Then the supernatant was discarded and the sediment was re-
158 suspended into 10mls of physiological saline, and centrifuge for 2 minutes at 1500 rpm. The
159 supernatant was discarded and the sediment re-suspended again in 7mls of 10% formaldehyde, 3mls
160 of ether (diethyl) was also added. The tube with a glass stopper was closed and shaken vigorously
161 to mix, and then the stopper was removed and centrifuge for 2minutes at 1500 rpm. The supernatant
162 was poured out and the sediment carefully placed on a clean glass slide and covered with cover slip
163 and this was examined at x10 and x40 objective. [19].

164

165 **3.9 DATA COLLECTION METHODS**

166 Data was collected by the researcher himself. Data were entered independently at two separate
167 occasions using Microsoft Excel 2016. Double data entry analysis was done to ensure data quality.

168 **3.10 ETHICAL COSIDERATION**

169 Ethical clearance was obtained from the ethics and research committee of UDUTH, Sokoto. And
170 consent was sought from the patient prior to sample collection.

171 **3.11 STATISTICAL ANALYSIS**

172 Data was entered into the statistical package for social sciences (SPSS) version 20. Analysis for
173 categorical variable was carried out using Chi-square test to determine the association. Simple and
174 multiple logistic regression analysis was used to determine associated risk factors of the infections.
175 Values were considered **statistically** significant at $p < 0.05$.

176

177 **4.0 RESULT**

178 **Characteristics of the study population**The result of this study revealed that from a total of 243
179 **participants selected for this study, an overall prevalence of (12%) were found corresponding**
180 **to 29 subjects positive for intestinal parasitic infections.**

Formatted: Font: Bold

181 Table 1, Shows the socio-demographic characteristics of variable with respect to marital status,
182 gender, age group and tribe. Of the total study subjects 160 (65.3%) ~~are-were~~ males and 83 (34.7%)
183 female. ~~among~~ Among the subjects 38 (15.5%) were married and 205 (84.5%) single. The age ranges
184 shows, 21-25 years have the highest frequency of 83 (33.9%) followed by 16-20 years then 0- 5 years
185 and 26-30 years 13 (5.3%) with the least.

186 Based on the source of water participants consumed, those that drink sachet water had 72 (31.4%)
187 followed by those that consume tap water 62 (25.3%) then followed by those that consumed
188 river/stream 49 (20%) and well water 37 (15.1%) and lastly those that consumed other source of
189 water with 23 (9.4%). With regards to occupation, higher frequency was recorded among 114 (47.3%)
190 students, likewise highest among those that are civil servant 47 (19.2%) then unemployed with 41
191 (16.7%), then those that are business men with 26 (10.6%), while the least was found among those
192 that are farming 15 (6.1%) (Table 1).

193 Table 2, Shows the socio-demographic characteristics of variable with respect to frequency of eating
194 vegetables, type toilet facility, diarrhoea, dysentery, abdominal pain, fever and do you walk bare foot.
195 Out of 243 participants about 124 (51.4%) have high frequency of diarrhoea and 119 (48.6%) had not.

196 With regards to washing of hand before eating about 193 (79.6%) do not wash their hand before
197 eating while 50 (20.4%) wash their before eating (Table 1).

198 **Prevalence and distribution of the intestinal parasitic infection according to variables**

Formatted: Font: Bold

199 **The result of this study revealed that from a total of 243 participants selected for this study, an overall**
200 **prevalence of 12% were found corresponding to 29 subjects positive for intestinal parasitic infections.**

201 **Distribution of infection rate according to age groups** Likewise, highest among age group ranges 11-15
202 **years of (36.8%) and the** least among 31 and above years. There was a statistically significant
203 difference by aged group ($p < 0.004$). The males showed a higher prevalence of intestinal parasite
204 infections of **11.9%** than the females that shows a prevalence rate of (11.8%). However, this is not
205 statistically significant ($p > 0.05$). (Table 43).

206 Prevalence of intestinal parasitic infection among study population based on characteristic of the
207 variable(s) shows that those that consumed river/stream water 12 (26.5%) have highest risk of
208 intestinal parasitic infection, followed by those that drink other source of water with prevalence of 8
209 (24.8%) then followed by those that drink well water with 4 (10.8%), Tap water (4.8) and lastly those
210 that consumed sachet water have the lowest prevalence of 1 (1.4%). Comparing the different
211 prevalence rates in relation to intestinal parasites by water source is statistically significant ($p < 0.001$)
212 (Table 43).

213 The distribution of intestinal parasitic infection among study population base on frequency of eating
214 vegetables, walking bare footed, type of toilet facility, diarrhoea, dysentery, abdominal pain and fever.
215 Out of 243 participants about 20 (16.8%) are recorded with diarrhoea and 8 (7.1%) had not. There
216 was a statistically significant difference of ($p < 0.019$) by diarrhoea (Table 2).

217 The distribution of intestinal parasitic infection among study population base on the characteristic
218 variable (s) as shown in table 82. It was observed that those who eat less hawked food had the
219 highest risk of parasitic infection- with 97 (65.5%) followed by those that do not eat hawked food at all
220 with 69 (27.6%); while those who engage more often in eating hawked food with 97 (65.5%) were the
221 least infected with intestinal parasite. Comparing the difference in mode of eating hawked food has no
222 statistically significant difference ($p < 0.060$) (Table 2).

Comment [UM01]: Table 2 don concern distribution of parasitic infection but distribution of population according to variables

Comment [UM02]: Only 29 subjects are suffering from intestinal parasitic infection. So your distribution must concern only these 29.

223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241

UNDER PEER REVIEW

242

243

244 **Table 1: Socio-demographic characteristics of variable with respect to marital status, gender,**
 245 **age group, and tribe.**

Variable(s)	Frequency (F)	Percentage (%)
Marital status		
Married	38	15.5
Single	205	84.5
Gender		
Male	160	65.3
Female	83	34.7
Age group (years)		
0-5	33	13.5
6-10	19	7.8
11-15	19	7.8
16-20	55	23.3
21-25	83	33.9
26-30	13	5.3
31 and above	21	8.6
Tribe		
Hausa	200	82
Igbo	9	3.7
Yoruba	6	2.4
Fulani	20	8.6
Others	8	3.3
Educational status		
None	84	34.3
Informal	16	6.5
Primary	30	12.2
Secondary	40	16.3
Tertiary	73	30.6
Occupation		
Business	26	10.6
Farming	15	6.1
Civil servant	47	19.2
Unemployed	41	16.7
Student	114	47.3
Monthly income		
High	31	12.7
Average	108	44.9
Low	104	42.4
Water source		
Tap water	62	25.3
Well water	37	15.1
River/stream	49	20.0
Sachet water	72	31.4
Others	23	9.4

246

247

249 **Table 2: Socio-demographic characteristics of variable with respect to population**
 250 **characteristics of the variables**

Variable(s)	Frequency (F)	Percentage (%)
Frequency of eating vegetables		
Frequent	68	27.8
Not frequent	161	66.5
Not at all	14	5.7
Do you walk bare foot		
Yes	180	73.5
No	63	26.5
Type of toilet facility		
Pit latrine	55	23.3
Bucket latrine	45	18.4
Open space	43	17.6
Flush	100	40.8
Diarrhoea		
Yes	119	48.6
No	124	51.4
Dysentery		
Yes	62	24.9
No	181	74.7
Abdominal pain		
Yes	87	34.7
No	156	64.1
Fever		
Yes	32	12.2
No	211	86.5
Headache		
Yes	44	18
No	197	81.2
Do you wash your hand before eating		
Yes	50	20.4
No	193	79.6
Vomiting		
Yes	25	10.2
No	218	89.0
Frequency of eating Hawk food		
Often	50	6.9
Less	97	65.5
Not at all	67	27.6
Intensity of Infection		
No infection	214	88.2
Light infection	29	11.8

Comment [UM03]: 44+197=241

Comment [UM04]: 50+97+67=214

252

253 **Table 3 : Distribution of intestinal parasitic infection among study population with respect to**
 254 **the characteristics variable(s) listed below.**

Variables	Intestinal parasites				p-value ^a
	Infection		No infection		
	N	%	n	%	
Marital status					
Married	3	(7.90)	35	(92.1)	0.905
Single	26	(12.6)	181	(87.4)	
Gender					
Male	19	(11.9)	141	(88.1)	0.413
Female	10	(11.8)	75	(88.1)	
Age group (years)					
0-5	5	(15.2)	28	(84.8)	0.004 *
6-10	4	(21.1)	15	(78.9)	
11-15	6	(36.8)	12	(63.2)	
16-20	6	(10.5)	50	(89.5)	
21-25	5	(6.00)	78	(94.0)	
26-30	2	(15.4)	11	(84.6)	
31 and above	0	(0.00)	21	(100.0)	
Educational status					
None	7	(8.3)	77	(91.7)	0.021*
Informal	2	(6.5)	14	(87.9)	
Primary	3	(10.0)	27	(90.0)	
Secondary	29	(72.5)	11	(27.5)	
Tertiary	6	(8.0)	69	(92.0)	
Occupation					
Bussiness	3	(11.5)	23	(88.5)	0.905
Farming	1	(6.7)	14	(93.3)	
Civil Servant	7	(14.9)	40	(85.1)	
Unemploye	4	(9.8)	37	(90.2)	
Students	14	(12.1)	102	(87.9)	
Monthly income					
High	2	(6.5)	29	(93.5)	0.408
Average	16	(14.5)	94	(85.5)	
Low	11	(10.6)	93	(89.4)	
Water Source					
Tap water	3	(4.8)	59	(95.2)	0.001*
Well water	4	(10.8)	33	(89.2)	
River/Stream	12	(26.5)	35	(44.8)	
Sachet water	1	(1.4)	73	(98.6)	
Others	8	(24.8)	15	(65.2)	

255 Key:

256 a = Pearson chi-square test

257 n = Number of parasites

258 * = Statistically significant

259

Comment [UM05]: the total is 28 instead of 29

260 **Table 4: Distribution of intestinal parasitic infection among study population base on**
 261 **Characteristic variable(s) as shown in the table.**

Variable(s)	Intestinal parasite				p-value ^a
	Infection		No infection		
	N	%	N	%	
Frequency of eating vegetables					
Frequent	11	(16.2)	57	(83.8)	0.201
Not frequent	18	(11.0)	145	(89.0)	
Not at all	0	(0.00)	14	(100)	
Do you walk bare foot					
Yes	25	(13.9)	155	(86.1)	0.098
No	4	(6.20)	61	(93.8)	
Type of toilet facility					
Pit latrine	9	(15.8)	48	(84.2)	0.379
Bucket latrine	39	(18.1)	6	(20.7)	
Open space	41	(19.0)	2	(6.90)	
Flush	12	(12.0)	88	(88.0)	
Diarrhoea					
Yes	20	(16.8)	99	(83.2)	0.019 *
No	8	(7.10)	116	(92.9)	
Dysentery					
Yes	5	(8.20)	56	(91.8)	0.500
No	24	(13.1)	159	(86.9)	
Abdominal pain					
Yes	5	(5.90)	80	(94.1)	0.530
No	24	(15.3)	133	(84.7)	
Fever					
Yes	4	(13.3)	26	(86.7)	0.808
No	25	(11.8)	187	(88.2)	
Headache					
Yes	4	(9.1)	40	(90.9)	0.364
No	25	(23.7)	174	(87.4)	
Do you wash your hand					
Yes	8	(16.0)	42	(84.0)	0.307
No	21	(10.8)	174	(89.2)	
Vomiting					
Yes	3	(12.0)	22	(88.0)	0.979
No	26	(11.8)	194	(88.2)	
Habit of eating Hawk food					
often	2	(3.8)	50	(96.2)	
Less	19	(16.4)	97	(83.6)	
Not at all	8	(10.4)	69	(89.6)	

Comment [UM06]: 9+39+41+12=101?

Comment [UM07]: 28 instead of 29?

262 Key:a = Pearson chi-square test

263 0 = No infection

264 **5.1 DISCUSSION**

265 This study reveals a parasitic prevalence rate of 12% among 243 patients attending the Usmanu
266 Danfodiyo University Sokoto Teaching Hospital, which were selected at random from May to
267 November, 2017.

268 The low prevalence of this study is in line with the study observed in North western Nigeria of 15.67%
269 by [19] and 12% in South India by [20]. However, the results are considerably lower than studies
270 reported in North western Nigeria by [21], North central Nigeria by [22], western Tajikistan by [23] and
271 North eastern Ethiopia by [24]. The lower prevalence might be due to improved environmental
272 sanitation, better knowledge of personnel health and hygiene, economic and educational status of the
273 subjects found in the study area.

274 The present study revealed that males gender are little more susceptible to infection (11.9%) than the
275 females (11.8%), this finding was found to be similar with that reported by [26]. This might be due to
276 the common feeding pattern in which a great number of men eat outside their homes while on daily
277 activities to earn a living. And also due to the contamination of soil by human faeces, use of raw
278 sewage for agricultural purposes; use of waste water irrigated vegetables and contaminated imported
279 vegetables [27].

280 Prevalence is not dependent on sex among the sampled population which disagrees with the work of
281 [26] who observed a higher prevalence of intestinal parasite in females than in males. And the work
282 contrast with that of [28], who reported that male was found to have higher prevalence rate in his
283 study carried out in North western Ethiopia.

284 However, 11-15 years aged group and 6-10 years had a highest prevalence of 36.8% and 21.1%
285 respectively. This finding was found to be similar with that reported by [21, 29]. This study is also
286 similar to the work of [30], who reported highest prevalence in the ages 9-10 years among children
287 [30]. Even though WHO confirmed that intestinal protozoan parasite (IPP) are dependent on age and
288 greater severity of the infection is found in the younger children [31]. This could be attributed to the
289 different host responses and other related factors such as the nutritional status [32].

290 The most common intestinal parasitic infection identified in the community include amongst others *H.*
291 *nana*, *Ascaris lumbricoides*, *G. lamblia*, *E. histolytica* and Hookworm specie. However Hookworm and
292 *A. lumbricoides* recorded the highest prevalence of 15 (51%) followed by *G. lamblia* and *E. histolytica*
293 9 (31%) and *H. nana* recorded the least prevalence of 5 (17.2%). This finding was similar to those
294 reported in Ethiopia [28], and in contrast with the study in Nigeria [33].

295 In this study, occupation, monthly income eating of hawked food, presence of latrine and frequency
296 eating of vegetables were not significantly associated with intestinal parasitic infections. However,
297 according to the study conducted by [34] and [35] were strongly associated with infections. This is
298 more likely due to high level of education, better sanitation condition, better knowledge about the
299 faeco-oral transmission of intestinal parasite through their unwashed hands and the
300 contamination of vegetables with faecal materials in the farm. Season could be another important
301 predictor of intestinal parasitic infections especially during rainy season where agricultural activities is
302 said to be highest. This finding is in agreement with the findings of other researchers that indicated
303 seasonal variations contributed to the higher prevalence of the disease [36,37].

304 **5.2 CONCLUSION**

305 This present study revealed that there is low prevalence of intestinal parasitic infection among
306 patients attending Usmanu Danfodiyo University Teaching Hospital, Sokoto.

307 The different potential risk factors assessed in the study include occupation, educational status, water
308 source, and type of toilet facility were strongly associated with intestinal parasitic infection. However,

309 the low prevalence might be attributed to proper management of organic refuse, public health
310 Enlightenment about the risk of intestinal parasitic infections, adequate supply of clean water, proper
311 drainage and use of sites for defaecation.

312 Therefore, all stakeholders should give attention to raise awareness about control of intestinal
313 parasitic infection, personal and environmental hygiene, and improving the quality of drinking water
314 source.

315 REFERENCES

316 [1] WHO (2000). World Health Report 2000 Conquering Suffering Enriching humanity. World Health
317 Organisation: Geneva.

318 [2] Nwosu ABC (1981). The community of soil transmitted helminth infection of humans in a hyper-
319 endemic area of southern Nigeria. *Annals Trop Med Parasite*,**75**: Pp 197-203.

320 [3] Udonsu, J. K. (1984). *Necator americanus*: a cross-sectional study of rural community in relay to
321 some clinical signs. *Annal. Trop. Med. Parasitol*.**78**: 443-445.

322 [4] Albonico, M. Wright, V. Jape, K. Haji, H. J. Taylor, M. Savioli, L. Bickle, O. (2002). Soil transmitted
323 nematode infections and mebendazole treatment in Mafia Island School children. *Annal Trop Med*
324 *Parasitol*, **96**:717-726.

325 [5] Alvañiz AFA, de Castro EC, Tablizo BJB, Flores MJC, Maghirang ESV, Esmeli LAR, Pacificador
326 AY, Sumalapao DEP. Prevalence, physiologic effects, and risk factors of soil-transmitted helminth
327 infections among grade school children. *National Journal of Physiology, Pharmacy and*
328 *Pharmacology*. 2017;**7**(9):907-913.

329 [6] Flores MJC, Bautista YM, Carandang LGE, Go KM, Olalia MBIT, Maghirang ESV, De Peralta AS,
330 Esmeli LAR, Sumalapao DEP. Risk factors associated with intestinal parasitic infections among
331 school children. *Annual Research & Review in Biology*. 2018;**24**(4):1-6.

332 [7] Nokes, C. S. M. Grantham-Mcgregor, A.W. Sawyer, E.S. Looer. D.A.P Bundy (1992). Parasitic
333 helminthic infection and cognitive function in school children.

334 [8] Nokes, C. Bundy, DAP (1994). Does helminth infection affect mental processing and educational
335 achievement? *Parasitol Today*.**10**: Pp 14-8.

336 [9] Sackey, M. E. Weigel, M. M. Armijos, R. X. (2003). Predictors and nutritional consequences of
337 intestinal parasitic infections in rural Ecuadorian children. *J. Trop. Pediatr*.**49**: 17-23.

338 [10] Yakubu, N. Musa, G. Yakubu, S. E. (2003). Seasonal changes in the distribution and infection
339 rate of *Schistosoma* intermediate hosts in River Kubanni and its tributaries. *Bio Res Com* 2003,
340 **15**:207-214.

341 [11] Wang, K. Tang, D. Wang, M. Lu, J. Yu, H. Liu, J. (2009). *Journal of cell science* 122(Pt12):20556

342 [12] WHO (1987). Report Series 749: Prevention and control of intestinal parasitic infections. Geneva:
343 WHO; 1987.

344 [13] Sokoto State Business Directory (SSBD) (2007). A Publication of the Commerce Department,
345 Ministry of Commerce, Industry and Tourism, Sokoto. Pp 14-18.

346 [14] UNFPA Annual report (2013). Publisher UNFPA ISBN 978-0-89714-994-5.

347 [15] Taofeek, I. (2010). Sample Size Determination. Research Methodology and Dissertation Writing
348 for Health and Allied Health Professionals 1st edition, Cress Global Link Limited, Abuja, Pp 74-75.

- 349 [16] Muhammad, I. M., Umoru, A. M., Isyaka, T. M., (2014). Intestinal parasitic infections among
350 patients attending a Tertiary Health Institution in North-eastern Nigeria. *American Journal of Research*
351 *Communication*, **2**(6): 88-96.
- 352 [17] Cheesbrough, M. (2005). District Laboratory Practice in tropical countries part 1, Cambridge
353 University Press, and International U.K. **Pp** 184-205. Children in southeast Nigeria. *Journal 7,*
354 *Environmental health*. **1**(2); 8-64.
- 355 [18] Ochei, J. Kolhatkar, A. (2000). Medical Laboratory Science. Theory and Practice. Tata Mcgraw-
356 Hill Publishing Company Limited: New Delhi. 2nd Edition, **Pp** 331-349.
- 357 [19] Garba, D. D. Thomas, H. Z. Jatau, E. D. Inabo, H. I. (2014). Prevalence of intestinal helminths
358 among primary school children in chikun and kaduna south local Government Area of Kaduna state,
359 Nigeria. *J. Med. Res. Vol 2*(2), Pp 6-1.
- 360 [20] Baragundi, M. C. Sonth, S. B. Solahannwar, S. Patil, C. S. (2011). The prevalence of parasitic
361 infections in patients attending tertiary care hospital, *Nat J Bas Med Sci*; **2**(1):314.
- 362 [21] Kabiru, M., Mohammed, R.A., and Julia O. (2015). Intestinal Parasitic Infection and Assessments
363 of Risk Factors in North-Western, Nigeria: A community based study. *International Journal of Pharma*
364 *Medicine and Biological Sciences*, **4**:2
- 365 [22] Ikeh, E., M. Obadofin, B. Brindeiro, (2006). Intestinal parasitism in rural and urban areas of
366 North Central Nigeria, an update, *Internet J. of Microbiology*, vol. 2, no. 1.
- 367 [23] Matthys, B. M. Bobieva, G. Karimova, Z. Mengliboeva, V. Jean, Richard, (2011). Prevalence
368 and risk factors of helminths and intestinal protozoa infections among children from primary
369 schools in Western Tajikistan, *Parasites and Vectors*, vol. 4, no. 195, **Pp**. 1-13.
- 370 [24] Missaye, A. D. Mulat, A. Alemu, and Agersew, A. (2013). Prevalence of intestinal parasite and
371 associated risk factors among HIV/AIDS Patients with pre-ART and on-ART Clinic, North-east
372 Ethiopia, *AIDS Research Therapy J*, vol. 10, no. 7, **Pp**. 1-9.
- 373 [25] Okon, K.O. Moses, A. E. Zailani, S. B. Tettey, E. N. (2003). "Prevalence of human intestinal
374 parasites in a semi-arid area of Nigeria: A five year review". *Nig. J exptl app bio* **17**: 503-24.
- 375 [26] Brooker, S, Hotez P. J. Bundy, D. A. (2008). Hookworm-Related Anaemia. Among Pregnant
376 Women: *A Systematic Review of Trop*. **Pp** 2-4.
- 377 [27] Atu, B.O. Galadima, M. and Alice, F. (2006). Prevalence of intestinal parasites in Etulo, Benue
378 State Nigeria. *Nig. J. of Parasitology*; **7**: **Pp** 1-16.
- 379 [28] Gelaw, A. A. Belay, Nigssie, B. (2013). Prevalence of intestinal parasitic infections and risk
380 factors among School children at the university of Gondar Community School, North-west Ethiopia, a
381 cross sectional study, *BMC Public Health*, vol. 13, no. 1, pp. 257.
- 382 [29] Abou-EL, F. R. Salama, Taha, N. (2009). "Predictors of intestinal parasitic infection among pre-
383 school children in Rural Lower Egypt, *The Egyptian Journal of Community Medicine*, vol. 27, no. 1,
384 **Pp**. 1-9.
- 385 [30] Oguoma, V. M. Anyasodor A. E. Mbata, T. I. (2008). Valence of Intestinal Protozoan parasites
386 among children aged 5-12 years in Owerri Metropolis South Eastern Nigerian. *Trop. J. Health Sci*;
387 **15**(2): **Pp** 43-48.
- 388 [31] WHO (2005). De-worming for health and development. Report of the third global meeting of the
389 partners for parasite control.

- 390 [32] Adekunle, L. (2002). Intestinal parasites and nutritional status of Nigerian children. Afr. J.
391 Biomed.Res, 5: **Pp** 115-119.
- 392 [33] Akinbo, F. O. Omorigie R. Eromwan, R. (2011). Prevalence of intestinal parasites among
393 patients in tertiary hospital in benin city, Nigeria. North American journal of medical science 3(10):462-
394 464.
- 395 [34] Erko, B. Medhin, G. (2003). Human helmenhasis in wodogenet, southern Ethiopia,with emphasis
396 on geohelmethasis, *Ethiop MED J*: 41: **Pp** 333-44.
- 397 [35] Asrat, A. Tewodros, D. Alemayehu, W. (2011). Prevalence and risk factors of intestinal paraties
398 among Delgi school children, North Gondar, Ethiopia.Journal of parasitology and vector Biology Vol.
399 3(5) Pp 75-81. ISBN 21-41-2510
- 400 [36]Miller SA, Rosario CL, Rojas E, Scorza JV. Intestinal parasitic infection and associated symptoms in
401 children attending day care centres in Trujillo, Venezuela. *Trop Med Int Health*. 2003;8:342-7.
- 402 [37] Nasiri V, Esmailnia K, Karim G, Nasir M, Akhavan O. Intestinal parasitic infections among
403 inhabitants of Karaj City, Tehran province, Iran in 2006-2008. *Korean J Parasitol*. 2009;47:265-8..

UNDER PEER REVIEW