

Original Research Article

Response of Upland Rice to Nitrogen and Phosphorus Fertilization on Vertisols of Tigray, Ethiopia

Abstract

Nitrogen and phosphorus are often cited as the most limiting nutrients in agricultural soils of Ethiopia. Their availability in the soil solution also determines the growth and productivity of the soils and crops. Thus, experiments were conducted to investigate the response of upland rice and determine the optimum rates of N and P fertilizers. ~~The e~~Experiments were conducted at four sites of Tselemti district (Tsaeda-Kerni, Mizikir, Maitsebri and Sekota-Mariam), Northwestern Tigray, Ethiopia. A factorial combination of two factors, four levels of nitrogen (0, 23, 46, 69, kg N ha⁻¹ designated as N0, N1, N2 and N3, respectively) and four levels of phosphorus (0, 10, 20, 30 kg ~~P₂O₅P~~ ha⁻¹ designated as P0, P1, P2 and P3, respectively) were conducted on complete randomized block design (CRBD). Data on yield and yield components of upland rice were collected and subjected to ANOVA through SAS software V9. ~~The r~~Results showed that, ~~the~~ plant height, number of spikelet and total seed per panicle of upland rice were significantly increased when nitrogen and phosphorus fertilizers interacted at higher doses. ~~It was also observed~~ ~~Therer were~~ significant differences of main effects on panicle length, grain yield and straw yields of rice. ~~A s~~Significantly higher grain and straw yields were obtained in plots ~~received-receiving~~ 69 kg N ha⁻¹ compared to preceding treatments, and application of either 20 or 30 kg ~~P₂O₅P~~ fertilizer ha⁻¹ can also boost the yields of upland rice in the district. However, the optimum grain yield of rice to the application of N and P fertilizers was not found in the response curve which indicates an extra application rates ~~would-should~~ be tried. -Further, ~~the~~ highest agronomic ~~efficiency-efficiencies~~ were also observed in the lower rates of both main effects.

Formatted: Font: Italic

Formatted: Font: Italic

Key words: agronomic efficiency, response curve, nutrients, upland rice, yield and yield components

28 Introduction

29 In Ethiopia, rice is among the target commodities that have received due emphasis in promotion
30 of agricultural production, and as such it is considered as the “millennium crop” expected to
31 contribute to ensuring food security in the country. Although rice ~~is~~ has been introduced to the
32 country very recently, ~~rice~~ it has proven to be a crop that can assure food security in Ethiopia, the
33 second most populous nation in sub-Sahara Africa (SSA) with about 74 million people in 2007
34 [1]. The trend of rice production is increasing both in area coverage, participant farmers, and
35 production [2]. -Productivity in quintal per hectare has increased from 28.91 in 2012 to 28.97 and
36 the number of participant farmers increased from 115,832 to 119, 4970 in 2013 cropping season
37 [3].

38 ~~The n~~ National average yield of rice in Ethiopia is 2.9 t ha⁻¹ [4], which is much lower than the
39 world’s average rice yield of 4.54 t ha⁻¹ [5]. This is due to insect pest and diseases occurrence
40 (rice blast and brown spot), weeds and environmental fluctuations. In addition, poor agronomic
41 practices; human and institutional capacity and shortage of adapted varieties for different agro-
42 ecologies are the major rice production constraints in the country. -According ~~to~~ [6]
43 improvement of rice production has not been possible due to low soil fertility, inadequate
44 nutrient management, continuous cropping, and application of suboptimal levels of mineral
45 fertilizers among other factors.

Comment [J1]: Write the name of scientist(s)

46 Nitrogen and phosphorus are often cited as the most limiting nutrients in agricultural soil of
47 Ethiopia [7]. Relatively those nutrients are deficient in vertisols where nitrogen is subjected to
48 leaching while the limited availability of phosphorus observed due to several factors. Nitrogen
49 and phosphorus are required in huge amounts by crops than other nutrients. They are constituents
50 of several plant organs and enzyme mediated biochemical processes. Their availability in the soil
51 solution determines the growth and productivity of soils and crops. Yield and growth
52 enhancements of crops have been reported due to application of N and P fertilizers in the country
53 and world. According ~~to~~ [8] and [9] application of N and P fertilizers significantly increased ~~Teff~~
54 teff and ~~Wheat-wheat~~ grain yields respectively compared to control plots. Additionally, studies
55 from Fogera; area; Amhara region [6], Bambasi and Kamashi area; Benshangul-Gumuz region
56 [10] and [11], respectively; ~~;~~ Gambella region [12] and Uganda [13] showed a positive rice yield
57 increments due to N and P fertilization.

Comment [J2]: Write the name of scientist(s)

58 However, farmers of the district are reluctant in to using chemical fertilizers due to several
59 reasons and the absence of site specific fertilizer guideline particularly for Rice-rice is the main
60 problem as the blanket recommendation was initially developed for the major cereal crops of the
61 country. In the study area, application of macro nutrients especially N and P were not clearly
62 documented. Furthermore, effective rate of application for N and P to increase rice productivity
63 at four districts has not been established. Therefore, the objectives of this study were to
64 determine N and P level on yield and yield components of rice and quantify agronomic
65 efficiency of the crop.

66 **Materials and Methods**

67 **Description of the Study Area**

68 The experiments were conducted at four sites of Tselemti district, Northwestern Tigray,
69 Ethiopia. ~~The e~~Experimental locations were: Tsaeda-Kerni (located at 13° 33' 04.29.29" N, 38°
70 01' 52.18" E and 1116 masl), Mizikir (located at 13° 32' 56.99" N, 38° 03' 22.97" E and 1145
71 masl), Maitsebri (located at 13° 35' 28.68" N, 38° 08' 57.01" E and 1325 masl), and Sekota-
72 Mariam (located at 13° 32' 58.64" N, 38° 13' 33.91" E and 1145 masl).

73 Tselemti district is ~~found~~ 1172 km far from the capital city Addis Ababa and geographically
74 located 13°48'N latitude and 38°15'E longitude. It is bordered with Asgede Tsimbla, Welkait,
75 Tanqua Abregelle districts and Amhara region to the north, west, east and south, respectively. The
76 district covers an altitude ranging from 800 to 2870 m above sea level. The mean annual
77 temperature of the area is 16 °C (November–January) and 38 °C (February–May) minimum and
78 maximum, respectively. Some of the major crops grown in the area include sorghum, finger
79 millet, maize, chickpea and sesame [14].

80 **Experimental Treatments, Design, and Procedures**

81 The treatments were factorial combination of two factors, four levels of nitrogen (0, 23, 46, 69,
82 kg N_{ha}⁻¹ designated as N0, N1, N2 and N3, respectively) and four levels of phosphorus (0, 10,
83 20, 30 kg P₂O₅P_{ha}⁻¹ designated as P0, P1, P2 and P3, respectively). Urea (46% N) and Triple
84 Super Phosphate (46% P₂O₅) were used as fertilizer sources for N and P, respectively. At each
85 site then, the field experiment had arranged in randomized complete block design with three
86 replications. Gross plot size for the trial was 12.8 m² (4 m* × 3.2 m) and the space between

blocks and plots were 1 m and 0.5 m, respectively. Planting was done in row method with 20 cm inter row spacing and 5 cm planting depth at a rate of 70 kg/ha¹. The newly adopted upland rice variety, i.e., NERICA-13 was used as a test crop. All of the phosphorus fertilizer for each treatment (except the control plots) were applied at planting while Nitrogen-nitrogen fertilizers was split applied (the first 1/3 was applied at sowing, and the remaining at tillering stage). All other agronomic, cultural and management practices including weeding, disease and pests were followed the recommended practices of the crop.

Soil Sampling and Analysis

Composite soil samples for chemical and physical characterization were collected up to 20 cm soil depth from each field before fertilizer application. Each parameter were analyzed using standard procedure as prescribed by [15]. Soil texture was determined by the hydrometer method. Soil pH and EC was also determined using a pH meter from 1:2.5 soil:water ratio suspension and using EC meter from 1:5 soil water saturation extract respectively. As there is a lot of information about the interpretation of E_{Ce} in relation to plant growth, it was useful to convert EC (1:5) to E_{Ce} [16].

$$E_{Ce} \text{ (dSm}^{-1}\text{)} = EC_{1:5} \text{ (dS}_m^{-1}\text{)} \times \text{multiplier factor}$$

Where; the multiplier factor was used for heavy clay i.e. 5.8

The determination of soil organic carbon was based on the Walkley-Black chromic acid wet oxidation method (Ref. ?). The Kjeldhal process (digestion, distillation, and titration) was followed to determine the total nitrogen (Ref. ?). -Olsen Method (Bicarbonate extractable P) was used to extract and determine available phosphorus, using 0.5 M NaHCO₃ at adjusted pH 8.5 (Ref. ?). Determination of CEC at pH 7 was carried out with Ammonium Acetate method (Ref. ?).

Data Collection and Analysis

Grain Data on grain yield, straw yield, plant height, panicle length, number of tillers, effective tillers, number of total seeds and unfertilized seeds per panicle, thousand seed weight and harvest index was were collected. These data was were arranged in factorial format and subjected to

analysis of variance (ANOVA) using SAS statistical software (Ref. ?). The response curve and Agronomic use efficiency were also worked out in excel worksheets.

Result and Discussion

Soil Physicochemical characteristics of soil of the Experimental studies

Textural classes of Sekota Silassie, Maitsebri and Tsaeda Kerni experimental sites were clay loam and clay for Mizikir experimental site (Table 1). The pH of soils of the experimental sites varied from 5.8 to 6.5, which ranges from slightly acidic to moderately acidic [17]. Thus, the pH of the experimental site soils was within the range for productive soils. According to the ratings of [17] the organic carbon content of soils of experimental sites were medium and total nitrogen was low in all fields [18].

Table - 1:- Physicochemical characteristics of the study sites

Soil Physicochemical physicochemical Parametersparameters	Locations			
	Sekota_Silassie	Mizikir	Maitsebri	Tsaeda_Kerni
pH (1:2.5 H ₂ O)	5.8	6.5	6.2	6.1
EC (dS m ⁻¹)	0.38	0.11	0.21	0.38
Texture	% Sand	28	30	28
	(%)			
	% Clay	28	50	38
	(%)			
	% Silt (%)	44	20	34
Textural Class	Clay Loamloam	Clay	Clay Loamloam	Clay Loamloam
%OC (%)	2.44	1.75	2.20	2.10
CEC (meq./ 100_gm Soilsoil)	35	57.2	46.4	40.8
%TN (%)	0.0230	0.0448	0.0148	0.0145
Available P (ppmmg kg ⁻¹)	11.88	5.38	4.42	7.34

Formatted: Superscript

Formatted: Superscript

127 Available soil phosphorous (Olsen P) at Maitsebri and Sekota_Silassie sites was very low and
128 medium, respectively while it was low at Mizikir and Tsaeda Kerni sites [19]. According to the
129 ratings of [16], the CEC were high to very high for soils of the experimental sites.

130

131 **Yield ~~Attributes~~ attributes of Upland-upland Rice**

132 Application of ~~Nitrogen-nitrogen~~ and ~~Phosphorus-phosphorus~~ fertilizers individually and/or in
133 combination at different rates does not show significant difference ($P \leq 0.05$) on number of
134 tillers and effective tillers per plant of ~~Up-upland~~ rice ~~on-in~~ the district (Table 2). [10] had also
135 reported insignificant differences over fertilized and unfertilized plots due to Phosphorus
136 fertilization on tiller production of rice at Benshangul-Gumuz region of Ethiopia. On other hand,
137 there were interaction effects of nitrogen and phosphorus fertilizer rates on plant height, number
138 of spikelet and total seed per panicle and the fertilized plots have shown an increase over the
139 control. The result showed that, ~~the~~ plant height of upland rice significantly increased at higher
140 doses of fertilizer N and P applications. The tallest rice plants (94.87 cm) were recorded at plots
141 fertilized with 69 kg N ha⁻¹ and 20 kg ~~P₂O₅P~~ ha⁻¹ while the shortest plants measured from the
142 unfertilized plots. Comparable with this, rice plants treated with N2-P3, N3-P3, N3-P1 and N1-
143 P1 were tallest by 43%, 41%, 35% and 33% respectively over the unfertilized plants. In line
144 with the findings of the present study, it has been reported that the application of nitrogen and
145 phosphorus fertilizers significantly enhanced the growth of upland rice [6], [10], [11]. The
146 growth enhancement might be in line with remarkable increase of fertilizer application which
147 could boost availability in the soil solution and crop uptake.

148 Combined application of highest doses of nitrogen and phosphorus were produced highest
149 number of spikelets and total seeds per panicle. ~~The high~~ Greater spikelets were counted from
150 plots received 46 kg N ha⁻¹ along with 30 kg P ha⁻¹ followed by combined applications of higher
151 doses of N and P fertilizers (N3-P2, N3-P3 and N2-P2) significantly. Application of 69 kg N ha⁻¹
152 and 20 kg ~~P₂O₅P~~ ha⁻¹ also resulted in highest seed numbers (126.9) per panicle of rice than did
153 other plots, with an advantage of 106.5% over control. These results were comparable with plots
154 received combined applications of 46 kg N ha⁻¹ with 20 and 30 kg ~~P₂O₅P~~ ha⁻¹. Similar trend were
155 also observed by [6], [11], [20].

Formatted: Font: Italic

Hence, there were no interaction effects on panicle length and number of unfertilized seed per panicle of upland-rice. However, there were significant differences of main effect nitrogen and phosphorus on panicle length. The longest panicles was 20.19 cm recorded from plots treated with 46 kg N ha⁻¹, while the shortest panicles were found in the control plots for main effect nitrogen. Fertilized treatments with main effect phosphorus fertilizer had a significant difference in panicle length ($P < 0.05$) from the unfertilized treatment but not among each other [12] also observed an increment in the panicle length of rice plants due to applied nitrogen and phosphorus fertilizer effects at southwestern Ethiopia. Number of unfertilized seeds per panicle showed significant response to the application of phosphorus fertilizer, whereas no significant difference among nitrogen fertilizer rates. The lowest unfertilized seed number was found in plots received 20 kg P₂O₅ ha⁻¹ and the highest from plots treated with low and/or no phosphorus fertilizer. It was understood that phosphorus had a key role in grain filling and fertility of seeds [21].-

Table - 2:-Main effects of nitrogen and phosphorus fertilizer levels on yield components of upland-rice

Treatments	NTP ⁻¹	NETP ⁻¹	PH	PL	NSP ⁻¹	NTSP ⁻¹	NUSP ⁻¹
Nitrogen Levels (kg N ha⁻¹)							
N0	4.20 ^A	3.46 ^A	74.46 ^C	18.44 ^C	7.14 ^C	69.25 ^C	10.59 ^A
N1	4.23 ^A	3.47 ^A	79.42 ^B	18.95 ^{BC}	8.35 ^B	79.30 ^B	9.10 ^A
N2	4.38 ^A	3.52 ^A	84.37 ^{AB}	20.19 ^A	10.48 ^A	102.50 ^A	9.63 ^A
N3	4.15 ^A	3.49 ^A	86.62 ^A	19.96 ^{AB}	10.94 ^A	104.55 ^A	8.24 ^A
LSD	NS	NS	4.96	1.23	0.71	5.33	NS
Phosphorus levels (kg P₂O₅ ha⁻¹)							
P0	4.19 ^A	3.53 ^A	73.77 ^B	18.27 ^B	8.01 ^B	73.37 ^C	10.26 ^{AB}
P1	4.48 ^A	3.60 ^A	83.06 ^A	19.82 ^A	8.65 ^B	82.79 ^B	10.84 ^A
P2	4.13 ^A	3.43 ^A	83.16 ^A	19.76 ^A	9.96 ^A	99.12 ^A	7.94 ^B
P3	4.18 ^A	3.39 ^A	84.90 ^A	19.69 ^A	10.30 ^A	100.32 ^A	8.50 ^{AB}
LSD	NS	NS	4.96	1.23	0.71	5.33	2.35
N*P	NS	NS	**	NS	*	***	NS
CV %	34.62	39.23	15.16	15.8	19.12	14.90	62.28

Means in a column with the same letter are not significantly different ($P \leq 0.05$); NS = Non significant difference. NTP⁻¹= Number of tillers per plant; NETP⁻¹= Number of Effective Tillers per plant; PH= Plant height (cm); PL= Panicle Length (cm); NSP⁻¹ = Number of spikelets per panicle; NTSP⁻¹= Number of Total seed per panicle; NUSP⁻¹= Number of unfertilized seed per panicle

Formatted: Font: Italic

Formatted: Left, None, Space Before: 0 pt, Line spacing: single, Don't keep with next, Don't keep lines together

Comment [J3]: Write the significant differences in small letter(s) as a, b, c, etc. in whole and all tables

Formatted: Left, Space After: 0 pt, Line spacing: single

Formatted: Font: Italic

174 | **Yield of Upland-upland rice**

175 | The interaction-interactive effect of nitrogen and phosphorus fertilizers at different levels showed
 176 | no significant differences on grain yield and straw yields of upland-rice. However, analysis of
 177 | variance from the four sites showed that both the main factor N and P fertilizers effects on grain
 178 | and straw yield were significant ($P < 0.0001$). Fertilization of upland-rice with nitrogen fertilizer
 179 | at a rate of 23, 46 and 69 kg ha⁻¹ consistently increased grain yield and were significantly higher
 180 | (24%, 46% and 63%, respectively) than the unfertilized fields. It is also observed that a
 181 | significant differences among each treatments. Rice fields treated with phosphorus fertilizer at
 182 | the rate of 20 and 30 kg ha⁻¹ had significantly higher grain yield (34% and 43%, respectively
 183 | over control) than preceding treatment, though P increase beyond 20 kg ha⁻¹ did not give
 184 | statistically significant ($P > 0.05$) grain yield increase. Application of 10 kg P₂O₅ ha⁻¹ also
 185 | had a significant effect compared to the control plots.

Formatted: Font: Italic

186 | **Table - 3:-** Main effects of nitrogen and phosphorus fertilizer levels on yields of upland-rice

Treatments	GYLD	SYLD	TSW	HI
Nitrogen Levels (kg N ha⁻¹)				
N0	2636.30 ^D	4029.80 ^D	35.38 ^A	39.63 ^B
N1	3263.30 ^C	4646.40 ^C	35.91 ^A	41.15 ^A
N2	3852.60 ^B	5520.60 ^B	35.37 ^A	41.37 ^A
N3	4306.30 ^A	6198.50 ^A	35.35 ^A	40.83 ^{AB}
LSD	389.37	575.12	NS	NS
Phosphorus levels (kg P₂O₅ ha⁻¹)				
P0	2851.90 ^C	4385.00 ^B	35.50 ^A	39.76 ^B
P1	3320.00 ^B	4807.20 ^B	35.13 ^A	40.95 ^{AB}
P2	3815.70 ^A	5483.60 ^A	36.17 ^A	40.71 ^{AB}
P3	4070.80 ^A	5719.60 ^A	35.21 ^A	41.54 ^A
LSD	389.37	575.12	NS	NS
N*P	NS	NS	NS	NS
CV %	27.5	27.99	9.02	8.87

Formatted: Left, Space After: 0 pt, Line spacing: single

Formatted: Left, Space After: 0 pt, Line spacing: single

187 | Means with the same letter are not significantly different ($P \leq 0.05$). GYLD= Grain Yield (kg ha⁻¹); SYLD= Straw
 188 | yield (kg ha⁻¹); TSW= Thousand Seed Weight (g); HI= Harvest Index (%)

Formatted: Font: Italic

Similar to grain yield, straw yield had similar increasing trend for both main effects. The results showed that application of 23, 46 and 69 kg N ha⁻¹ produced highest straw yield, and it was significantly higher by 15%, 37% and 54% over control. A significantly higher straw yield was also obtained in plots received 30 kg P₂O₅P fertilizer ha⁻¹ than other treatments; on par with plots treated with 20 kg P fertilizer ha⁻¹. Increments in grain and straw yields of rice due to main effect nitrogen and phosphorus fertilizers were reported in different regions of the country [6], [10], [12] and elsewhere [13]. Similarly, [22] and [8] also reported a significant yield difference for Teff and Wheat respectively. The thousand grain weight and harvest index of rice were not significantly affected by interactions as well as the main effects of N and P fertilizers in this experiments.

The response curve revealed that optimum grain yield of rice to the application of N and P fertilizers were not found. Application of N and P fertilizers showed increasing trend (Figure 1). Application of N and P at a rate of 69 kg N ha⁻¹ and 30 kg P₂O₅P ha⁻¹ were not enough to boost up the production and productivity of rice in the district and application of higher rates of N and P should be tried.




Figure - 1:- Response curve of rice to Nitrogen-nitrogen and phosphorus fertilizer

levels

Comment [J4]: Give yield in t ha⁻¹

Agronomic efficiency of rice to the application of N and P

N (kg ha ⁻¹) Rate	AEN (kg kg ⁻¹)	P (kg ha ⁻¹) Rate	AEP (kg kg ⁻¹)
0		0	-
23	27.26	10	46.81
46	26.44	20	48.19
69	17.62	30	40.63

208 Agronomic efficiency is the amount of additional yield obtained for each additional kg of
 209 nutrient applied [21]. Agronomic efficiency of rice showed decreasing trend with increasing rate
 210 of N fertilizer application (Table 4). Therefore, application of one kg of N caused increase in
 211 grain yield of rice by 27.26 kg from plots treated with 23 kg N ha⁻¹. Even though there were
 212 some inconsistent trends, agronomic efficiency of rice to application of P decreased with P rates.
 213 The highest agronomic efficiency was recorded on plots treated with 20 kg P_2O_5 ha⁻¹. The result
 214 from site was nearly in line with the findings of [9], where AE decreases with N and P rates.

215 **Table - 4:-** Agronomic efficiency of rice to the application of N and P

216 AEN (kg kg⁻¹) = Agronomic Efficiency of rice to Nitrogen fertilizer and AEP (kg kg⁻¹) = Agronomic Efficiency of
 217 rice to Phosphorus fertilizer

218 Conclusion

219 The growth and yield of upland rice were enhanced significantly due to application of nitrogen
 220 and phosphorus fertilization on vertisols of Tselemti area. Plant height, number of spikelet and
 221 total number of seeds per panicle were affected by the interaction of higher doses, while the
 222 panicle length, unfertilized seeds, grain yield as well as straw yield by the main effect of nitrogen
 223 and phosphorus fertilizers. Thus, according to the results, an optimum grain and straw yield of
 224 upland rice can be achieved in the district by applying nitrogen and phosphorus fertilizer at a rate
 225 of 69 kg N ha⁻¹ and 30 kg P_2O_5 ha⁻¹, respectively. However, these application rates were not
 226 enough to boost up the production and productivity of rice in the district and application of
 227 higher rates of N and P should be tried. Additionally, higher agronomic efficiency of rice was
 228 also obtained from fertilization of 23 kg N ha⁻¹ and 20 kg P_2O_5 ha⁻¹.

229 References

- 230 [1] MoA, *National Rice Research and Development Strategy of Ethiopia*. Addis Ababa,
231 Ethiopia, 2010.
- 232 [2] MoA, "Towards Implementation of National Rice Development Strategies of Ethiopia,"
233 Addis Ababa, 2011.
- 234 [3] CSA, "REPORT ON AREA AND PRODUCTION OF MAJOR Crops," Addis Ababa,
235 2014.
- 236 [4] CSA, "Statistical REPORT ON AREA AND PRODUCTION OF CROPS (2007- 2016),"
237 Addis Ababa, 2016.
- 238 [5] FAO, *FAO Statistical Yearbook*. Rome, Italy, 2013.
- 239 [6] H. Gebrekidan and M. Seyoum, "Effects of Mineral N and P Fertilizers on Yield and
240 Yield Components of Flooded Lowland Rice on Vertisols of Fogera Plain , Ethiopia," *J.*
241 *Agric. Rural Dev. Trop. Subtrop.*, vol. 107, no. 2, pp. 161–176, 2006.
- 242 [7] M. Tekalign, I. Haque, and C. S. Kamara, "Phosphorus status of some Ethiopian highland
243 Vertisols," in *Management of vertisols in Sub-Saharan Africa*, 1998.
- 244 [8] F. Mirutse, M. Haile, F. Kebede, A. Tsegay, and C. Yamoah, "Response of Teff [
245 *Eragrostis* (teff) Trotter] to Phosphorus and Nitrogen on Vertisol at North Ethiopia," *J.*
246 *Drylands*, vol. 2, no. 1, pp. 8–14, 2009.
- 247 [9] B. Haileselassie, D. Habte, M. Haileselassie, and G. Gebremeskel, "Effects of mineral
248 nitrogen and phosphorus fertilizers on yield and nutrient utilization of bread wheat (
249 *Triticum aestivum*) on the sandy soils of Hawzen District , Northern Ethiopia," vol. 3, no.
250 3, pp. 189–198, 2014.
- 251 [10] B. A. Fayisa and G. D. Welbira, "Influence of Phosphorous and Nitrogen Fertilizer Rate
252 on Grain Yield of Rice at Kamashi Zone of Benshal-gul Gumuz Region, Ethiopia," *J.*
253 *World Econ. Res.*, vol. 5, no. 2, pp. 8–14, 2016.
- 254 [11] G. Dereje, B. Walegn, A. Giddisa, H. Solomon, A. Hagos, A. Dabi, R. Dibaba, and D.
255 Alemu, "Participatory Evaluation and Determination of N and P Fertilizer Application
256 Rate on Yield and Yield Components of Upland Rice (NERICA-4) at Bambasi District,
257 Benishangul-Gumuz Regional State," *Adv. Crop Sci. Technol.*, vol. 5, no. 4, 2017.
- 258 [12] S. Nesgea, H. Gebrekidan, J. J. Sharma, and T. Berhe, "East African Journal of Sciences (
259 2012) Effects of Nitrogen and Phosphorus Fertilizer Application on Yield Attributes ,
260 Grain Yield and Quality of Rain Fed Rice (NERICA-3) in Gambella , Southwestern
261 Ethiopia," vol. 6, pp. 91–104, 2012.
- 262 [13] V. A. Ochwoh and P. Abulo, "Influence of nitrogen and phosphorus fertilizer application
263 on grain yield of upland rice in Eastern Uganda," *African J. Crop Sci.*, vol. 3, no. 9, pp.
264 230–233, 2015.
- 265 [14] H. Hagos, E. Ndemo, and J. Yosuf, "Factors affecting adoption of upland rice in Tselemti
266 district , northern Ethiopia," *Agric. Food Secur.*, vol. 7, no. 59, pp. 1–9, 2018.
- 267 [15] S. Sahlemedhin and B. Taye, "Procedures for Soil and Plant Analysis," Addis Ababa,
268 2000.
- 269 [16] P. Hazelton and B. Murphy, *Interpreting Soil Test Results: What do all the numbers*
270 *mean?* Collingwood VIC 3066, Australia: CSIRO Publishing, 2007.
- 271 [17] T. Tekalign, "Soil, plant, water, fertilizer, animal manure and compost analysis," Addis
272 Ababa, 1991.
- 273 [18] D. Berhanu, "The physical criteria and their rating proposed for land evaluation in the

- 274 highland region of Ethiopia,” Addis Ababa, 1980.
- 275 [19] A. Cottenie, “Soil and plant testing as a basis of fertilizer recommendations,” *FAO soil*
- 276 *Bull.*, vol. 38, no. 2, 1980.
- 277 [20] S. Yoseftabar, “Effect of Nitrogen and Phosphorus Fertilizer on Growth and Yield Rice (
- 278 *Oryza Sativa L*),” *Int. J. Agron. Plant Prod.*, vol. 3, no. 12, pp. 579–584, 2012.
- 279 [21] K. Mengel and E. A. Kirkby, *Principles of Plant Nutrition Edited by and*, 5th Editio.
- 280 Kluwer Academic Publishers, 2001.
- 281 [22] F. Khan, S. Khan, S. Fahad, S. Faisal, S. Hussain, and S. Ali, “Effect of Different Levels
- 282 of Nitrogen and Phosphorus on the Phenology and Yield of Maize Varieties,” *Am. J. Plant*
- 283 *Sci.*, vol. 5, no. July, pp. 2582–2590, 2014.
- 284