

SCIENCEDOMAIN international

www.sciencedomain.org

SDI Review Form 1.6

| Journal Name: | Annual Research & Review in Biology |
|--------------------------|---|
| Manuscript Number: | Ms_ARRB_34737 |
| Title of the Manuscript: | PREVALENCE OF GEOHELMINTH PARASITES OF HORSES IN DUTSINMA METROPOLIS, KATSINA STATE NIGERIA |
| Type of the Article | |

General guideline for Peer Review process:

This journal's peer review policy states that <u>NO</u> manuscript should be rejected only on the basis of '<u>lack of Novelty'</u>, provided the manuscript is scientifically robust and technically sound.

To know the complete guideline for Peer Review process, reviewers are requested to visit this link:

(http://www.sciencedomain.org/page.php?id=sdi-general-editorial-policy#Peer-Review-Guideline)

SCIENCEDOMAIN international



SDI Review Form 1.6

PART 1: Review Comments

| | Reviewer's comment | Author's comment (if agreed with reviewer, correct the manuscript and highlight that part in the manuscript. It is mandatory that authors should write his/her feedback here) |
|------------------------------|---|--|
| Compulsory REVISION comments | The manuscript Ms_ARRB_34737 has been reviewed. The study deals with the coprological prevalence of geohelminth in horses in an area of Nigeria. The number of horses examined (n=48) is too small. Statistical analysis section is lacking in Materials and Methods. However, the authors report the finding of Ascaris lumbricoides eggs in horse faeces. This is the first time I read something like that. Ascaris lumbricoides is well known to be a parasite affecting human beings not horses. Moreover, although the authors report the finding of A. lumbricoides eggs, they show a picture of Ascaris larvae (?) where only plant debris can be seen. Similarly, it is also amazing that the authors report the finding of Strongyloides stercoralis larvae in horse faeces. Strongyloides stercoralis is also well known to be a parasite affecting human beings not horses. Although the authors report the finding of S. stercoralis larvae in horse faeces, they show a picture where only a strongyle-like egg can be seen. Strongyloides species other than S. stercoralis parasitize horses but they release embryonated eggs not larvae in faeces. The authors seem to confuse gastrointestinal strongyles of horses and Strongyloides species. Gastrointestinal strongyles of horses and Strongyloides species are completely different parasites with different life cycle and different host range, even if the name sounds similar. | |



SCIENCEDOMAIN international

www.sciencedomain.org

SDI Review Form 1.6

| Minor REVISION comments | |
|----------------------------|--|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| Optional/General comments | |
| optional/octional comments | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |

Reviewer Details:

| Name: | Roberto Amerigo Papini |
|----------------------------------|---------------------------|
| Department, University & Country | Università di Pisa, Italy |