


## Original Research Article


# Effect of Shea Nut Shell Biochar on Root Knot Nematodes (*Meloidogyne* spp.) of Tomato (*Solanum lycopersicum* L.)

### ABSTRACT

Effect of shea nut shell biochar on root knot nematodes and performance of tomato was investigated under nematode infested and inoculated soils. Tomato variety (Petomech-GH) was planted in potting medium of soil to biochar ratio of one part of biochar is to one part of soil (1B1S), one part of biochar is to two parts of soil (1B2S), two parts of biochar is to one part of soil (2B1S), and no biochar application (control). Nematode infested soil was amended with biochar as well as steam sterilized soil amended with biochar inoculated with 1000  The result indicated that, biochar increased the pH of the soil, lessened the adverse effects of nematodes, resulting in decline in galling and improvement in growth and yield of tomato. Increased biochar concentration resulted in decreased nematode gall formation on the roots of the tomato plant. Biochar amended soils resulted in lower egg masses. Increased biochar concentration resulted in decreased performance of tomato plant. Tomato plants treated with low biochar concentrations (1B2S and 1B1S) produced higher fruit numbers and weights, and plant biomass.

*Keywords: Biochar; shea nut shell; root knot nematode; tomato.*

### 1. INTRODUCTION

Tomato (*Solanum lycopersicum* L.) is one of the most popular vegetable consumed in almost every Ghanaian household [1]. It is an important component of balanced diet of most Ghanaians that provide vitamin A and C, lycopene which serves as antioxidant and can help reduce the risk of cardiac diseases and some types of cancer [2]. Tomato production in Ghana has been significantly affected by the incidence of pest and diseases [3] notably among which are root knot nematodes [4, 5]. Crop damages more than 27% ~~take place~~ in tomato [6] and in excess of \$100 billion loss globally [7]. At the Bontanga irrigation zone in the northern region of Ghana, total crop loss of tomato occurs and currently, most farmers do not cultivate tomato in this area [1]. Soil fumigants and chemical nematicides are used in controlling nematodes. These are **however** expensive and poses threats to environment and human health resulting in its withdrawal. Several reports indicated the use of botanicals, aqueous and crude plant extracts for nematode control, which contain minimum bioactive concentration against RKN [8, 9, 10]. Biochar  most agro by products has now been directed to manage nematodes. Biochar addition to soil increases the pH of the soil to become alkaline [11]. The

29 increased pH results in a sharp decrease in nematode population when the soil becomes alkaline  
30 [12]. There is one published report that biochar soil amendment at the concentration of 1.2% delays  
31 the development of root knot nematode [13].

32 The objective of the present study, therefore, was to evaluate the effect of biochar on root knot  
33 nematode and the performance of tomato.

34

## 35 **2. MATERIALS AND METHODS**

### 36 **2.1 Experimental Site**

37 The study was carried out at the plant house of the University for Development Studies (UDS),  
38 Nyankpala campus which lies within latitude 9° 25' 41" and longitude 0° 58'42" W. The entire  
39 experiment was conducted from September to December, 2017.

### 40 **2.2 Source of Study Materials**

41 Tomato seeds (Petomech GH) were obtained from the local farmers in Nyankpala. The shea nut shell  
42 used to make the biochar was sourced from Cheyohi, a suburb of UDS Nyankpala campus.  
43 Nematode infested soil sample was collected from Bontanga irrigation farm in the Kumbungu district  
44 of the northern region of Ghana.

### 45 **2.3 Biochar Preparation**

46 Shea nut shells were placed in a barrel with holes under and a chimney on top which serves as a  
47 pyrolyzer. Dried leaves were lighted on top of the shea nut shell for a few minutes and covered with a  
48 chimney to allow charring or incomplete burning of the shells which will eventually form biochar. It is a  
49 slow process which takes some time but very efficient when done in small quantities.

### 50 **2.4 Experimental Approach**

51 Two methods of nematode study were employed, using nematode infested and inoculated soils.

### 52 **2.5 Soil Sampling, Extraction and Identification of Nematodes**

53 Soil was sampled from each plot at the start of the trial and at harvest with a 5 cm diameter soil  
54 auger to a depth of 20 cm. Initial soil nematodes were extracted from 200 cm<sup>3</sup> of soil before  
55 planting using the modified Baermann tray method [14]. Various nematode species were  
56 identified under a compound microscope at a magnification of 100x. Nematode species were  
57 identified [15] morphologically.


## 58 **2.6 Soil Sterilization**

59 Soil for inoculated experiment was sterilized using the steam barrel sterilization method. Gravels were  
60 removed from sandy loam soil by sieving, which was then packed into a jute sack. Three stones were  
61 laid in a triangular form above the ground level to provide space for fire wood. Water was poured into  
62 a tank about one quarter. Tripod wooden slaps were placed little above the water surface to provide  
63 room for vapor to form. The soil was then placed on this wooding slaps and the tank covered with  
64 polythene. Fire was set under the tank and the heat produced was used to generate steam below the  
65 soil in the tank which was then allowed to stand for 6 hours.

## 66 **2.7 Nursing of Seeds and Transplanting**

67 Tomato seeds were sown in steam sterilized soil placed in a wooden box measuring 1.0 m by 0.6 m.  
68 Cultural practices such as watering and shading was done to ensure proper germination. The most  
69 uniform seedlings were transplanted three weeks after emergence.

## 70 **2.8 Application of Nematode Inoculum Level to Potted Tomato Seedlings**

71 The potted seedlings were inoculated with 3 ml of the nematode solution per pot two weeks after  
72 transplanting (approximately 100 ). Three holes were made in a triangular form 2 cm equidistant  
73 from the base of each plant. The nematode solution was homogenized by gentle shaking the test  
74 tubes containing the nematode solution and then dispensed into the holes.

## 75 **2.9 Experimental Design and Treatments**

76 The experiments were laid out in completely randomized design with five replications. Treatment was  
77 prepared into a 2 L size pot. The pots were filled with 1.6 L of the infested soil-biochar combination  
78 and sterilized soil-biochar combination in different proportions (v/v). The control was without biochar.  
79 Watering was done early mornings or evenings. Too much watering was avoided to prevent water  
80 logging. Detailed treatment descriptions (v/v) were as follows: one part of biochar is to one part of soil  
81 (1B1S); one part of biochar is to two parts of soil (1B2S); two parts of biochar is to one part of soil  
82 (2B1S); no biochar application (control)

## 83 **2.10 Data Collection and Statistical Analysis**

84 Plant growth parameters such as plant height, number of leaves and root weight were taken at two  
85 weeks interval after transplanting. Yield characteristics such as shoot weight, number of fruits, fruit  
86 weight and plant biomass were also taken. Nematode induced parameters such as root galling, egg  
87 mass indices and final nematode population were also taken.

88 Data collected were subjected to analysis of variance (ANOVA) using Genstat (18<sup>th</sup> Edition) statistical  
89 package. Treatment means was separated using least significant difference (LSD) at 5% level of  
90 significance.

### 91 **3. RESULTS AND DISCUSSION**

92 No significant difference on the plant height of tomato was observed among the biochar treatments  
93 which were significantly different from the control (Table 1). However, it was observed that, as  
94 concentration of the biochar increased, the height of tomato plant decreased. Lower mean height was  
95 observed in the highest biochar concentration (2B1S). Similar observation was made by Howard [16]  
96 in corn and soybean, where he reported reduced growth in higher biochar weights investigated and  
97 suggested that, increment in alkalinity of the soil, the holding of too many nutrients, potential toxic ions  
98 and microbes upon too much biochar addition may have negative effect on plant growth. Grabber et  
99 al. [17] similarly reported enhanced plant height of tomato following biochar application.

100 The reduction of plant height under control condition was due to root knot nematode infection.  
101 Sharma and Sharma [18] reported significant reduction in plant height of tomato due to KN infection  
102 (1000 J2).

103 The effect of biochar on the number of leaves was only significant at 2WAP and 4WAP (Table 1). At  
104 2WAP, 1B2S treatment recorded the highest average leaf number while 2B1S treatment recorded the  
105 lowest. This might be attributed to the fact that, at 2WAP, root knot nematode may have penetrated  
106 the roots of tomato but may have not cause significant infection. At 4WAP, similar observation was  
107 made but in this case, the average leave number for 1B1S treatment was higher than the control  
108 whereas 2B1S treatment recorded the lowest. It was observed that, as the concentration of the  
109 biochar increased, leave number decreased.

110 Root weight generally differs based on the concentration of biochar with 1B2S treatment recording  
111 higher significant mean values followed by 1B1S, 2B1S and the control respectively (Table 1). The  
112 root weight of the control plant was significantly low because of the lack of formation of lateral roots  
113 due to root knot nematode infection. This agree with the findings of Sharma and Sharma [18], whose  
114 report indicated significant reduction in root weight and root length of tomato as a result of root knot  
115 nematode infection.

116

117

118

119 **Table 1. Effect of biochar concentrations on growth characteristics of tomato**

Treatment	Plant height			Number of leaves			Root weight		
	2WAP	4WAP	6WAP	2WAP	4WAP	6WAP	2WAP	4WAP	6WAP
1B1S	23.80	33.52	48.30	5.20 <sup>b</sup>	8.20 <sup>bc</sup>	13.80 <sup>a</sup>	0.96	1.18	1.98
1B2S	25.12	36.10	52.34	6.00 <sup>bc</sup>	10.00 <sup>bc</sup>	17.80 <sup>a</sup>	1.27	1.88	2.47
2B1S	22.94	26.58	39.50	4.40 <sup>a</sup>	4.80 <sup>a</sup>	7.50 <sup>a</sup>	0.92	1.26	1.60
Control	25.54	31.54	39.25	5.60 <sup>bc</sup>	7.60 <sup>b</sup>	8.00 <sup>a</sup>	1.26	1.18	1.03
LSD $\alpha=0.05$	7.11	10.05	18.86	0.43	2.11	11.69	0.85	0.86	1.32
<i>P</i> values	0.56	0.05	0.14	< 0.01	< 0.01	0.05	0.51	0.05	0.06

120 *Means followed by the same letter(s) in a column are not significantly different ( $P > .05$ ).*121 **3.2 Yield and Yield Parameters**

122 There was significant effect of biochar on the number of fruit, fruit weight and plant biomass of tomato  
123 (Table 2). This varied according to the biochar treated with 1B2S recording the highest average mean  
124 value followed by 1B1S and control respectively. 2B1S treatment produced no fruits and at the same  
125 time recorded the lowest dry plant biomass which may be due to the high biochar concentration. This  
126 agree with the findings of Grabber et al. [17] whose reports indicated that, biochar contain chemicals  
127 most of which are phytotoxic or biocidal at high concentration and therefore may affect plant growth.  
128 1B2S recorded the highest increment in plant biomass and fruit weight followed by 1B1S with control  
129 recording the least. Grabber et al. [17] reported significant improvement in plant growth in low biochar  
130 concentration. Hossain et al. [19] also reported improved growth and productivity of cherry tomato at  
131 10t/ha biochar application. The observed low biomass of control was due to root knot nematode  
132 infection. Sharma and Sharma [18] reported reduced growth as a result of root knot nematode  
133 infection in tomato. Similarly, Maleita et al. [20] reported stunted growth and reduction in yield on root  
134 knot nematode heavily infested fields. Moreover, application of 1000 J2 per plant significantly reduced  
135 growth and yield in a trial by Haider et al. [21] using French bean and pea.

136

137 **Table 2. Effect of biochar concentrations on yield and yield parameters of tomato**

Treatment	Shoot weight (g)			Mean plant biomass (g)			Fruit number	Fruit weight (g)
	2WAP	4WAP	6WAP	2WAP	4WAP	6WAP		
1B1S	1.71 <sup>a</sup>	4.07 <sup>ab</sup>	6.63 <sup>a</sup>	0.34 <sup>a</sup>	0.86 <sup>ab</sup>	2.96 <sup>a</sup>	3.00 <sup>a</sup>	37.60 <sup>b</sup>
1B2S	2.27 <sup>a</sup>	4.50 <sup>ab</sup>	9.20 <sup>a</sup>	0.42 <sup>a</sup>	1.19 <sup>ab</sup>	5.28 <sup>b</sup>	7.00 <sup>b</sup>	170.00 <sup>c</sup>

2B1S	1.88 <sup>a</sup>	2.79 <sup>a</sup>	4.43 <sup>a</sup>	0.34 <sup>a</sup>	0.81 <sup>a</sup>	1.91 <sup>a</sup>	0.00 <sup>a</sup>	0.00 <sup>a</sup>
Control	1.60 <sup>a</sup>	3.10 <sup>ab</sup>	3.76 <sup>a</sup>	0.37 <sup>a</sup>	1.46 <sup>ab</sup>	1.92 <sup>a</sup>	2.00 <sup>a</sup>	25.00 <sup>b</sup>
LSD $\alpha=0.05$	1.39	1.71	7.95	0.20	0.65	2.70	4.07	17.78
<i>P</i> values	0.44	0.02	0.20	0.55	0.02	<0.01	<0.01	<0.01

138 *Means followed by the same letter(s) in a column are not significantly different (P > .05).*

139 **3.3 Root knot Nematode Population and Reproduction Factor**

140 Final nematodes population and reproduction factor is an indication of nematode multiplication.  
 141 Biochar treatment resulted in significant reduction in final nematode population over the control at  
 142 termination of both experiments (Table 3). 2B1S recorded the highest reduction in final nematode  
 143 population which is significantly different from 1B1S and 1B2S. The control however showed a  
 144 significant increase in final nematode population at the end of both experiment ( $P < .05$ ).

145 Nematode reproduction factor, as indicated in Table 2, also showed significant differences among the  
 146 treatments with 2B1S recording the lowest reproduction factor less than 1 for both experiment,  
 147 followed by 1B1S and 1B2S respectively. The control recorded the highest reproduction factor which  
 148 was greater than 1 for both experiment. This suggested that, root knot nematode may not multiply in  
 149 biochar amended soils. It is generally observed that, nematode population and reproduction factor  
 150 decreased as the concentration of biochar in the medium increased showing the nematicidal potential  
 151 of biochar against RKN. Biochar soil amendments was targeted to highly weathered and acidic soil  
 152 because biochar has been reported to increase soil pH and moisture content [22, 11]. Kung et al. [12]  
 153 reported a sharp decrease in nematode population when the pH of the soil became alkaline. 2B1S  
 154 recorded the highest alkaline pH of 8.62, followed by 1B1S with a pH of 7.46 and 1B2S with a pH of  
 155 7.12 at the end of the experiment. The control soil was however acidic with a pH of 6.20. Since  
 156 biochar amended soil becomes alkaline at the end of both experiments, the reduction in final  
 157 nematode population and decreased reproduction factor in biochar amended soil may be attributed to  
 158 increased pH of the medium.

159 **Table 3. Effect of biochar concentration on final nematode population and reproduction factor**

Treatment	Final nematode population per ml		Reproduction factor	
	Infested soil	Inoculated soil	Infested soil	Inoculated soil
1B:1S	21.00 <sup>b</sup>	23.00 <sup>b</sup>	0.66 <sup>b</sup>	0.72 <sup>ab</sup>
1B:2S	27.00 <sup>c</sup>	32.00 <sup>c</sup>	0.84 <sup>b</sup>	1.00 <sup>b</sup>
2B:1S	0.00 <sup>a</sup>	13.00 <sup>a</sup>	0.00 <sup>a</sup>	0.41 <sup>a</sup>
Control	47.00 <sup>d</sup>	61.00 <sup>d</sup>	1.47 <sup>c</sup>	1.91 <sup>c</sup>
LSD $\alpha=0.05$	2.14	2.33	0.18	0.56

*P* values <0.001 <0.001 <0.01 <0.01

160 Means followed by the same letter(s) in a column are not significantly different ( $P > .05$ ).

### 161 3.4 Root Galling

162 Root knot nematode infection is manifested by the development of galls or giant cells on the root  
 163 accompanied by stunted growth, chlorosis and loss of energy by the plant [23]. Biochar lessened the  
 164 adverse effects of nematodes, resulting in decline in galling and an improvement in the growth and  
 165 yield of the tomato, but the effect differs based on the treatment applied and parameters measured.  
 166 Application of biochar treatment significantly reduced the formation of galls on the root of tomato as  
 167 shown in Table 4. The number of galls or knots varied with the concentration of the biochar treatment.  
 168 Results revealed that, extent of gall formation on the roots were significantly lower in higher biochar  
 169 treated medium with 2B1S recording the lowest root galling followed by 1B1S and 1B2S respectively.  
 170 It may be observed that, as the biochar concentration increased, the extent of gall formation on the  
 171 roots of tomato decreased. Root galling was not observed during the first 2WAP, at 4WAP, significant  
 172 galling occurred on the roots which increased at 6WAP. The absence of galls during the first 2WAP  
 173 may be due to the fact that most RKN has a life cycle of at least three (3) weeks [24]. The root knot  
 174 nematodes may have penetrated the roots but may have not reproduced to establish permanent  
 175 feeding sites in the roots which lead to the formation of galls.

176 Moreover, the control recorded higher number of root galls, where most of the plants showed  
 177 symptoms of wilting during the day and most died before maturity. This agrees with the findings of  
 178 Mitkowski and Abawi [25] who reported wilting and stunted growth in lettuce as a result of RKN  
 179 infection. It is observed that, the extent of gall formation on the root correlated with egg mass indices  
 180 analyzed. Treatments that recorded higher root gall index had higher egg masses (Table 4). Biochar  
 181 amended soils had lower egg masses in which no significant differences occur among the three  
 182 biochar concentrations but all were significantly different from the control. Hence biochar may have  
 183 the potential to manage gall formation on the roots of tomato.

184

185 **Table 4. Effect of biochar concentration on root gall formation and egg masses**

Treatment	Root gall index				Egg mass index	
	Infested soil		Inoculated soil		Infested soil	Inoculated soil
	4WAP	6WAP	4WAP	6WAP		
1B1S	1.80 <sup>b</sup>	2.00 <sup>a</sup>	2.10 <sup>a</sup>	2.60 <sup>b</sup>	1.10 <sup>a</sup>	1.60 <sup>a</sup>
1B2S	2.80 <sup>c</sup>	2.80 <sup>a</sup>	3.00 <sup>a</sup>	3.10 <sup>b</sup>	1.62 <sup>a</sup>	1.81 <sup>a</sup>
2B1S	1.00 <sup>a</sup>	1.00 <sup>a</sup>	1.20 <sup>a</sup>	1.30 <sup>a</sup>	0.00 <sup>a</sup>	0.60 <sup>a</sup>
Control	6.80 <sup>d</sup>	8.00 <sup>b</sup>	6.40 <sup>b</sup>	8.10 <sup>c</sup>	3.67 <sup>b</sup>	3.50 <sup>b</sup>
LSD $\alpha=0.05$	0.77	2.68	2.17	1.57	1.65	1.29

P Value <0.01 <0.01 <0.01 <0.01 <0.01 <0.01

186 *Means followed by the same letter(s) in a column are not significantly different (P > .05).*

187

#### 188 4. CONCLUSION

189 The effectiveness of biochar against root knot nematodes may be confirmed by an increment in shoot  
 190 growth, plant biomass, fruit number and weight which are due to decline in nematode attack as  
 191 indicated by decreased final nematode population in biochar treated soils. It was assessed from the  
 192 study that, root knot nematodes density decreased whilst plant growth parameters were enhanced  
 193 significantly due to biochar application. Biochar increased the pH of the soil to become alkaline at the  
 194 end of the experiment. Soil pH control should be carried out after biochar application to a range that is  
 195 suitable for the growth of tomato.

196

#### 197 REFERENCES

198

- 199 1. Asare-Bediako E, Showemimo FA, Buah JN, Ushawu Y. Tomato production constraint at  
 200 Bolgatanga. *J Appl Sci.* 2007;7(3):459–461.
- 201 2. Campbell JK, Canene-Adams K, Lindshield BL, Boileau TWM, Clinton SK, Erdman J. Tomato  
 202 phytochemicals and prostate cancer risk. *J Nutr.* 2004;134(12):3486S–3492S.
- 203 3. Clotey VA, Karbo N, Gyasi KO. The tomato industry in northern Ghana; production  
 204 constraints and strategies to improve competitiveness. *Afr J Food Agric Nutr Dev.* 2009;  
 205 9(6):1436–1451.
- 206 4. Sikora RA, Fernandez E. Nematodes parasites of vegetables. In: Luc, M., Sikora. A. and  
 207 Bridge, J. eds. *Plant parasitic nematodes in subtropical and tropical agriculture.* Wallingford,  
 208 C.A.B. International. 2005;319–392.
- 209 5. Kankam F, Adomako J. Influence of inoculum levels of root knot nematodes (*Meloidogyne*  
 210 spp.) on tomato (*Solanum lycopersicum* L.). *Asian J Agric Food Sci.* 2014;2(2):171–177.
- 211 6. Kaur DN, Sharma SK, Sultan MS. Effect of different chemicals on root knot nematode in seed  
 212 beds of tomato. *Plant Dis Res.* 2011;26:170–170.
- 213 7. Bird DM, Opperman CH, Williamson VM. Plant infection by root knot nematode. In: Berg, R.  
 214 H. and Taylor, C. G. eds. *Cell biology of plant nematode parasitism.* Berlin, Springer Verlag.  
 215 2008;1–13.
- 216 8. Mashela WP, Dirk De Waele D, Pofu KM. Use of Indigenous Cucumis technologies as  
 217 alternative to chemical nematicides in management of root knot nematodes in low input  
 218 agricultural farming systems: A review. *Scientific Research and Essays.* 2011;6(33):6762–  
 219 6768.
- 220 9. Kankam F, Sowley ENK, Dankwa IN. Management of root knot nematode (*Meloidogyne*



- 221 *incognita*) on cowpea (*Vigna unguiculata* L. Walp.) with oil cakes. Int J Biosci.  
222 2014;5(12):413–419.
- 223 10. Kankam F, Sowley ENK, Mohammed A. Management of root knot nematode (*Meloidogyne*  
224 *spp.*) on okra (*Abelmoschus esculentus* (L.) Moench) with aqueous sesame seed extract. Int  
225 J Agron Agric Res. 2015;6(4):24–31.
- 226 11. Novak JM, Busscher WJ, Laird DL, Ahmedna M, Watts DW, Niandou MAS. Impact of biochar  
227 amendment on fertility of a southeastern coastal plain soil. Soil Sci. 2009;174:105–112.
- 228 12. Kung SP, Gauger R, Kaya KK. Effect of soil temperature, moisture and relative humidity on  
229 entomopathogenic nematodes persistence. J Invertebr-Pathol. 1990;57(2):242–249.
- 230 13. Wen-kun, H., Hong-li, J., Godelieve, G., Jane, D. and Tina, K. Biochar amended potting  
231 medium reduces the susceptibility of rice to root knot nematodes infections. BMC Plant Biol.  
232 2015;15:267.
- 233 14. Whitehead AG, Hemming Jr. A comparison of some quantitative methods of extracting small  
234 vermiform nematodes from soil. Ann Appl Biol. 1965;55:25–38.
- 235 15. CIH (Commonwealth Institute of Helminthology). Description of plant parasitic nematodes.  
236 CAB International, Wallingford, UK.1978.
- 237 16. Howard T. The effect of biochar on the root development of corn and soybeans in Minnesota  
238 soil and sand. International Biochar Initiative. 2011;1–23.
- 239 17. Grabber ER, Yigal E, Eddie C, Yael MH, Beni L. The biochar effect: plant resistance to biotic  
240 stresses. A review. Phytopathol Mediterr. 2011;50:335–349.
- 241 18. Sharma IP, Sharma AK. Effect of initial inoculums levels of *Meloidogyne incognita* J2 on  
242 development and growth of tomato cv. PT-3 under control conditions. Afr J microbiol Res.  
243 2015;9(20):1376–1380.
- 244 19. Hossain MK, Strezov V, Yin-Chan, K, Nelson PF. Agronomic properties of wastewater sludge  
245 biochar and bioavailability of metals in production of cherry tomato (*Lycopersicon*  
246 *esculentum*). Chemosphere. 2010;78(9):1167–1171.
- 247 20. Maleita CMN, Curtis RHC, Powers SJ, Abrantes IMO. Inoculum levels of *Meloidogyne*  
248 *hispanica* and *M. javanica* affect nematode reproduction, and growth of tomato genotypes.  
249 Phytopathol Mediterr. 2012;51(3):566–576.
- 250 21. Haider MG, Dev LK, Nath RP. Comparative pathogenicity of root knot nematode, *Meloidoyne*  
251 *incognita*, on different pulse crops. Ind J Nematol. 2003;33:152–153.
- 252 22. Lehmann J, Gaunt J, Rondon M. Biochar sequestration in terrestrial ecosystems. A review.  
253 Mit Adapt Strat Global Chang. 2006;11:403–427.
- 254 23. Babu AM, Vineet K, Tomy P. Root knot nematode- A hard to kill parasite study. Indian Silk  
255 1999;38:11–12.
- 256 24. Moens T, Yeates GW, Ley P. Use of carbon and energy sources by nematodes. In:  
257 Proceeding of the Fourth International Congress of Nematology. 2004;2:529–545.
- 258 **25.** Mitkowski NA, Abawi GS. Reproductive fitness on lettuce of populations of *Meloidogyne hapla*  
259 from New York State vegetable fields. Nematology. 2003;5(1):77–83.