

Original Research Article

Proximate and phytochemical profile of *Melanthera biflora*

ABSTRACT

The proximate and phytochemical composition of *Melanthera biflora* was investigated, using standard methods. From the obtained results, the leaves had high moisture contents (71.1± 0.2%) and crude fiber (3.91 ± 0.5), while containing moderate protein (70 ± 0.03%) lipid (1.10 ± 4%), ash (2.8 ± 0.2%), total carbohydrate (6.09 ± 0.2%) and caloric value (62.26±0.14 kcal/100g). Eleven Phytochemical families were detected with tannin as the most abundant (27.82%) consisting 100% tannic acid. Thirteen alkaloids (13.65%) were detected consisting mainly of morphine (28.05%), methylmorphine (16.22%), dephnoline (12.02%) biflorin, (20.63%), aromoline (12.61%) homoaromaline (7.79%) and others insignificant amount. Twenty three flavonoid (5.71%) chief among which were quercetin (44.21%), kaemferol (28.94%), dandzein (7.20%), letuolin (10.17%), salvagenin (6.76%), sinensetin 8.20%, and others in insignificant amount. The ten known carotenoids (2.48%), consisting of lutein (40.76%), carotene (17.90%), malvidin 5.63%, zeaxanthin (16.5%), viola-xanthin (9.5%), and others in insignificant amount, were detected. Sixty one terpenoid including linalool (40.98%), germacrene (12.74%), Alpha-terpineal 6.40%, terpinen – 4-01 (5.62%), and Gamma terpine, and others in insignificant amount, were detected. Six phenolic acids (16.26%), consisting of vanilic acid (45.8%), ferulic acid (53.94%), and others in significant amount, were detected. Seven phytosterol (2.25%), consisting of sitosterol (65.3%), savenasterol (14.19%) stigmasterol (12.70%), and others were detected. The leaves had very low hydroxycinnamic acid content (8.93x10⁻⁴%)—content, consisting of eight known compounds of which caffeic acid (71.93%) and p-coumaric acid (27.91%) were the most abundant. They also had a very low alllicins (1.94x10⁻⁴%) content, consisting of daillylthiosulphunate (97.05%), and methyl thiosulphinate (2.6%) and allylthiosulphin and allylthiosulphinate (0.3%). The leaves had a very low content of glycosides consisting of eight known compounds, of which quabain (78.54%) were detected and they include gitogenin (22.04%), diosgenin (20.02), neohegen (20.79%). Their rich contents of nutrients and many bioactive molecules suggest strong nutraceutical potential of these leaves, further suggesting their likely use as a functional food

Keyword: proximate, phytochemical, vegetable

INTRODUCTION

The importance and awareness of nutrition as a prerequisite for good health and longevity has undoubtedly lead to the increase quest for

41 knowledge about the nutritional content of food. Green leafy vegetables occupy
42 an important place among the food crops as they provide adequate amount of
43 vitamins and minerals for human consumption. In addition to their nutritional
44 value, vegetables also contain phytochemicals, which exhibit some protective
45 and disease preventive effect, thus, making them serve a dual function against
46 a number of biochemical, physiological and metabolic disorder. (Aletor and
47 Adeogun 1995), Green leafy vegetables constitute an indispensable constitute
48 of human diet in Africa generally and West Africa in particular (Osagie and
49 Offiong, 1988). Low consumption of green leafy vegetable in the diet is one of
50 the major factor which leads to deficiency of vitamin and iron. Nigeria is
51 blessed with a great natural tropical ~~rainforest~~~~rain forest~~ that is characterized
52 ~~by~~~~with~~ viable soil where vegetables of high nutritional value are grown. This is
53 even more pronounced in Southeastern~~South Eastern~~ Nigeria. There are edible
54 inexpensive leafy vegetables found in this zone (South Eastern Nigeria) whose
55 chemical, nutritional and phytochemical potentials are yet to be adequately
56 studied and utilized. Among this vegetable is “akuwa” (*Melanthera biflora*). The
57 present study therefore is aimed at evaluating the proximate and
58 phytochemical compositions of this tropical leafy vegetable found in South East
59 Nigeria.

60 *Melanthera biflora* is a perenial herbaceous plant which belongs to the
61 family of Asterecae, its common name is beach daisy, it is known among the
62 Igbos as “akwuwa” and “akwuba” among the Efiks in Cross Rivers State
63 Nigeria. It produces a luxiorous edible leaves which is used in making soup.

64 **MATERIALS AND METHODS**

65 **Sample collection**

66 The leaves of *Melanthera biflora* were harvested fresh from Ude plantation in
67 Okon-Aku, in Ohafia Local Government Area of Abia State and was later
68 identified by a taxonomist in the herbarium of the department of plant science,
69 university of Port Harcourt. Dr. Edwin Nwosu.

70 **Sample Preparation**

71 The harvested vegetable leaves destalked, washed with cold running
72 water and divided into two. The first portion was used for proximate analysis
73 while the other portion were dried in an oven at 60°C for 24 hours, after the
74 drying, the leaves were ground into a fine powder using mortar and a pestle
75 and sieved to pass through a 40 mesh sieve and stored in an airtight~~air tight~~
76 container under refrigerated temperature for further use.

77 **Determination of chemical composition**

78 The proximate analysis (carbohydrate, fats, protein, moisture and ash) of
79 the leaves were determined by using AOAC (1995) methods. Carbohydrate was
80 determined by the difference method~~difference method~~ (100- (protein + fat +
81 moisture + ash). The nitrogen value, which is the precursor for protein of a
82 substance, was determined by micro-Kjeldah/method (Guebel et al 1991). The
83 Nitrogen value was converted to protein by multiplying to a factor of 6.25. The
84 moisture and ash were determined using the weight difference method~~weight~~

85 | ~~difference method~~, while determination of crude lipid of the sample was done
86 using soxhlet type and the direct solvent extraction method. Energy value was
87 calculated using Atwater factor method [(9 x fat) + (4xcarbohydrate) +
88 (4xprotein)] as described by Osborne and voogt (1978), and Ihekoronye and
89 Ngoddy (1985). All the proximate values were reported in percentage (AOCS,
90 2000; Okwu and Morah, 2004).

91 **Determination of phytochemicals profile**

92 Phytochemicals were determined using gas chromatography after their
93 individual extractions. (plz write the details of GC and condition)

94 **RESULTS**

95 **Table 1 The proximate composition of *Melanthera biflora* leaves**

Constituent	Composition (%)
Protein (g)	7.00±12
Lipid (g)	1.10±0.16
Crude fiber fibre (g)	3.91±0.01
Ash (g)	2.80±0.14
Moisture (g)	71.10±0.03
Total carbohydrate (g)	6.09±0.12
Total caloric content (kcal)	62.26±0.14

96 Results are means ±S.D of triplicate determination.

97 **Phytochemical profile of *Melanthera biflora* leaves**

98 **Table 2.1 Alkaloid composition of *Melanthera biflora* leaves**

Compounds	Amounts ($\times 10^{-3}$) (mg/100g)	% Composition
Morphine	17882	28.05
Methyl morphine	10340	16.22

Formatted: Highlight

Formatted: Highlight

Papaverine	47.40	0.074
Biflorin	13154	20.63
Narcotine	7.699	0.012
Daphnoline	7664	12.02
Aromoline	8056	12.64
Homoaromoline	4914	7.71
Ambelline	2.309	0.003
6-Hydroxybuphanidine	0.981	0.002
Monocrotalline	9.025	0.001
6-Hydroxy powelline	2.012	0.003
Nitidine	1666	2.613
Total	63751	

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

99 (plz sure these results)

100 **Table 2.2 Flavonoid composition of *Melanthera biflora* leaves**

Compounds	Amount X10 ⁻⁴ (mg/100g)	% Composition
Catechin	0.033	1.219 x 10 ⁻⁵
Resveratrol	1.107	4.15 x 10 ⁻⁴
Apigenin	1880	0.705
Daidzein	19210	7.203
Butein	2.443	9.16 x 10 ⁻⁴
Naringenin	6.454	2.42 x 10 ⁻³
Biochanin	2.65	9.93 x 10 ⁻⁴
Luteolin	27110	10.165
Kaempferol	77190	28.943
(-) - Epicatechin	7.979	2.99 x 10 ⁻³
Salvagenin	18040	6.764
(-) - Epicatechin-3-galleate	5.90	2.212 x 10 ⁻³
Gallocatechin	3.052	1.144 x 10 ⁻³
Quercetin	117920	44.214
Isorhamnetin	36.14	1.355 x 10 ⁻³
Myricetin	5.077	1.904 x 10 ⁻³
Sinensatin	21860	8.19
Kaempferol-3-arabinoside	1.842	0.691
Naringerin	2.841	1.065 x 10 ⁻³
Quercitrin	830.6	0.311
Isoquercetin	415.1	0.156
Orientin	0.409	1.534 x 10 ⁻⁴
Isoorientin	278.5	0.1044
Total	266700	

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

101

102 **Table 2.3 The tannic acid composition and *Melanthera biflora* leaves**

Compound	Amount (mg/100g)
Tannic acid	129.8803

103

104 **Table 2.4 The glycosides composition of *Melanthera biflora* leaves**

Compound	Amount (mg/100g) (X10 ⁻⁶)	% Composition
Kampferol-3-O-rhamnoside	1.490	0.268
Arbutin	6.848	1.234
Salicin	10.64	1.917
Amygdalin	71.85	12.946
Quabain	435.910	78.544
Digitoxin	3.986	0.718
Vitexicarpin	19.962	3.597
Digoxin	0.625	0.43
Costrugenin	3.952	0.712
Total	5.5499	

105

106 **Table 2.5 The phytosterol composition of *Melathera biflora* leaves**

Compound	Retention time (min)	Amount (mg/100g) (X10 ⁻⁵)	% Composition
Cholesterol	19.488	0.0035	0.033
Cholestenol	20.521	6.834	0.64
Ergosterol	21.393	6.877	0.65
Camfesterol	21.954	84190	7.93
Stigmasterol	23.221	134700	12.70
S-Avenasterol	24.018	149900	14.10
Sitosterol	25.260	693200	63.3
Total	-	1062000	-

107

108

109

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

110 **Table 2.6 Allicins composition of *Melanthera biflora* leaves**

Compound	Amount (mg/100g) (X10 ⁻⁶)	% Composition
Diallyl thiosulphinate	8.765	97.05
Metthl allyl thiosulphinate	0.234	2.591
Allyl methyl thiosulphinate	0.031	0.343
Total	9.031	

111

112 **Table 2.7 The carotenoid composition of *Melanthera biflora* leaves**

Compounds	Amount (X10 ⁻³) (mg/100g)	% Composition
Malvidin	651.4	5.627
Carotene	2080	17.968
Lycopene	1.060	0.091
Beta-cryptanxanthin	343.9	2.971
Lutein	4718	40.757
Zeaxanthin	1910	16.500
Anthera-xanthin	3.416	0.030
Asta-xanthin	4.549	0.039
Viola-xanthin	1082	9.347
Neo-xanthin	330.7	2.857
Total	11,576	

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

113

114 **Table 2.8 The saponin composition *Melathera biflora* leaves**

Compounds	Amount (mg/100mg) (X10 ¹)	% Composition
Gitogenin	2.578	22.044
Solagenin	0.0028	0.195
Diosgenin	2.339	20.024
Tigogenin	0.00149	0.042
Neohecogenin	2.429	20.794
Hecogenin	1.764	15.101
Sapogenin	1.659	12.205
Euphol	0.055	0.471
Saponine	0.857	7.337
Total	11.68	

115

116 **Table 2.9 Hydroxycinnamic acid composition of *Melanthera biflora* leaves**

Compounds	Amount (mg/100g) (X10 ⁻⁴)	% Composition
Cinnamic acid	3.278	0.078
Coumarin	0.692	0.017
p-Coumaric acid	11.6	27.914
o-Coumaric acid	2.314	0.056
Caffeic acid	2999	71.918
Sinapinic acid	0.0856	0.002
Chlorogenic acid	0.1937	0.005
Cichoric acid	0.1735	0.004
Total	0.417	

Formatted: Highlight

117

118 **Table 2.10 The phenolic acid composition of *Melanthera biflora* leaves**

Compound	Amount (mg/100g) (X10 ⁻⁴)	% Composition (10)
Vanillic acid	3480	45.85
Ferullic acid	4093	63.94
Syringic acid	1.713	20.24x10 ⁻⁴
Piperic acid	4.410	50.8 x 10 ⁻⁵
Ellagic acid	8.444	1.111 x 10 ⁻⁴
Rosmarinic acid	2.258	2.258
Total	7.590	

Formatted: Highlight

Formatted: Highlight

119

120 **Table 2.11 Terpenes composition of *Melathera biflora* leaves**

Compounds	Amount (Norm. %)
Butanol	0.083
2-Hydroxy-3-butanone	0.366
Butanoic acid	0.116
Sabinene	0.117
2-Methylbutenoic acid	0.095
2-Methylbutanoic acid	0.271
2- Methylbutanoic acid ethyl	0.290
Azulene	0.299
2-methylbutanoic acid ethyl	0.210
Alpha pinene	1.688
Beta pinene	1.788
Benzyl alcohol	0.593

Cis ocimene	3.756
Myrane	0.209
Allo ocimene	0.246
Pinene-2-ol	0.000
Alpha thujene	0.645
Gama terpinene	4.198
2,6-O-dimethyl-1-5 heptanal	0.310
Citral	0.366
Camphor	0.201
Neral	0.519
Geranial	0.405
Iboartemisia	0.245
1,8-Cineole	0.592
Borneol	0.500
Linalool	40.984
Citronellal	0.196
Nerol	0.196
Alpha terpineol	6.395
Terpinen-4-ol	5.620
Citronellol	0.359
Ascaridole	0.468
Linalyl acetate	0.449
Alpha terpinenyl acetate	0.310
Ethyl cinnamate	0.583
Borneol acetate	0.733
Neryl acetate	0.2098
Geranyl acetate	0.311
Beta bisabolene	0.661
Germacrene D	12.735
Gama cadinene	1.690
Beta caryophyllene	0.968
Cyprene	0.143
Beta elemene	0.143
[6]-Shogaol	0.565
Alpha gurgunene	0.469
Alpha copane	0.211
Beta selinene	0.209
Itumulene	0.396
Vacencene	0.310
Caryophyllene oxide	3.856
Alpha selinene	0.491
[6]-Paradol	0.084
Beta selinene	0.248
Aromadendrene	0.370

Gama muurolene	0.314
Aristolone	0.310
Viridiflorol	0.304
Taraxeron	0.325
Lupeol	0.319
Total	100

121

122 **Table 2.12 Percentage composition of group phytochemicals in**
 123 ***Melanthera biflora***

Phytochemicals	Amount (mg/ 100g)	% Composition
Alkaloids	63.75	13.654
Flavonoids	26.670	5.712
Tannic acid	129.88	27.818
Glycosides	5.55 x 10 ⁻⁴	0.001
Terpenoids	100.00	21.418
Phytosterols	10.620	2.275
Allicins	9.031 x10 ⁻⁶	1.937 x 10 ⁻⁶
Carotenoids	11.576	2.480
Saponins	116.81	2.502
Hydroxycinnamic acids	4.170 x 10 ⁻⁴	89.3 x 10 ⁻⁴
Phenolic acids	7.590	16.26
Total	466.898	

124

125 **End Note:** Percentages are based on the weight of the compounds per the
 126 total extract of its family.

127 (plz all labled yellow results must sure them)

128 (the comment on the results must be)

129 **Discussions**

130 The moisture content of *Melanthera biflora* is higher than that of *Talinum*
 131 *triangulare* and *Telferia occidentalis* (Oguntana, 1988), but less than
 132 *Pennisetum purpureum* (Okaraonye and Ikewuchi, 2009). The moisture content
 133 of any food is an index of its water activity (Olutiola et al., 1991) and it is used

134 as a measure of stability and susceptibility to microbial contamination (Uriah
135 and Izuagbe, 1990). The higher moisture content provides for greater activity of
136 water soluble enzymes and coenzymes needed for metabolic activities of leafy
137 | vegetables. The implication of this is that, the leaf will have a higher shelf life
138 than *Pennisetum purpureum*, but a lower one than *Talinum triangulare* and
139 *Telferia occidentalis*. This suggests that the leaves will not be stored for a long
140 time as higher water content enhances microbial action.

141 The crude protein of *Melanthera biflora* is greater than that of *Pennisetum*
142 *purpureum* (Okaraonye and Ikewuchi, 2009), *Amarantus hybridus*, *T.*
143 *occidentalis* and *T. triangulare* (Oguntona, 1998). The leaf protein is rich in
144 essential amino acids. These amino acids serve as an alternative source of
145 energy when carbohydrate availability in the body is impaired. A 100 g of this
146 sample can meet the daily protein requirement of 23-56 g (FAO/WHO/UNU,
147 1991; Chaney, 2006a). Regular uses of plant food rich in protein make an
148 invaluable addition to a diet (Wardlaw, 1999). The ash content of *Melanthera*
149 *biflora* was greater than that reported for *T. occidentalis*, *T. triangulare*
150 (Oguntona, 1998) and *P. purpureum* (Okaraonye and Ikewuchi, 2009), but less
151 than *A. hybridus* (Oguntona, 1998). The ash composition of a food is the
152 amount of minerals substances left after the carbon material must have been
153 burnt of (Onyeike and Osuji, 2013).

154 *Melanthera biflora* leaves contain comparable lipid content to *P.*
155 *purpureum* (Okaraonye and Ikewuchi, 2009) and *A hybridus* (Oguntona, 1998),

156 but greater one than *T. occidentalis*, *T. Triangulare* (Oguntona, 1998) and
157 *Sansevieria liberica* (Ikewuchi et al., 2010).

158 The total carbohydrate content of *Melanthera biflora* was less than those
159 reported for *A. hybridus* (Oguntona, 1998) and *P. tuberregium* sclerotia
160 (Ikewuchi and Ikewuchi, 2009), but more than *P. purpureum* (Okaraonye and
161 Ikewuchi, 2009). A 100 g of the leaves can provide 6-10% of the recommended
162 daily allowance for carbohydrate. *Melanthera biflora* contains higher fibre
163 content than *A. hybridus*, *T. triangulare*, *T. occidentalis* (Oguntona, 1998) and
164 *P. purpureum* (Okaraonye and Ikewuchi, 2009).

165 | Results from epidemiological studies reveal that increased ~~fiber~~fibre
166 | consumption may help in the reduction of certain diseases such as diabetes,
167 | coronary heart diseases, colon cancer, obesity, high blood pressure and various
168 | digestive disorders (Walker 1978; Food and Agriculture Organization;
169 | Eriyamremu and Adamson, 1994; Scientific Advisory Committee on Nutrition,
170 | 2008). Dietary ~~fiber~~fibre has been associated with alternations of the colonic
171 | environment that protect against colorectal diseases. It provides protection by
172 | increasing faecal bulk, which dilates the increased colonic bile concentration
173 | that occurs with a high-fat diet (Dillard and German, 2000). This is one benefit
174 | derivable from the consumption of *Melanthera biflora*.

175 The total caloric content of *Melanthera biflora* was higher than *P.*
176 *purpuerum* (Ikewuchi and Okaraonye, 2009), but less than *P. tuberregium*

177 sclerotia (Ikewuchi and Ikewuchi, 2009). This result shows that *Melanthera* is a
178 good source of nutrient.

179 | **Phytochemical composition of ~~the leaves of~~ *Melanthera Biflora* leaves as**
180 **determined by gas chromatography**

181 The phytochemical screening revealed that *Melanthera Biflora* is rich in
182 tannic acid. Tannic acid is an antioxidant, hepatoprotective, hypocholesteromic
183 and hypoglycemic agent (Liu et al 2005) Tannin is used in the treatment of
184 inflamed or ulcerated tissues. *Melanthera Biflora* is rich in alkaloid, prominent
185 which is morphine used as n analgesic, local anaesthetic and anti-leishmanial
186 | agent (Carrol and starmer 1967). Flavonoid are of a particular importance in
187 the human diet as there are evidence that they act as antioxidants, antiviral
188 and anti-inflamentory agent. (Soetan 2008) and are associated with reduced
189 risk of cancer and cardiovascular diseases. (Middleton et al 2000). Terpenes are
190 used as flavor enhancers in food, frangrances in perfuming and in traditional
191 and alternative medicines such as aromatherapy (Kappers et al 2005). They
192 have anticancer (Dewick 2004) Antimicrobial (Islam et al 2003) and anti-
193 oxidant Dillard and German 2000).

194 The leaves have low saponin, very low glycoside and moderate allicin
195 content. Saponins are reported to have broad range of pharmcological
196 properties (Soetan 2008). Allicin is reported to have an anti-inflammatory,
197 antimicrobial, anti oxidation, anti-thrombotic, ` anti-ulcer, cardioprotective,

198 hypolipidemic, hypotensive and insecticidal properties (Elilat et al, 1995;
199 Elkayam et al 2003).

200 *Melathera Biflora* has moderate phytosterol content. Phytosterol reduce
201 cholesterol levels by competing with cholesterol absorption in the gut of
202 humans (Tilvis and Miethinen 1986). The sample has phenolic acid, which are
203 important for cell structure, signaling and pigmentation (Adyanthaya, 2007).
204 They are known to act as allelochemicals (Yoshioka et al, 2004), protect plant
205 against environmental and biological stress such as high energy radiation,
206 bacterial infection or fungal attacks (Tuzen and Ozdemir, 2003), cold, stress
207 hyperthermia and oxidation stress (Dillard and German, 2000). Thus, their
208 presence in *melanthera biflora* may suggest a likely allelopathic potential of the
209 plant.

210 **CONCLUSION**

211 These results suggest strong nutraceutical potential of this plant and
212 suggest further research in its therapeutic uses in the management and
213 prevention of disease as a result of its rich phytochemical composition.

214 It is a potential pharmaceutical which will help to alleviate some certain
215 kind of diseases and infections such as cancer, cardiovascular diseases, type 2
216 diabetics, cough, hypertension, piles, asthma, malaria etc.

217 **REFERENCES**

218
219 Adyanthaya, I. (2007). Antioxidant response mechanism in apples during post-
220 harvest storage and implications for human health benefits. M.Sc.
221 Thesis, University of Massachusetts, Amherst.
222

223 Aletor, V,A and Adeogun, O.A (1995). Nutrients and anti-nutrients Component
224 of some tropical leafy vegetables. *Food chemistry* 54(4): 375-379
225

226 AOAC International (2006).Official Methods of Analysis of the AOAC (W.
227 Horwitz, Editor).18th edition. AOAC International: Washington DC, USA.
228

229 ACOS American oil Chemist Society (2000) official methods of Analysis 5th edn,
230 Association of official Analytical chemist Washington, DC .USA
231

232 Carrol P.R and Starmer, G.A (1967) studies on the pharmacology
233 *conopharyngine*, an indole alkaloid of the vacanya series. British Journal
234 of *Pharmaceutical and chemotherapy*, 30.173-185
235

236 Chaney, S.G. (2006a). Principles of nutrition I: Macronutrients. In T.M. Devlin
237 (Ed.), *Textbook of biochemistry, with clinical correlation*, 6thedn (pp. 1071-
238 1090). New York: John Wiley and sons.
239

240 Dewick, P.M. (2005). Tumor inhibitors from plants. In W.C. Evans (Ed.).*Trease*
241 *and Evans Pharmacognosy*, 15thedn (pp. 394-406). India: Elsevier.
242

243 Dillard, C.J. and German, J.B. (2000). Phytochemicals: nutraceuticals and
244 human health. *Journal of the Science of Food and Agriculture*, **80**(12):
245 1744 – 1756.
246

247 Eilat, S., Oestraicher, Y., Rabinkov, A., Ohad, D., Mirelman, D., Battler, A.,
248 Eldar, M. and Vered, Z. (1995).Alteration of lipid profile in hyperlipidemic
249 rabbits by allicin, an active constituent of garlic.*Coronary Artery Disease*,
250 **6**: 985–990.
251

252 Elkayam, A., Mirelman, D., Peleg, E., Wilchek, M., Miron, T., Rabinkov, A.,
253 Oron-Herman, M. and Rosenthal, T. (2003).The effects of allicin on
254 weight in fructose-induced hyperinsulinemic, hyperlipidemic,
255 hypertensive rats.*American Journal of Hypertension*, **16**(12): 1053–1056.
256

257 Eriyamremu, G.E. and Adamson, I. (1994). Early changes in energy metabolism
258 in rats exposed to an acute level of deoxycholate and fed a Nigerian-like
259 diet.*Annals of Nutrition and Metabolism*, **38**: 174-183.
260

261 FAO/WHO/UNU (1991). Energy and protein requirements: Report of a Joint
262 FAO/WHO/UNU Expert Consultation. WHO Technical Report Series 724.
263 ISSN: 0512-3054.
264 [Http://www.fao.org/docrep/003/aa040e/AA040E01.htm](http://www.fao.org/docrep/003/aa040e/AA040E01.htm)
265

266 Guebel DV, Nudel BC, Giulietti F (1991). A simple and rapid micro-Kjeldahl
 267 method for total nitrogen analysis. *Biotech. Technol.*, 5(6): 427-430.

268 Ihekoronye, A.I and Ngoddy, P.O.(1985) Integrated Food and Science and
 269 Technology for the Tropics. Macmillian publisher, London P257-264
 270

271 Ikewuchi, C.C. and Ikewuchi, J.C. (2009) chemical profile of pleurotusberregiui
 272 (Fr) Sings Sclerotia. *Pacific journal of science and technology for the*
 273 *tropics.* Macmillian publishers, London p 257-264.
 274

275 Ikewuchi, C.C. and Ikewuchi, J.C. (2009).Chemical profile of *Pleurotu*
 276 *stuberregium* (Fr) Sing's Sclerotia.*Pacific Journal of Science and*
 277 *Technology*, **10**(1): 357-362.

278 Ikewuchi, C.C., Ikewuchi, C.J.,Ayalogu, O.E. and Onyeike, N.E.
 279 (2010).Proximate and phytochemical profile of *Sansevieria liberica*
 280 Gérôme and Labroy. *Journal of Applied Science and Environmental*
 281 *Management*, **14**(2): 103 – 106.
 282

283 Islam, A.K., Ali, M.A., Sayeed, A., Salam, S.M., Islam, A., Rahman, M., Khan,
 284 G.R., and Khatun,S. (2003). An anti-microbial terpenoid from
 285 *Caesalpinia pulcherrima* Swartz: Its characterization, antimicrobial and
 286 cytotoxic activities. *Asian Journal of Plant Science*, **2**: 17-24.
 287

288 Kappers, I.F., Aharoni, A., Van Herpen, T., Luckerhoff, L., Dicke, M.
 289 andBouwmeester, H.J. (2005). Genetic engineering of terpenoid
 290 metabolism attracts bodyguards to Arabidopsis. *Science*, **309**: 2070-
 291 2072.
 292

293 Liu, X., Kim, J.-K., Li, Y., Li, J., Liu, F. and Chen, X. (2005). Tannic acid
 294 stimulates glucose transport and inhibits adipocyte differentiation in
 295 3T3-L1 cells. *Journal of Nutrition*, **135**: 165–171.
 296

297 Middleton, E. Jr., Kandaswami, C. andTheoharides, T.C. (2000). The effects of
 298 plant flavonoids on mammalian cells: implications for inflammation,
 299 heart disease and cancer. *Pharmacology Reviews*, **52**(4): 673-751.
 300

301 Oguntona, T. (1998).Green leafy vegetables.In A.U. Osagie and O.U. Eka (Eds.),
 302 *Nutritional quality of plant foods* (pp. 120-133). Nigeria: Post Harvest
 303 Research Unit, Department of Biochemistry, University of Benin, Benin
 304 City.
 305

306 Okaraonye, C.C. and Ikewuchi, J.C. (2009).Nutritional and antinutritional
 307 components of *Pennisetum purpureum* (Schumach).*Pakistan Journal of*
 308 *Nutrition*, **8**(1): 32-34.
 309

310
311 Olutiola, P.O., Famurewa, O. and Sonntag, H.-G. (1991). *An introduction to*
312 *general microbiology, a practical approach*. Germany: Heidelberger
313 Verlagsanstalt und Druckerei GmbH Heidelberg.
314
315 Onyeike, E.N. and Osuji, J.O.(2003).Techniques of proximate analysis of foods
316 and food products.In E.N. Onyeike and J.O. Osuji (Eds.), *Research*
317 *techniques in biological and chemical sciences* (pp. 226-260).Owerri:
318 Springfield Publishers.
319
320 Osagie, A.U. and Ofiong U.E. (1998) Nutritional quality of plant food. Amik
321 press. Post-harvest Unit. University of Benin, Benin City Nigeria.
322
323 Osborne, D.R. and Voogt, P (1978).*The analysis of nutrients in foods*.London:
324 Academic press, London.
325
326 Scientific Advisory Committee on Nutrition.(2008). Draft SACN position
327 statement on dietary fiber and health and the dietary fiber definition -
328 August
329 2008.SACN/08/20.[http://www.sacn.gov.uk/pdfs/final_draft_sacnstate](http://www.sacn.gov.uk/pdfs/final_draft_sacnstatement_on_dietary_fibre_for_website.pdf)
330 [ment_on_dietary_fibre_for_website.pdf](http://www.sacn.gov.uk/pdfs/final_draft_sacnstatement_on_dietary_fibre_for_website.pdf)
331
332 Soetan, K.O. (2008). Pharmacological and other beneficial effects of anti-
333 nutritional factors in plants - a review.*African Journal of Biotechnology*,
334 **7**: 4713-4721.
335
336 Tilvis, R.S. and Miettinen, T.A. (1986).Serum plant sterols and their relation to
337 cholesterol absorption.*American Journal of Clinical Nutrition*,**43**:92-97.
338
339 Tüzen, M. And Özdemir, M. (2003).Chromatographic determination of phenolic
340 acids in the Snowdrop by HPLC.*Turkish Journal of Chemistry*, **27**: 49 -54.
341
342 Uraih, N. And Izuagbe, Y. (1990).*Public health, food and industrial microbiology*.
343 Nigeria: Uniben Press.
344
345 Walker, A.R.P. (1978). The relationship between bowel cancer and fibre content
346 in the diet. *American Journal of Clinical Nutrition*, 31: S248-S251.
347
348 Wardlaw, G.M. (1999). *Perspectives in nutrition*, 4thedn. London: McGraw-Hill.
349
350 Yoshioka, T., Inokuchi, T., Fujioka, S. and Kimura, Y. (2004).Phenolic
351 compounds and flavonoids as plant growth regulators from fruit and leaf
352 of *Vitexrotundifolia*.*Z. Naturforsch.*, **59c**: 509-514.
353

