

# Evaluation of the nutrients, antinutrients and metals content of five leafy vegetables in Dengi metropolis

## ABSTRACT

**Aim:** To quantify the levels of amino acids, some metals, and phytochemicals/antinutritional factors in leafy vegetables: *Cucurbita pepo*, *Vitex doniana*, *Hibiscus cannabinus*, *Leptadenia hastata*, *Balanites aegyptiaca*.

**Study design:** The research work is descriptive.

**Place and duration of study:** Department of Biochemistry, University of Jos between March 2016 and September 2016.

**Methodology:** 5 samples each of the vegetables were collected at random from different locations in the area of study. Atomic Absorption Spectrophotometry was applied to analyse for metals. Levels of amino acids in samples were determined using the PTH amino acids analyser. Antinutrients were quantified using standard methods. One way ANOVA was used to analyse the data obtained at 95% level of significance.

**Results:** Mean levels of Pb and Cd were  $0.030 \pm 0.01$  ppm and  $0.004 \pm 0.001$  ppm respectively ( $P > 0.05$ ) relative permissible limits. Mean Mg, Mn and Fe content were  $1.084 \pm 0.02$  ppm,  $0.069 \pm 0.01$  ppm and  $1.534 \pm 0.10$  ppm respectively ( $P = .05$ ). Amino acids profile indicated mean values (g/100g proteins) of Glutamate 8.34, Aspartate 8.14, Leucine 8.34, Lysine 4.03, Isoleucine 3.30, Phenylalanine 4.17, Tryptophan 2.25, Valine 4.6, Methionine 1.12, Proline 2.84, Arginine 4.99, Tyrosine 2.75, Histidine 2.23, Cysteine 1.09, Alanine 3.71, Glycine 4.08, Threonine 2.88, Serine 2.99. Mean range of antinutrients were: tannins,  $0.51 \pm 0.13$  % to  $0.60 \pm 0.12$  %, oxalates,  $0.14 \pm 0.14$  % to  $0.60 \pm 0.20$  %, phytates,  $1.70 \pm 0.01$  mg/100g to  $4.10 \pm 0.01$  mg/100g, saponins,  $11.85 \pm 1.85$  % to  $15.13 \pm 1.50$  %, cyanogenic glycosides,  $4.82 \pm 1.30$  % to  $7.59 \pm 1.20$  %, total alkaloids and total flavonoids were  $16.22 \pm 1.61$  % to  $19.37 \pm 1.23$  % and  $9.87 \pm 1.32$  % to  $14.71 \pm 2.30$  % respectively.

**Conclusion:** Although samples analysed contained significant amounts of antinutrients, they are very good sources of amino acids; especially Lysine, Methionine, Leucine, tryptophan which are essential; and mineral elements. Levels of lead and cadmium in the samples were lower than safe limits. These vegetables are good sources of nutrients. Their consumption will replenish nutrients to the cell thereby improving the well being of consumers.

**Key words:** amino acids, antinutrients, nutrients, metals, vegetables

42

**43 INTRODUCTION**

44 Leafy vegetables have been shown to be valuable sources of nutrients (Nesamvuni *et al.*,  
45 2001) with some having medicinal properties (Hilou *et al.*, 2006). These vegetables serve as valuable  
46 sources of nutrients especially in rural areas where they exist in the open country. They might contain  
47 significant levels of trace elements, heavy metals, amino acids as well as antinutrients. Leafy  
48 vegetables can contribute substantially to food security in the rural areas where people's diet is based  
49 on mostly carbohydrates and legumes which are high in calories but deficient in essential  
50 micronutrients and proteins (Yiridoe and Anchirinah, 2005).

51 Heavy metals such as arsenic, cadmium, lead are toxic to cells (Schumacher *et al.*, 1991),  
52 thus it is important to determine their levels in especially in plant-based foods and diets. In general,  
53 information on edibility and therapeutic properties of wild plants is scanty but data on their nutritional  
54 composition and mineral content is negligible (Aloskar *et al.*, 1992). Manganese is an essential trace  
55 element, which plays an important role as a cofactor for many enzyme systems such as hexokinase  
56 and superoxide dismutase. At high level however, it can cause damage to the brain (Aschner and  
57 Aschner, 2005). Magnesium is another nutrient required in the plasma and extra cellular fluid, where it  
58 helps in maintaining osmotic equilibrium. It is required in many enzyme-catalysed reactions,  
59 especially those in which nucleotides participate where the reactive species is the magnesium salt (eg  
60 MgATP<sup>2-</sup>). Lack of Mg is associated with abnormal irritability of muscle and convulsions whereas  
61 excess levels implicated in depression (Aschner and Aschner, 2005).

62 Iron is necessary for the formation of haemoglobin and also plays an important role in oxygen  
63 and electron transfer in human body (Kaya and Incekara, 2000). Also, in the functioning of the central  
64 nervous system as well as oxidation of carbohydrates, proteins and fats (Adeyeye and Otokiti,  
65 1999). Cadmium is a heavy metal that causes both acute and chronic poisoning; adverse effect on  
66 kidney, liver, vascular and immune system (Heyes, 1997). Chronic exposure to chromium may result  
67 in liver, kidney and lung damage (Zayed and Terry, 2003).

68 Lead causes both acute and chronic poisoning with the kidney and liver as primary targets. It inhibits  
69 the catalytic action of  $\delta$ -amino levulinic acid dehydratase (Porphobilinogen Synthase) in haem  
70 biosynthetic pathway; toxic on the vascular and immune system (Heyes, 1997).

71 Antinutritional factors reduce the nutritive values of many plants due to their natural inherence  
72 in the plants. They are capable of eliciting deleterious effect in man and animals (Kubmarawa *et al.*,  
73 2008). Oxalate tends to render calcium unavailable by binding to the calcium ion to form complexes  
74 (Nkafamiya *et al.*, 2009). Phytic acid acts as a strong chelator forming protein and mineral-phytic acid  
75 complexes thereby decreasing protein and mineral bioavailability (Fasusi *et al.*, 2006). Phytate is  
76 associated with nutritional diseases such as rickets and osteomalacia in children and adult,  
77 respectively. Tannins are water soluble phenolic compounds with a molecular weight greater than 500  
78 and with the ability to precipitate proteins from aqueous solution. They occur in all vascular plants.  
79 Tannin binds to proteins making them bio unavailable (Bagepallis *et al.*, 1993). This seeks to quantify  
80 both the nutrients and antinutrients in these plants samples.

81

**82 MATERIALS AND METHODS**83 **Materials**84 **Chemicals and reagents**

85 All reagents and chemicals were of analytical grade from BDH.

**86 Equipment used**

87 OHUAS digital balance, applied biosystems PTH amino acid analyzer and Soxhlet assembly set up.

88

89 Raw samples of the leafy vegetables were collected from different farms in and around Dengi  
90 metropolis. They were destalked, washed and air dried. Samples were pulverised in ceramic mortar  
91 and pestle. This was followed by sieving the samples to obtain fine particles.

**92 Amino acids analysis**

93 The amino acid profile was determined using the method described by Sparkman *et al.*(1958). Where  
94 each of the dried samples were defatted(by refluxing 100g of the air-dried powdered samples with  
95 250ml petroleum ether (600-800) for 4 hours, the resulting residue was then dried and subjected to  
96 aqueous extraction using Soxhlet assembly. Extracts were thereafter evaporated and loaded into the  
97 "applied biosystems PTH amino acid analyzer" which separated and analysed free acidic, neutral and  
98 basic amino acids of the hydrolysate.

**99 Determination of mineral elements content**

100 The minerals content of the different samples was evaluated for Mn, Mg, Fe, Cd, Cr and Pb by dry  
101 ashing of dried powdered sample (5g) in a muffle furnace set at 775°C. The ash obtained was  
102 dissolved in 5 mL of 20% HCl and analysed using the atomic absorption spectrophotometer at their  
103 respective wavelengths of maximum absorption ( $\lambda_{max}$ ) thus: 385nm, 285.5nm, 405nm, 357.8nm,  
104 582nm, 389.6nm for Mn, Mg, Fe, Cd, Cr and Pb in that order.

**105 Determination of antinutrients**

106 Tannins were quantified according to Bainbridge *et al.* (1996), total oxalate quantified applying Day  
107 and Underwood (1986) method. Phytate content was determined by the Reddy and Love method  
108 (1999).

**109 Statistical Analysis**

110 The statistical method employed for all the analysis was one way ANOVA and all results are means of  
111 three determinations ( $\pm$ SD).  $P = .05$  was considered significant.

112

**113 RESULTS AND DISCUSSION**

114 This research sought to assess the amino acids, antinutrients, phytochemicals and metals  
115 composition of five commonly consumed leafy vegetables in Dengi metropolis. Green leafy  
116 vegetables constitute an indispensable constituent of human diet in Africa generally and West Africa  
117 in particular; the varieties of leafy vegetables utilized are diverse, ranging from leaves of annuals and  
118 shrubs to leaves of trees. Leafy vegetables are generally good sources of nutrients, important  
119 protective foods, highly beneficial for the maintenance of health and prevention of diseases as they  
120 contain valuable food ingredients which can be utilized to build up and repair the body. They are  
121 valuable in maintaining alkaline reserve in the body and are valued mainly for their high vitamin,  
122 dietary fibre and mineral content (Olubunmi *et al.*, 2017). The dark green leaves and deep yellow  
123 fruits provide a high amount of carotene, ascorbic acid and micro-minerals which play important roles  
124 in nutrient metabolism and slowing down of degenerative diseases (Olubunmi *et al.*, 2017).  
125

126 Eighteen amino acids were analysed in different proportions in the vegetables. The amino  
127 acid contents are generally high in all samples with the highest, based on dry weight, observed in  
128 *Letadenia hastata* (93.20g/100g protein) others range between 71.41g to 74.62g/100g proteins. Level  
129 of Glutamic Acid was the highest amongst other amino acids in all samples (with an average 11.19  
130 g/100g proteins) followed by Aspartic Acid with an average of 7.92g/100g proteins whereas Cysteine  
131 and Methionine were low with average of 1.37g/100g proteins. All the samples analysed contained  
132 high levels of Glutamate with *Letadenia hastata* have the highest level at 12.11g/g proteins. Levels of  
133 Cysteine and Methionine in *Balanites aegyptiaca* were 0.78g/100g and 0.78g/100g proteins  
134 respectively. The samples contain essential amino acids such as Methionine, Lysine, Leucine,  
135 Isoleucine, Tryptophan, Phenylalanine, Valine and Histidine albeit the levels were lower than the non  
136 essential amino acids. Leucine stimulates muscle strength and growth, regulate blood sugar level by  
137 moderating insulin into the body during and after exercise and can even help prevent depression by  
138 the way it acts on neurotransmitters in the brain (Guoyao, 2010). Lysine is responsible for proper  
139 growth and in the production of carnitine (a nutrient responsible for converting fatty acids into fuel to  
140 lower cholesterol). It also helps the body absorb calcium for further bone strength and also aids in  
141 collagen production. Methionine helps form cartilage in the body through the use of sulphur. Histidine  
142 is involved in transport neurotransmitters to the brain and also helps overall muscle health within each  
143 muscle cells. Valine is needed for optimal muscle growth and repair(Guoyao, 2010).

144 Table 2 shows the mineral content of the vegetables. The permissible limit of iron set by  
145 FAO/WHO (1984) in edible plants is 20 ppm. Iron is necessary for the formation of haemoglobin and  
146 also plays an important role in oxygen and electron transfer in human body (Kaya and Incekara,  
147 2000) and normal functioning of the central nervous system and in the oxidation of carbohydrates,  
148 proteins and fats (Adeyeye and Otokiti, 1999), the highest iron content of 2.223 ppm was found in the  
149 leaves of *Leptadenia hestata* while the leaves of *Balanites aegyptiaca* contain the least iron content of  
150 0.772 ppm, the leaves of *Vitex doniana*, *Hibiscus cannabinus* and *Cucurbita pepo* have significantly  
151 higher iron content of 1.081, 0.916 and 0.831 ppm, respectively. An average culinary preparation  
152 contains about 300g of the fresh leaves of vegetables and this would result in an intake of 30g dry  
153 weight leaves per serving portion. Therefore 1.081, 0.916 and 0.831 ppm in *Vitex doniana*, *Hibiscus*  
154 *cannabinus* and *Cucurbita pepo* respectively, will contribute up to, in mg, 3.2, 2.7 and 2.3/serving  
155 portion of Fe respectively, to the recommended dietary allowance of Fe (10- 15mg/day) (FAO/WHO,  
156 2002).

157 Manganese level, in ppm, was found to be 0.133 in *Leptadenia hastata*, 0.065 in *Hibiscus*  
158 *cannabinus*, 0.041 in *Balanites aegyptiaca*, 0.061 in *Cucurbita pepo* and 0.051 in *Vitex doniana* with  
159 respectively.

160  
161 In table 2, level of Mg, in ppm, was 1.091 in *Cucurbita pepo*, which is the highest compared to others  
162 whose content ranged from 0.915 to 1.080. Mg is required in the plasma and extra cellular fluid,  
163 where it helps in maintaining osmotic equilibrium (Ryan, 1991). It is required in many enzyme –  
164 catalysed reactions, especially those in which nucleotide participate where the reactive species is the  
165 magnesium salt, MgATP<sup>2-</sup>. Deficiency of Mg is associated with abnormal irritability of muscle, and  
166 convulsions. Excess Mg predisposes to depression (Ryan, 1991).

167 Cadmium concentration, in ppm, of the samples ranged from 0.003 in *Hibiscus cannabinus* to  
168 0.004 in *Leptadenia hestata*. These values are below the permissible limit of 0.212 set by FAO/WHO  
169 (1984) in edible plant. In medicinal plants however, the permissible limit by WHO is 0.310. The low  
170 level of Cd in all the samples means they are safe for consumption.

171 As for Pb content of the samples, *Leptadenia hestata* had 0.002 ppm whereas *Balanites*  
 172 *aegyptiaca* contained 0.055 ppm. These levels in the samples are below the permissible level,  
 173 0.43ppm (FAO/WHO 2004).

174 Results of the phytochemical analysis are presented in Table 3. Antinutritional factors have  
 175 been shown to limit the use of many plants due to their ubiquitous occurrence as natural compounds  
 176 capable of eliciting deleterious effect in man and animals (Kubmarawa *et al.*, 2008). The major  
 177 antinutritional factors commonly found in green leafy vegetables are phytic acid, oxalic acid and  
 178 tannins (Osagie and Offiong, 1998). High levels of phytates and oxalates have been shown to inhibit  
 179 the absorption and utilization of minerals such as calcium by animals including man (Grases *et al.*,  
 180 2017). Tannins decrease protein quality by reducing the digestibility and palatability; they interfere  
 181 with absorption of iron and a possible carcinogenic effect (Spalinger *et al.*, 2010).

182 The oxalate content in these vegetables ranged between 0.14mg/100g in *Vitex doniana* to  
 183 0.60mg/100g in *Cucurbita pepo*. These values are below the established toxic level (FAO/WHO,  
 184 2004). The phytate level was between 1.7 mg/100g in *Cucurbitapepo* 4.1 mg/100g to *Leptadenia*  
 185 *hastata*. Results obtained are below the toxic level (FAO/WHO, 2004). According to Oke (1989) a  
 186 phytate diet of 1-6% over a long period of time decreases the bioavailability of mineral elements in  
 187 mono gastric animals.

188 Cyanogens are glycosides of a sugar, sugars and cyanide containing aglycone. Cyanogens  
 189 can be hydrolyzed by enzymes to release a volatile HCN gas (Nkafamiya and Manji, 2006). Excess  
 190 cyanide ion inhibits the cytochrome oxidase which stops ATP formation and so tissues suffer energy  
 191 deprivation and death follows rapidly. High level of HCN has been implicated for cerebral damage and  
 192 lethargy in man and animal (Nkafamiya and Manji, 2006). In table 3, the levels were 7.59 % in  
 193 *Cucurbita pepo* and 4.82% in *Balanites aegyptiaca* which are below lethal level as indicated on the  
 194 table 3. As for saponins, level ranged between 11.85 % in *Leptadenia hastata* to 15.13 % in *Vitex*  
 195 *doniana*. Saponins are glycosides containing polycyclic aglycone moiety of either C<sub>27</sub> steroid or  
 196 C<sub>30</sub>triterpenoids attached to a carbohydrate. High saponin level has been associated with  
 197 gastroenteritis manifested by diarrhoea and dysentery (Sodipo *et al.*, 2000).

198 Tannins are water soluble phenolic compounds with a molecular weight greater than 500 and  
 199 with the ability to precipitate proteins from aqueous solution. They occur in all vascular plants. Tannin  
 200 binds to proteins making them bio unavailable (Bagepallis *et al.*, 1993). From the results, the level  
 201 obtained was between 0.51% in *Balanites aegyptiaca* and 0.60 % in *Vitex donina*. There was no  
 202 significant difference in the Tannin level among the vegetables (p>0.05).

203 Alkaloids have been implicated in the inhibition activities of many bacterial species  
 204 (Nuhu *et al.*, 2000). The levels of alkaloids in the various leafy vegetables analysed ranges from  
 205 16.22% observed in *Letadenia hastata* and 19.37% in *Vitex doniana*  
 206 Plant phenolics such as flavonoids have been shown to have antioxidant properties and also  
 207 contribute to their medicinal significance (Wong *et al.*, 2006). In this work, the levels of flavonoids  
 208 ranged from 9.87 % in *Balanites aegyptiaca* to 14.71 % in *Vitex doniana*. Consumption of these  
 209 vegetables would further enhance the capacity of the cell to mop up the highly reactive oxygen  
 210 radicals generated due to oxidative metabolic reactions that occur in cells.

211

212 **TABLE 1:** Amino acid composition of the samples

AMINO ACIDS g/100g proteins	<i>Cucurbita pepo</i>	<i>Vitex doniana</i>	<i>Hibiscus cannabinus</i>	<i>Letadenia hastata</i>	<i>Balanites aegyptiaca</i>	MEAN	S.D	C.V %
L	8.14	6.48	7.29	8.34	6.48	7.35	0.88	12.03
K	4.03	3.63	3.97	5.54	4.37	4.31	0.74	17.11
I	3.30	3.01	2.94	4.78	3.40	3.49	0.75	21.47
F	4.17	4.43	4.34	5.58	4.70	4.64	0.56	12.00
W	2.25	1.26	2.10	2.86	1.12	1.92	0.72	37.78
V	4.50	3.59	4.00	5.61	4.09	4.36	0.77	17.69

M	1.12	1.23	1.71	2.00	0.80	1.37	0.48	34.95
P	2.84	3.14	2.84	4.57	2.23	3.12	0.87	27.96
R	4.99	6.11	5.33	7.83	4.99	5.85	1.20	20.47
Y	2.75	3.09	3.09	4.13	3.09	3.23	0.52	16.23
H	2.23	2.17	2.39	2.11	2.65	2.31	0.22	9.39
C	1.09	1.21	1.57	2.18	0.78	1.37	0.54	39.22
A	3.71	4.40	3.94	5.40	4.28	4.35	0.65	14.95
E	9.99	11.28	11.58	12.11	10.97	11.19	0.79	7.06
G	4.08	4.11	3.75	4.46	3.89	4.06	0.27	6.61
T	2.88	2.99	3.27	3.38	2.88	3.08	0.23	7.51
S	2.99	3.29	3.13	3.83	3.25	3.30	0.32	9.69
D	8.31	8.00	7.38	8.49	7.44	7.92	0.50	6.33
<b>TOTAL</b>	<b>73.37</b>	<b>73.42</b>	<b>74.62</b>	<b>93.20</b>	<b>71.41</b>	<b>77.20</b>	<b>9.02</b>	<b>11.68</b>

213

214

215 **TABLE 2:** Levels of metals in the samples

SAMPLES	Pb	Cd	Mg	Mn	Fe
concentration(ppm)					
<i>Cucurbita pepo</i>	0.0229±0.004	0.003±0.001	1.0871±0.002	0.0557±0.001	0.7997±0.003
<i>Vitex doniana</i>	0.0302±0.002	0.0036±0.001	1.1371±0.010	0.0499±0.001	1.0799±0.100
<i>Hibiscus cannabinus</i>	0.0396±0.021	0.029±0.001	1.1255±0.004	0.0653±0.004	0.9159±0.012
<i>Letadenia hastata</i>	0.0021±0.001	0.037±0.010	0.9951±0.010	0.1331±0.000	2.2231±0.900
<i>Balanites aegyptiaca</i>	0.0552±0.012	0.041±0.001	1.0753±0.002	0.0405±0.001	0.7723±0.123
<i>permissible levels</i>	0.10	0.05	-	-	-

216 Values represent means of triplicate determination± SD

217 Source for permissible limits: FAO (Mg/L) recommended maximum Concentration of trace elements  
218 for vegetables (1985).

219

220 **TABLE 3:** Phytochemical content of the samples

SAMPLES	Tannins (%)	Oxalate (mg/100g)	Phytate (mg/100g)	Saponins (%)	Cyanogenic Glycosides (mg/100g)	Alkaloids (%)	Flavonoids (%)
<i>Cucurbita pepo</i>	0.53±.012	0.60±0.20	1.70±0.01	12.49±1.20	7.59±1.20	17.45±2.51	13.72±1.21
<i>Vitex doniana</i>	0.60±0.12	0.14±0.01	2.10±0.21	15.13±1.50	5.33±1.11	19.37±1.23	14.71±2.30
<i>Hibiscus cannabinus</i>	0.59±0.13	0.32±0.12	3.14±0.12	13.35±2.35	4.93±2.13	17.10±2.50	10.38±1.30
<i>Letadenia hastate</i>	0.55±0.21	0.22±0.03	4.10±0.01	11.85±1.85	6.21±1.20	16.22±1.61	13.22±1.23
<i>Balanites aegyptiaca</i>	0.51±0.13	0.31±0.01	3.08±0.02	13.48±3.12	4.82±1.30	18.13±2.31	9.87±1.32

221 The values represent the mean of three determinations± SD.



222

223 **CONCLUSION AND CONTRIBUTION TO KNOWLEDGE**

224 From these results, leaves of the vegetables constitute rich sources of amino acids and mineral  
225 elements. The vegetables are therefore rich alternatives (and/or supplements) which can replenish  
226 the cellular requirements of the nutrients. Also, samples contain antinutritional factors such as  
227 oxalates and phytates but at low levels. The content of flavonoids and alkaloids in appreciable  
228 amounts in the samples is critical given the therapeutic/medicinal use of phytochemicals. Hence, leafy  
229 vegetables could contribute to the alleviation of protein malnutrition and micronutrient deficiencies if  
230 consumed.

231

232

233

234 **REFERENCES**

235 1. Adeyeye EI, Otokiti, MKO. Proximate composition and some nutritionally valuable minerals of two  
236 varieties of *Capsicum annum* (Bell and Cherry peppers). *Discovery and Innovation* 11:75-81 1999;  
237 Aegean region. *J. Turk. Weed Scie* 3:56-6.

238

239 2. Aloskar LV, Kakkar KK, Chakra OJ. Second Supplement to Glossary of Indian Medicinal Plants  
240 with Active Principles, 1992. Part-I (A-K), NISC, CSIR, New Delhi, pp: A.O.A.C. (Association of Official  
241 Agriculture Chemists).

242

243 3. Aschner JL, Aschner M. Nutritional aspect of manganese homeostasis. *Molecular aspects of*  
244 *Medicine*. 2005

245

246 4. Bagepallis S, Narasinga R, Tatinemi P. Tannin contents of foods commonly consumed in India and  
247 its influence on ionisable iron. *J. Sci. Food Afric*. 1993; 33:89-96.

248

249 5. Bainbridge ZK, Tomlins A, Westby A. Analysis of condensed tannins using acidified vanillin. *J.*  
250 *Food Sci. Agric*. 1996; 29: 77-79.

251

252 6. Day RA, Underwood AL. Quantitative analysis. 5th ed. Prentice Hall, 701. 1986.

253

254 7. Donald ES, Collins WB, Thomas AH, Cassara NE, Carnaha AM. The impact of tannins on protein,  
255 dry matter, and energy digestion in moose (*Alces alces*). *Canadian Journal of Zoology*. 2010; 88: 977-  
256 987.

257

258 8. FAO/WHO. Contaminants. In: *Codex Alimentarius* (1st ed, XVII), FAO/WHO, 1984.

259

260 9. FAO/WHO. United Nations Food and Agriculture Organization/World Health Organization: Fruit and  
261 Vegetables for Health. Report of a Joint FAO/WHO Workshop, Kobe, Japan, 2004.

262

263 10. Fasusi AO. Nutritional potentials of some tropical vegetable meals. Chemical characterization and  
264 functional properties. *Afr. J. Biotechnol*. 2006; 5(1): 49-53.

265

266 11. Grases F, Rafaei MP, Antonis C. Dietary Phytate and Interactions with Mineral Nutrients. In book:  
267 *Clinical Aspects of Natural and Added Phosphorus in Foods* 10.1007/978-1-4939-6566-3\_12. 2017.

267

268 12. Guoyao, W. Functional Amino Acids in growth, reproduction and health. *Advances in Nutrition*.  
2010; 1(1): 31-37.

269

270 13. Heyes RB. The Carcinogenicity of metals in humans: *Cancer Causes Control* 8:371-385. 1997.

- 271 14. Hilou A, Nacoulma OG, Guiguemde TR. *In vivo* antimalarial activities of extract from  
272 AmaranthusspinosusL and BoerhaaviaaerectaL. J. Ethnopharmacol. 2006; 103: 236-240.  
273
- 274 15. Kaya I, Incekara N. Contents of some wild plants species consumed as food in Aegean region.  
275 2010; J. Turk. Weed Scie 3:56-64.  
276
- 277 16. Kubmarawa DI, Andenyang FH, Magomya AM. Proximate composition and amino acid profile of  
278 two non-conventional leafy vegetables (*Hibiscus cannabinus* and *Haematostaphisbarteri*). Pakistan  
279 Journal of Nutrition. 2013; 12 (10): 949-956 ISSN 1680-5194.  
280
- 281 17. Nesamvuni C, Steyn NP Potgieter MJ. Nutritional value of wild, leafy plants consumed by the  
282 Vhavenda. South African Journal Science. 2001; 97: 51-54.  
283
- 284 18. Nkafamiya II, Manji AJ. A study of cyanogenetic glucoside contents of some edible Nuts and  
285 Seeds. J. Chem. Soc. 2006; Niger 31 (1 and 2) 12-14  
286
- 287 19. Nkafamiya II, Modibbo UU, Manji AJ, Haggai D. Nutrient Content of Seeds of Some Wild Plants.  
288 Afr. J. Biotechnol. 2007; 6(15): 1665-1669.  
289
- 290 20. Nuhu AM, Mshelia MS, Yakubu Y. Antimicrobial screening of the bark extract of *Pterocarpus*  
291 *serinaceus* tree. J. Chem. Soc. Nig. 2000; 28: 85-87.  
292
- 293 21. Oke, OL. Chemical studies on the more commonly used vegetables in Nigeria. Afr.Sci. Ass.  
294 11:42-48 Okiek of Kenya. Economic Botany 1989; 55, 32–46.
- 295 22. Olubunmi AO, Olaofe O, Olunayo RA. Amino Acid Composition of Ten Commonly Eaten  
296 Indigenous Leafy Vegetables of South-West Nigeria. World Journal of Nutrition and Health. 2017;  
297 3(1).
- 298 23. Osagie AU, Offiong UE. Nutritional Quality of Plant foods, Ambik Press, Benin City, Edo State,  
299 Nigeria 1998: pp. 131-221.  
300
- 301 24. Reddy MB, Love M. The impacts of food processing on the nutritional quality of vitamins and  
302 minerals. Adv. Exp. Med. Bio.459 1999; 99-106.  
303
- 304 25. Ryan, MF. The role of magnesium in clinical biochemistry: an overview. Ann Clin Biochemistry.  
305 1991; 28:19-26.  
306
- 307 26. Schumacher M, Bosque, MA, Domingo JL, Corbella J. Dietary intake of lead and cadmium from  
308 food in Tarragona Province, Spain. Bull. Env. Cont.Toxicol. 1991; 46:320-328.  
309
- 310 27. Sodipo OA, Akiniyi JA, Ogunbamosu JU. Studies on certain characteristics of extracts of bark of  
311 *Pansinystali amacruceas* (K schemp) picrre Exbeille. Global J. Pure Appl. Sci. 2000; 6:83-87.  
312
- 313 28. Sparkman DH, Stein EH, Moore S. Automatic recording apparatus for use in Chromatography of  
314 amino acids, Anal. Chem. 1958; 30; 119.  
315
- 316 29. Wong SP, Leong LP, Koh, JH. Antioxidant activities of aqueous extracts of selected plants.Food  
317 Chemistry.2006; 775-783.  
318
- 319 30. Yiridoe EK, Anchirinah VM. Garden production systems and food security in Ghana:  
320 Characteristics of traditional knowledge and management systems. Renew. Agric. Food Syst.2005;  
321 20: 168-180.  
322
- 323 31. Zayed AM, Terry N. Chromium in the environment: factors affecting biological remediation Plant.  
324 Plant Soil. 2003; 249:139-156.