

1 **Original Research Article**  
2 **Response of sugar beet to graded levels of**  
3 **nitrogen, phosphorous and potash on nutrient**  
4 **uptake and economics**  
5  
6  
7  
8  
9  
10

11 **ABSTRACT**

Field experiment was undertaken during 2005-06 to 2006-07 to study the various agrotechniques for sugar beet cultivation for Northern Karnataka at Agricultural Research Station, Bailhongal, Belgaum district (Karnataka) under irrigated condition. The experiment consisted of 28 treatment combinations comprising of graded levels of nitrogen, phosphorus and potassium. Design of the experiment was randomized block design with factorial concept. Application of 180, 90 and 120 kg ha<sup>-1</sup> of nitrogen, phosphorus and potassium, respectively registered significantly higher nitrogen, phosphorus and potassium uptake by sugar beet compared to other levels of nutrient. The same dose of nutrient application also improved gross returns, net returns and BC ratio.

12 Keywords: Sugar beet, nutrient uptake, economics, nitrogen, phosphorus and potassium  
13  
14  
15

16 **1. INTRODUCTION**  
17

18 Sugar beet is a long day plant, which requires adequate moisture and bright  
19 sunshine for good growth. Seeds germinate between soil temperature range of 12-15° and  
20 high sugar accumulation is observed in temperature of 20-22°C whereas, temperature  
21 exceeding 30°C adversely affect sugar accumulation. However, recently developed tropical  
22 sugar beet varieties require an optimum temperature range of 20-25°C for germination, 30-  
23 35°C for growth and development and 25-35°C for sugar accumulation, wherein the night  
24 15-20°C is suitable. The crop does not prefer high rainfall or continuous heavy rain which  
25 may affect development of tuber and sugar synthesis [1]. Tropicalised varieties of sugar beet  
26 developed make it possible to grow the crop in the tropical and subtropical areas. The crop  
27 matures within 5 to 6 months, requires moderate water requirement of 60- 80 cm, tolerant to  
28 soil water stress [2], less fertilizer requirement, provides about 60-80 tonnes of roots tuber  
29 yield per hectare. Sugar beet root contains 16-19 per cent sucrose with a recovery of 12-14  
30 per cent in the process of sugar extraction. Besides the sugar beet crop matures in March-  
31 April when the crushing season is nearly over as the harvesting period of sugar beet  
32 coincides with the off season of sugar factories. Thus, the supply of sugar beet can extend  
33 the crushing period of mills by nearly 2 months in the off season. It helps in continuous  
34 functioning of the sugar mills and thus reduces the cost of sugar production.

35 Owing to concerns and problems associated with sugarcane cultivation and  
36 potential production feasibilities associated with the sugar beet production indicated greater  
37 perspectives for the sugar beet cultivation as economically viable and potential sugar crop  
38 for crop diversification in the sugarcane grown area. Decision making process in crop  
39 production like selection of best genotypes, date of sowing, fertilizer application and date of  
40 maturity for harvesting which form prime agronomic practices for evaluating the performance

**Comment [A1]:** But not the highest BC !! Make recommendation accordingly.

41 of crop and extending hand in improvement of yield as well as the quality parameters needs  
42 critical adjustment. The scientific information on different agro-techniques to be adopted for  
43 cultivation of sugar beet is not available as it is completely new to this region. The technical  
44 information regarding the cultivation of sugar beet will be helpful for the cultivators of the  
45 region to harvest good yield. Being an introduced crop in the country, there is an urgent  
46 need to undertake research on tropical sugar beet in the country in general and north  
47 Karnataka in particular. Hence, the research work has major focus on analyzing the  
48 optimum fertilizer requirement for higher yield and quality of sugar beet.  
49

## 50 2. MATERIAL AND METHODS

51  
52 Field experiment was undertaken during 2005-06 to 2006-07 to study the optimum  
53 fertilizer requirement for higher yield and quality of sugar beet for Northern Karnataka at  
54 Agricultural Research Station, Bailhongal, Belgaum district (Karnataka) under irrigated  
55 condition. The experiment consisted of 28 treatment combinations comprising of graded  
56 levels of nitrogen, phosphorus and potassium. The initial soil pH was 7.20, available N, P<sub>2</sub>O<sub>5</sub>  
57 and K<sub>2</sub>O were 216, 17 and 270 kg ha<sup>-1</sup>. The organic carbon was 0.48 % and EC 0.23 dSm<sup>-1</sup>.  
58 For analyzing growth and development of the crop, five plants were selected at random from  
59 each net plot area in each treatment and were tagged to record various biometric  
60 observations. The average values were used for analysis. Fischer's method of analysis of  
61 variance was used for analysis and interpretation of the data as outlined by [3]. The level of  
62 significance used in 'F' and 'T' tests was p=0.05. Critical differences were calculated  
63 wherever 'F' test was significant.  
64

Comment [A2]: Add details.

### 65 2.1.1 Plant analysis

66  
67 The plant samples of sugar beet collected for dry matter production studies at  
68 harvest were analyzed for nitrogen, phosphorus and potash contents after drying in hot air  
69 oven at 70°C and powdered in micro-wiley mill. Nitrogen estimation was done by Kjeldahl's  
70 method (4) phosphorus by vanado molybdate phosphoric yellow colour method and  
71 potassium by flame photometric method.  
72

73 Based on nutrient content of plants and dry matter production, uptake of nitrogen,  
74 phosphorus and potassium were worked out by using following formula  
75

$$\text{Nutrient uptake} = \frac{\text{Per cent nutrient concentration}}{100} \times \text{Biomass (kg ha}^{-1}\text{)}$$

### 76 2.1.2 Economics of the system

#### 77 2.1.2.1 Cost of cultivation

78  
79 It was worked out on the basis of cost of labour, inputs and other costs for sugar  
80 beet.  
81

#### 82 2.1.2.2 Gross return (Rs. ha<sup>-1</sup>)

83  
84 It was worked out on the basis of market rates prevailing at the time of harvest of the  
85 produce.  
86

#### 87 2.1.2.3 Net return (Rs. ha<sup>-1</sup>)

88  
89 Net return was calculated by subtracting the cost of cultivation (Rs. ha<sup>-1</sup>) from  
90 the gross return.  
91  
92

93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147

### 3. RESULTS AND DISCUSSION

#### 3.1 Effect of graded levels of nitrogen, phosphorus and potassium on nutrient uptake by sugar beet

Nutrient uptake by of sugar beet differed significantly due to graded levels of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O application in beet tops, roots and total (Table 1,2 and 3).

Application of nitrogen @ 180 kg ha<sup>-1</sup> recorded significantly higher N uptake in beet tops (48.6 kg ha<sup>-1</sup>), beet roots (212.3 kg ha<sup>-1</sup>) and total uptake (260.9 kg ha<sup>-1</sup>) The uptake of N was significantly low in the level 60 kg ha<sup>-1</sup> in top (33.7 kg ha<sup>-1</sup>), roots (128.4 kg ha<sup>-1</sup>) and total (162.1 kg ha<sup>-1</sup>).

Among the phosphorus levels, application of P at 90 kg ha<sup>-1</sup> recorded significantly higher P uptake in beet tops (44.4 kg ha<sup>-1</sup>), beet roots (187.7 kg ha<sup>-1</sup>) and total uptake (232.1 kg ha<sup>-1</sup>) The uptake of P was significantly low in the level 30 kg ha<sup>-1</sup> in top (36.9 kg ha<sup>-1</sup>), roots (155.8 kg ha<sup>-1</sup>) and total (192.7 kg ha<sup>-1</sup>).

Application of potassium @ 120 kg ha<sup>-1</sup> recorded significantly higher K uptake in beet tops (19.10 kg ha<sup>-1</sup>), beet roots (160.4 kg ha<sup>-1</sup>) and total uptake (179.5 kg ha<sup>-1</sup>) The uptake of K was significantly low in the level 90 kg ha<sup>-1</sup> in top (18.5 kg ha<sup>-1</sup>), roots (158.1 kg ha<sup>-1</sup>) and total (176.6 kg ha<sup>-1</sup>).

The optimum dose of nitrogen, phosphorus and potassium was essential for getting higher yield below which the yield reduces and above which the cost of production increases. The present study revealed that 180, 90 and 120 kg ha<sup>-1</sup> nitrogen, phosphorus and potassium was found economically viable for sustainable production of sugar beet, Similar findings were obtained by [5, 6 and 7].

#### 3.2 Effect of graded levels of nitrogen, phosphorus and potassium on economics

##### 3.2.1. Gross returns

The gross returns obtained from the sugar beet was varied significantly due to application of different levels of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O during both the years of experimentation and in their pooled data (Table 3).

Among the N levels, significantly higher gross returns was obtained with the application of nitrogen @ 180 kg ha<sup>-1</sup> (Rs. 1,28,437 ha<sup>-1</sup>) as compared to lower N levels @ 60 kg ha<sup>-1</sup> (Rs. 1,02,705 ha<sup>-1</sup>). However, it was on par with N applied @ 120 kg ha<sup>-1</sup> (Rs. 1,28,010 ha<sup>-1</sup>). Application of phosphorus at higher dose @ 90 kg ha<sup>-1</sup> (Rs. 1,22,944 ha<sup>-1</sup>) recorded significantly higher gross returns as compared to lower dose @ 30 kg ha<sup>-1</sup> (Rs. 1,13,992 ha<sup>-1</sup>). However, it was at par with P<sub>2</sub>O<sub>5</sub> applied @ 60 kg ha<sup>-1</sup> (Rs. 1,22,216 ha<sup>-1</sup>). The application of potassium @ 120 kg ha<sup>-1</sup> recorded significantly higher gross returns (Rs. 1,22,902 ha<sup>-1</sup>) as compared to its lower dose @ 60 kg ha<sup>-1</sup> (Rs. 1,14,575 ha<sup>-1</sup>). However, it was on par with K<sub>2</sub>O applied @ 90 kg ha<sup>-1</sup> (Rs. 1,21,674 ha<sup>-1</sup>).

The interaction effect of N × P<sub>2</sub>O<sub>5</sub> and N × K<sub>2</sub>O at different levels of application had significant influence on gross returns obtained from sugar beet. Among the N × P<sub>2</sub>O<sub>5</sub> interaction, 180:30/60/90 or 120:60/90 kg and P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> recorded significantly higher gross returns as compared to interactions and were on par with each other. Application of N and K<sub>2</sub>O @ 180/120:90/120 kg ha<sup>-1</sup> recorded on par gross returns and were significantly superior than other treatment combinations. As compared to fertilized treatments control treatment recorded significantly lower gross returns (Rs. 65,040 ha<sup>-1</sup>). The higher dose of nutrient improved the vegetative growth and enhanced the rate of production of assimilates

148 from source to sink, which ultimately increased the nitrogen uptake. The improved yield also  
149 increased the gross returns. Similar results were obtained by [8 and 9]  
150

### 151 3.2.2. Net returns

152 The net returns obtained from the sugar beet was varied significantly due to  
153 application of different levels of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O during both the years of experimentation  
154 and in their pooled analysis (Table 3).  
155

156 Among the N levels, significantly higher net returns were obtained with the  
157 application of nitrogen @ 120 kg ha<sup>-1</sup> (Rs. 97,369 ha<sup>-1</sup>) as compared to lower dose of N @  
158 60 kg ha<sup>-1</sup> (Rs. 72,589 ha<sup>-1</sup>). However, it was on par with N applied @ 180 kg ha<sup>-1</sup> (Rs.  
159 97,271 ha<sup>-1</sup>). Application of P<sub>2</sub>O<sub>5</sub> @ 90 kg ha<sup>-1</sup> resulted in significantly higher net returns  
160 (Rs. 91,783 ha<sup>-1</sup>) as compared to lower dose of P<sub>2</sub>O<sub>5</sub> @ 30 kg ha<sup>-1</sup> (Rs. 83,871 ha<sup>-1</sup>).  
161 However, it was on par with application of P<sub>2</sub>O<sub>5</sub> @ 60 kg ha<sup>-1</sup> (Rs. 91,575 ha<sup>-1</sup>). Application  
162 of K<sub>2</sub>O @ 120 kg ha<sup>-1</sup> resulted in significantly higher net returns (Rs. 92,036 ha<sup>-1</sup>) as  
163 compared to K<sub>2</sub>O applied @ 60 kg ha<sup>-1</sup> (Rs. 84,159 ha<sup>-1</sup>). However, it was on par with K<sub>2</sub>O  
164 applied @ 90 kg ha<sup>-1</sup> (Rs. 91,033 ha<sup>-1</sup>).  
165

166 The combined application of N × P<sub>2</sub>O<sub>5</sub> and N × K<sub>2</sub>O at different levels of application  
167 had significant influence on net returns obtained by sugar beet. Among the N × P<sub>2</sub>O<sub>5</sub>  
168 combinations, significantly higher net returns were obtained with the application of 120:60,  
169 120:90, 180:30, 180:60 and 180:90 kg ha<sup>-1</sup> as compared to other treatment combinations  
170 and were on par with each other. Among the N × K<sub>2</sub>O interactions, N applied @ 120/180  
171 irrespective of the K<sub>2</sub>O levels recorded significantly higher net returns as compared to N  
172 applied in lower dose (60 kg ha<sup>-1</sup>) irrespective of K levels. As compared to fertilizer applied  
173 treatments, control with no fertilizer recorded significantly lower net returns (Rs. 37,164 ha<sup>-1</sup>).  
174 Improved yield of the crop with lesser cost of production, consequently improved the net  
175 returns. Similar results were obtained by [10 and 11]  
176

### 177 3.2.4. BC ratio

178 The benefit cost ratio obtained from the sugar beet cultivation differed significantly  
179 due to graded levels of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O application during both the years of  
180 experimentation and in their pooled analysis (Table 3).  
181

182 Among the N levels, significantly higher B:C ratio was obtained both the application  
183 of N @ 120 kg ha<sup>-1</sup> (4.06) as compared to N applied @ 60 kg ha<sup>-1</sup> (3.31). However, it was  
184 on par with N applied at higher doses *i.e.*, 180 kg ha<sup>-1</sup> (4.03). Application of P<sub>2</sub>O<sub>5</sub> @ 60 kg  
185 ha<sup>-1</sup> recorded significantly higher B:C ratio (3.88) as compared to P<sub>2</sub>O<sub>5</sub> @ 30 kg ha<sup>-1</sup> (3.68).  
186 However, it was at par with P<sub>2</sub>O<sub>5</sub> applied @ 90 kg ha<sup>-1</sup> (3.84). Among the K<sub>2</sub>O levels,  
187 application of K<sub>2</sub>O @ 120 kg ha<sup>-1</sup> recorded significantly higher B:C ratio (3.89) as compared  
188 to K<sub>2</sub>O applied @ 60 kg ha<sup>-1</sup> (3.67). However, it was on par with K<sub>2</sub>O applied @ 90 kg ha<sup>-1</sup>  
189 (3.85).  
190

191 The combined application of N × P<sub>2</sub>O<sub>5</sub> and N × K<sub>2</sub>O had significant influence on  
192 B:C ratio. Among the N × P<sub>2</sub>O<sub>5</sub> applied @ 120:60 kg ha recorded significantly higher  
193 B:C ratio (9.27). However, it was on par with 120:90 and 180:30 kg N and P<sub>2</sub>O<sub>5</sub> ha. Among  
194 the N × K<sub>2</sub>O interactions significantly higher B:C ratio was obtained with the application of  
195 120:90 kg N and K<sub>2</sub>O ha<sup>-1</sup> (4.14) and was on par with all other treatments except N applied  
196 at lower dose (60 kg ha<sup>-1</sup>) irrespective of K<sub>2</sub>O levels. The benefit from the rupees investment  
197 was higher in 120, 90 and 120 kg nitrogen, phosphorus and potassium which is ascribed to  
198 improved net returns and yield of the crop. Similar results were noticed by [12 and 13]  
199

200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250

## CONCLUSION

The present study inferred that application of nitrogen 120 kg, phosphorus 90 kg and potassium 120 kg ha<sup>-1</sup> was found optimum for getting higher nutrient uptake and economically viable approach of sugar beet.

1. Ali MA, Alvi SM, Cheema SA, 2004, Sowing date and plant spacing effect on agro-qualitative traits of sugar beet (*Beta vulgaris*) in different ecological zones of Punjab [Pakistan]. *J. Agric. Res.*, 42(1) : 41-52.
2. Hills FJ, Broadbent FE, Fried, M, 1990, Timing and rate of fertilizer nitrogen for sugar beet related to nitrogen uptake and pollution potential. *J. Environ.*, 7 : 368-372.
3. Gomez KA, Gomez AA, 1984, *Statistical Procedures for Agricultural Research*, 2<sup>nd</sup> Edn. John Wiley and Sons, New York, USA.
4. Jackson ML, 1973, *Soil Chemical Analysis*, Prentice Hall of India Pvt. Ltd., New Delhi : 498.
5. Hellal FA, Taalab AS, Safaa AM, 2009, Influence of nitrogen and boron nutrition on nutrient balance and Sugar beet yield grown in calcareous Soil. *Ozean J. Appl. Sci.*, 2(1) : 1-10.
6. Majumdar B, Venkatesh MS, Kailashkumar, Patiram, 2005, Effect of potassium and farmyard manure on yield, nutrient uptake and quality of ginger (*Zingiber officinale*) in typic hapludalf of Meghalaya. *Indian J. Agric. Sci.*, 75(12) : 809-811.
7. Nemeat Alla EAE, EL-Geddawy IHM, 2001, Response of sugar beet to foliar spraying time with micronutrients under different levels of nitrogen and phosphorus fertilization. *J. Agric. Sci., Mansoura Univ.*, 26(4) : 670-681.
8. Albert L, Sims, 2010, Sugar beet response to broadcast and starter phosphorus applications in the Red River Valley of Minnesota. *Agron. J.*, 102(5) : 1369-1378.
9. Ali MK, Nujma M, 2011, Sugar beet crop - an alternative cane. *New Agri Technology*, pp 2831.
10. Balakrishnan A, 2006, Introduction of tropical sugar beet cultivation with suitable varieties in Tamil Nadu. In : Scheme completion report, 2006 centre for soil and crop management studies, Tamil Nadu Agricultural University, Coimbatore, India.
11. Camas N, Crak C, Albayrak S, 2007, Yield and quality component of sugar beet grown under Northern Turkey conditions. *Intl. J. Agric. Res.*, 2(3) : 296-301.
12. EL-Harriri DM, Mirvat EG, 2001, Response of growth, yield and quality of sugar beet to nitrogen and potassium fertilizers under newly reclaimed sandy soil. *J. Agric. Sci., Mansoura Univ.*, 26(10) : 5895-5907.
13. EL-Zayat MMT, 2000, Effect of irrigation regimes and fertilization on sugar beet. *Ph. D. Thesis*, in Agric. Sci. (Agron.), Fac. of Agric., Kafr EL-Sheikh, Tanta Univ.

Comment [A3]: Not the highest BC !!

**Table 1. N uptake by sugar beet as influenced by graded levels of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O (Pooled data of 2005-06 and 2006-07)**

Treatment		N uptake by beet top (kg/ha)				N uptake by tuber (kg/ha)				Total N uptake (kg/ha)			
		N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	Mean	N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	MEAN	N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	MEAN
P <sub>30</sub>	K <sub>60</sub>	18.4	33.3	35.8	29.2	78.7	148.8	177.6	135.0	97.0	182.1	213.4	164.2
	K <sub>90</sub>	25.6	43.5	55.5	41.5	108.5	176.4	208.5	164.5	134.1	220.0	264.0	206.0
	K <sub>120</sub>	38.3	40.8	41.0	40.1	124.5	164.4	214.8	167.9	162.8	205.2	255.8	207.9
	Mean	27.4	39.2	44.1	36.9	103.9	163.2	200.3	155.8	131.3	202.4	244.4	192.7
P <sub>60</sub>	K <sub>60</sub>	29.4	33.9	48.0	37.1	109.0	167.9	192.0	156.3	138.4	201.7	240.0	193.4
	K <sub>90</sub>	36.0	49.5	41.7	42.4	133.2	203.1	214.0	183.4	169.2	252.7	255.7	225.9
	K <sub>120</sub>	35.7	40.8	58.4	45.0	148.2	179.6	211.3	179.7	183.8	220.4	269.7	224.7
	Mean	33.7	41.4	49.4	41.5	130.1	183.5	205.8	173.1	163.8	224.9	255.1	214.6
P <sub>90</sub>	K <sub>60</sub>	38.5	40.9	51.1	43.5	129.4	169.4	223.1	174.0	167.8	210.3	274.2	217.5
	K <sub>90</sub>	44.7	42.4	47.6	44.9	170.9	201.0	236.2	202.7	215.6	243.3	283.8	247.6
	K <sub>120</sub>	37.0	39.3	58.4	44.9	153.4	172.5	233.3	186.4	190.4	211.8	291.7	231.3
	Mean	40.1	40.9	52.4	44.4	151.2	180.9	230.9	187.7	191.3	221.8	283.2	232.1
Mean of K	K <sub>60</sub>	28.8	36.0	45.0	36.6	105.7	162.0	197.6	155.1	134.4	198.1	242.5	191.7
	K <sub>90</sub>	35.4	45.1	48.3	42.9	137.5	172.1	219.6	178.0	172.9	212.5	267.8	221.3
	K <sub>120</sub>	37.0	40.3	52.6	43.3	142.0	193.5	219.8	183.5	179.0	238.7	272.4	226.5
Mean	33.7	40.5	48.6		128.4	175.9	212.3		162.1	216.4	260.9		
Control		20.1				39.1				59.5			
For comparison of means	S.Em+	CD @ 5%		S.Em+	CD @ 5%		S.Em+	CD @ 5%					
Nitrogen (N)	0.98	2.77		3.67	10.41		4.16	11.81					
Phosphorus (P)	0.98	2.77		3.67	10.41		4.16	11.81					
Potassium (K)	0.98	2.77		3.67	10.41		4.16	11.81					
N x P	1.72	4.88		6.47	NS		7.34	20.82					
N x K	1.72	NS		6.47	NS		7.34	NS					
P x K	1.72	4.88		6.47	NS		7.34	NS					
N x P x K	2.98	8.46		11.21	NS		12.71	NS					
Control vs Treatments	2.98	8.46		11.21	31.82		12.71	36.07					

1 **Table 2. P uptake by sugar beet as influenced by graded levels of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O (Pooled data of 2005-06 and 2006-**  
 2 **07)**

Treatment	P uptake by beet top (kg/ha)				P uptake by tuber (kg/ha)				Total P uptake (kg/ha)				
	N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	Mean	N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	MEAN	N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	MEAN	
P <sub>30</sub>	K <sub>60</sub>	2.2	4.1	5.0	3.8	18.0	27.6	37.8	27.8	20.3	31.6	42.8	31.6
	K <sub>90</sub>	3.0	5.4	7.5	5.3	20.5	32.8	41.6	31.6	23.5	38.2	49.2	37.0
	K <sub>120</sub>	4.6	5.7	5.8	5.4	24.0	35.1	43.9	34.3	28.6	40.8	49.7	39.7
	Mean	3.3	5.1	6.1	4.8	20.8	31.8	41.1	31.3	24.1	36.9	47.2	36.1
P <sub>60</sub>	K <sub>60</sub>	3.5	4.4	6.7	4.9	20.2	33.7	39.4	31.1	23.7	38.1	46.1	36.0
	K <sub>90</sub>	4.3	6.6	5.6	5.5	24.8	40.8	41.7	35.8	29.2	47.4	47.3	41.3
	K <sub>120</sub>	4.5	6.1	8.1	6.3	29.3	43.1	42.1	38.1	33.8	49.2	50.2	44.4
	Mean	4.1	5.7	6.8	5.6	24.8	39.2	41.1	35.0	28.9	44.9	47.9	40.6
P <sub>90</sub>	K <sub>60</sub>	4.6	5.5	6.5	5.5	23.9	34.9	40.6	33.1	28.5	40.3	47.2	38.7
	K <sub>90</sub>	5.3	5.9	6.2	5.8	29.8	42.9	42.6	38.4	35.1	48.9	48.8	44.3
	K <sub>120</sub>	5.4	6.2	7.9	6.5	34.9	43.6	43.5	40.7	40.2	49.8	51.4	47.1
	Mean	5.1	5.9	6.9	5.9	29.5	40.5	42.2	37.4	34.6	46.3	49.1	43.4
Mean of K	K <sub>60</sub>	3.4	4.6	6.1	4.7	20.7	32.0	39.3	30.7	24.2	36.7	45.4	35.4
	K <sub>90</sub>	4.2	6.0	6.5	5.6	25.1	38.8	42.0	35.3	29.3	44.8	48.4	40.8
	K <sub>120</sub>	4.8	6.0	7.3	6.0	29.4	40.6	43.2	37.7	34.2	46.6	50.4	43.8
Mean	4.2	5.6	6.6		25.0	37.2	41.5		29.2	42.7	48.1		
Control	2.4				10.5				12.9				
For comparison of means	S.E.m+	CD @ 5%		S.E.m+	CD @ 5%		S.E.m+	CD @ 5%					
Nitrogen (N)	0.13	0.37		0.62	1.75		0.68	1.92					
Phosphorus (P)	0.13	0.37		0.62	1.75		0.68	1.92					
Potassium (K)	0.13	0.37		0.62	1.75		0.68	1.92					
N x P	0.23	NS		1.09	3.09		1.19	3.39					
N x K	0.23	NS		1.09	NS		1.19	NS					
P x K	0.23	NS		1.09	NS		1.19	NS					
N x P x K	0.40	1.14		1.89	NS		2.07	NS					
Control vs Treatments	0.40	1.14		1.89	5.36		2.07	5.87					

3  
4

5 **Table 3. K uptake by sugar beet as influenced by graded levels of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O (Pooled data of 2005-06 and 2006-**  
 6 **07)**

Treatment		K uptake by beet top (kg/ha)				K uptake by beet tuber (kg/ha)				Total K uptake (kg/ha)			
		N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	Mean	N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	MEAN	N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	MEAN
P <sub>30</sub>	K <sub>60</sub>	9.8	15.6	16.5	14.0	108.4	147.1	169.4	141.6	118.2	162.7	185.9	155.6
	K <sub>90</sub>	13.0	18.6	23.4	18.3	123.1	152.6	172.5	149.4	136.1	171.2	195.9	167.7
	K <sub>120</sub>	18.6	18.8	16.9	18.1	135.3	156.3	171.1	154.2	153.9	175.1	188.0	172.3
	Mean	13.8	17.7	18.9	16.8	122.3	152.0	171.0	148.4	136.1	169.7	189.9	165.2
P <sub>60</sub>	K <sub>60</sub>	14.4	15.9	21.0	17.1	117.9	166.5	167.3	150.6	132.3	182.3	188.3	167.6
	K <sub>90</sub>	17.5	21.3	17.1	18.6	139.5	176.4	169.7	161.9	157.0	197.7	186.8	180.5
	K <sub>120</sub>	16.8	19.3	23.7	20.0	148.8	181.3	164.5	164.9	165.6	200.6	188.3	184.8
	Mean	16.2	18.8	20.6	18.6	135.4	174.7	167.2	159.1	151.7	193.6	187.8	177.7
P <sub>90</sub>	K <sub>60</sub>	18.1	19.3	20.8	19.4	127.8	167.8	172.9	156.2	145.8	187.1	193.7	175.5
	K <sub>90</sub>	19.1	18.1	18.2	18.5	146.6	175.3	167.4	163.1	165.7	193.4	185.7	181.6
	K <sub>120</sub>	17.4	18.6	21.7	19.3	153.1	174.3	159.2	162.2	170.5	192.9	181.0	181.5
	Mean	18.2	18.7	20.3	19.0	142.5	172.5	166.5	160.5	160.7	191.1	186.8	179.5
Mean of K	K <sub>60</sub>	14.1	16.9	19.4	16.8	118.0	160.5	169.8	149.4	132.1	177.4	189.3	166.3
	K <sub>90</sub>	16.5	19.3	19.6	18.5	136.4	168.1	169.9	158.1	152.9	187.4	189.5	176.6
	K <sub>120</sub>	17.6	18.9	20.8	19.1	145.7	170.6	165.0	160.4	163.3	189.5	185.8	179.5
Mean	16.1	18.4	19.9		133.4	166.4	168.2		149.5	184.8	188.2		
Control		11.32				63.8				72.9			
For comparison of means		S.Em+		CD @ 5%		S.Em+		CD @ 5%		S.Em+		CD @ 5%	
Nitrogen (N)		0.38		1.08		2.28		6.48		2.34		6.64	
Phosphorus (P)		0.38		1.08		2.28		6.48		2.34		6.64	
Potassium (K)		0.38		1.08		2.28		6.48		2.34		6.64	
N x P		0.67		NS		4.03		11.43		4.12		11.70	
N x K		0.67		NS		4.03		11.43		4.12		11.70	
P x K		0.67		NS		4.03		NS		4.12		NS	
N x P x K		1.16		NS		6.98		NS		7.14		NS	
Control vs Treatments		1.16		3.30		6.98		19.80		7.14		20.27	



**Table 4. Economics of sugar beet as influenced by graded levels of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O (Pooled data of 2005-06 and 2006-07)**

Treatment		Cost of cultivation (Rs./ha)				Gross returns (Rs./ha)				Net returns (Rs./ha)				B:C ratio			
		N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	Mean	N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	MEAN	N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	MEAN	N <sub>60</sub>	N <sub>120</sub>	N <sub>180</sub>	MEAN
P <sub>30</sub>	K <sub>60</sub>	29959	30484	31009	30484	82685	112279	129044	108002	52726	81796	98035	77519	2.76	3.68	4.16	3.54
	K <sub>90</sub>	30184	30709	31234	30709	93805	116337	131756	113966	63621	85629	100523	83258	3.11	3.79	4.22	3.71
	K <sub>120</sub>	30409	30934	31459	30934	103041	119155	130486	117561	72632	88222	99027	86627	3.39	3.85	4.15	3.80
	Mean	30184	30709	31234	30709	93177	115924	130429	113176	62993	85215	99195	82468	3.09	3.78	4.18	3.68
P <sub>60</sub>	K <sub>60</sub>	30479	31004	31529	31004	90094	127013	127626	114911	59615	96009	96098	83907	2.96	4.10	4.05	3.70
	K <sub>90</sub>	30704	31229	31754	31229	106499	134508	129476	123494	75796	103280	97722	92266	3.47	4.31	4.08	3.95
	K <sub>120</sub>	30929	31454	31979	31454	113442	138193	124832	125489	82514	106740	92854	94036	3.67	4.40	3.91	3.99
	Mean	30704	31229	31754	31229	103345	133238	127312	121298	72642	102010	95558	90070	3.37	4.27	4.01	3.88
P <sub>90</sub>	K <sub>60</sub>	30999	31524	32049	31524	97579	128026	130809	118805	66580	96503	98760	87281	3.15	4.06	4.08	3.77
	K <sub>90</sub>	31224	31749	32274	31749	111812	133611	126021	123814	80588	101863	93747	92066	3.58	4.21	3.91	3.90
	K <sub>120</sub>	31449	31974	32499	31974	116645	133000	121312	123652	85197	101026	88814	91679	3.71	4.16	3.73	3.87
	Mean	31224	31749	32274	31749	108678	131546	126047	122090	77455	99797	93774	90342	3.48	4.15	3.91	3.84
Mean of K	K <sub>60</sub>	30479	31004	31529	31004	90119	122439	129160	113906	59641	91436	97631	82902	2.96	3.95	4.10	3.67
	K <sub>90</sub>	30704	31229	31754	31229	104038	128152	129084	120425	73335	96924	97331	89196	3.39	4.10	4.07	3.85
	K <sub>120</sub>	30929	31454	31979	31454	111043	130116	125543	122234	80114	98663	93565	90781	3.59	4.14	3.93	3.89
Mean	30704	31229	31754		101733	126903	127929		71030	95674	96176		3.31	4.06	4.03		
Control	28464				65065				36602				2.29				
For comparison of means	S.Em <sub>±</sub>		CD @ 5%		S.Em <sub>±</sub>		CD @ 5%		S.Em <sub>±</sub>		CD @ 5%		S.Em <sub>±</sub>		CD @ 5%		
Nitrogen (N)	-		-		1740		4937		1740		4937		0.06		0.16		
Phosphorus (P)	-		-		1740		4937		1740		4937		0.06		0.16		
Potassium (K)	-		-		1740		4937		1740		4937		0.06		0.16		
N x P	-		-		3068		8708		3068		8708		0.10		0.28		
N x K	-		-		3068		8708		3068		8708		0.10		0.28		
P x K	-		-		3068		NS		3068		NS		0.10		NS		
N x P x K	-		-		5315		NS		5315		NS		0.17		NS		
Control vs Treatments	-		-		5315		15082		5315		15082		0.17		0.49		

UNDER PEER REVIEW

