

## Effect of GA<sub>3</sub>, spacing and nutrient sprays on growth and flowering in snapdragon (*Antirrhinum majus* L.) cv. Rocket Pink

### ABSTRACT

An experiment was conducted to study the effect of different concentrations of GA<sub>3</sub>, spacing and nutrient sprays on vegetative and floral parameters of snapdragon (*Antirrhinum majus* cv. Rocket pink) at Urban Technological Park, Habak, Srinagar, J&K during two successive years in 2017 and 2018. Eighteen different treatments with 3 concentrations of GA<sub>3</sub> (0 ppm, 100 ppm and 200 ppm), 2 spacings (15 cm x 15 cm and 15 cm x 20 cm) and nutrient sprays (3 sprays, 4 sprays and 5 sprays) were replicated thrice in Completely Randomized Block Design. The investigation revealed that GA<sub>3</sub> at 200 ppm proved best among all in vegetative including height (91.07 cm), number of primary branches (4.11) and number of leaves (174.60) as well as in floral parameters including days to first inflorescence initiation (47.77), days to first inflorescence opening (50.81), days to 50% flowering (54.30), length of flower stalk (45.91 cm), length of inflorescence (38.80 cm), number of spikes per plant (7.72), number of florets per spike (22.35), number of fully developed florets (5.17), duration of flowering (52.62 days) and vase life (13.00 days). Spacing of 15 cm x 15 cm proved better in terms of height (85.23 cm) in vegetative parameters and length of flower stalk (43.81 cm) and length of inflorescence (37.65 cm) in floral parameters while spacing of 15 cm x 20 cm proved better in number of primary branches (3.64) and number of leaves (170.68) as well as days to first inflorescence initiation (49.49), days to first inflorescence opening (52.33), days to 50% flowering (55.42), number of florets per spike (17.64), number of fully developed florets (4.47), duration of flowering (50.75 days) and vase life (10.71 days). Among different nutrient sprays application of 5 sprays proved best in all vegetative including height (87.06 cm), number of primary branches (4.26) and number of leaves (182.92) and floral parameters including length of flower stalk (42.88 cm), length of inflorescence (36.18 cm), number of spikes per plant (7.80), number of florets per spike (19.21), number of fully developed florets (4.67), duration of flowering (52.26 days) and vase life (10.92 days). While minimum days to first inflorescence initiation (49.05), first inflorescence opening (51.89) and 50% flowering were obtained by 3 nutrient sprays.

**Key words:** Snapdragon, GA<sub>3</sub>, Spacing, Nutrient sprays

### INTRODUCTION

*Antirrhinum majus* L. commonly known as snapdragon or dog flower is native to Mediterranean region. It is abundant in the temperate regions of the world. In India it is represented by 273 species and is commercially cultivated in the states of Assam, Gujrat, Uttar Pradesh, Maharashtra, Karnataka and Tamil Nadu. It is one of the principal cut flower crops wherein the flowers are borne on terminal long spikes of many colours and shades. The variation in plant

height (12-36 inches) in different types and groups provide wide scope of using *Antirrhinum* for different purposes.

Spacing of plants is an important aspect that needs to be carefully considered. Proper spacing between the plants can efficiently utilize the available area for healthy growth and thus result in the better yield. Inaba and Ohshiro [1] reported the yield of cut flowers per plant decreases with an increase in the planting density. However, yield per unit area increases. Inaba *et al.* [2] reported decrease in the weight of cut flowers and the number of axillary buds with the increase in planting density.

Fertilizers give nutrition to the plant and can develop tolerance against pests and diseases. Fertilizers don't only assist in increasing yields and promoting healthy growth of plants but also in their development. Snapdragon shows a great response to different types of fertilizers. With the increase in the concentration of nitrogen fertilizers there is increase in the number of leaves, number of branches, longevity of inflorescence and number of lateral inflorescence Sanjeeta and Maiti [3]. Majeed and Ali [4] also reported that with an increase in the concentration of nitrogen there is an increase in plant height, fresh weight of the plant, stem diameter, number of leaves per plant, number of branches per plant, number of spikes per plant, number of florets per spike and flower spike length.

Ornamental crops find extensive use of growth regulators for modifying their developmental processes. The major areas where growth regulators have successfully played their roles in ornamental plants are in vegetative propagation, inhibition of abscission, prevention of bud dormancy, growth control, promotion of flowering, prolonging the vase life of flowers and retarding their senescence Prakash and Jha [5]. Growth regulators have a great impact on the growth and flowering of snapdragon. GA<sub>3</sub> has been found to increase the stem length and vase life of snapdragon Sarhan and El-Sayed [6] and advance flowering by 6-7 days Verzilov and Rinkova [7]. Te-Sen and Toop [8] reported significant increase in the length of inflorescence, overall height of the plant at maturity and the fresh of top of snapdragon by GA<sub>3</sub>. Therefore, GA<sub>3</sub>, spacing and nutrient sprays were manipulated to improve growth and flowering of snapdragon.

## **MATERIALS AND METHODS**

The experiment was carried out at Urban Technology Park, Habak, Srinagar during 2017 and 2018. It is situated between 35.5° - 34.7° North latitude and 74.8° - 74.9° East longitude at an altitude of 1,588 m above mean sea level. In general, the climate of the area is temperate-cum-mediterranean and of continental type characterized by hot summers and severe winters. Hottest months are July and August during which temperature shoots up 36°C. Winter is severe, extending over 70 days from the middle of December to March, when the temperature often goes below the freezing point and the whole of Kashmir valley remains covered under snow. A Randomized Complete Block Design was used comprising of three replications. The land was divided into three blocks each with a width of 0.6 m and length of 16.2 m, leaving a path of 0.3 m between the blocks. Each block was divided into 18 plots of 0.6 x 0.6 m size separated by a

path of 0.3 m. Uniform dose of fertilizers N, P and K= 10 g each) and FYM was added to each plot at the final preparation prior to planting of seedlings. Uniform sized seedlings of snapdragon (*Antirrhinum majus*) cv. Rocket pink were planted in the field on 22<sup>nd</sup> of march at a spacing of 15 cm x 15 cm and 15cm x 20 cm in the alternative beds/plots and were slightly watered by rose can. Factors used were GA<sub>3</sub> (0 ppm, 100 ppm and 200 ppm), Spacing (15 cm x 15 cm and 15 cm x 20cm) and Nutrient sprays (19:19:19+Calmax Gold) (3 sprays, 4 sprays and 5 sprays). The stock solution of GA<sub>3</sub> and nutrient sprays were prepared by dissolving them in appropriate solvents and followed by dilution with distilled water. Spraying of micronutrient solutions (CalMax Gold and 19:19:19) was done uniformly with the help of garden sprayer. First spray of GA<sub>3</sub> (100 and 200 ppm) was done 40 days after planting (1st May). First dose of 19:19:19 and CalMax Gold was sprayed on 3<sup>rd</sup> of May. Plants were taken randomly from each treatment in every replication at the appropriate time to record the observations. The observations were recorded for growth parameters such as plant height, number of leaves and number of primary branches; for flowering parameters days to first inflorescence initiation, days to first inflorescence opening, days to 50% flowering, length of flower stalk (cm), length of inflorescence (cm), number of spikes per plant, number of florets per spike, number of fully developed florets, duration of flowering (days), vase life (days). Data analysis was done using Microsoft Excel and OPSTAT.

## Result

**Gibberellic acid:** Results presented in table 1 revealed that plant height, number of primary branches and number of leaves were significantly increased by GA<sub>3</sub>. Highest plant height (91.07 cm), maximum number of primary branches (4.11) and maximum number of leaves (174.60) were obtained at 200 ppm, while the lowest (79.07cm, 3.24 and 160.58, respectively) were obtained at control. GA<sub>3</sub> significantly decreased the number of days to first inflorescence initiation, first inflorescence opening and 50% flowering. Minimum number of days to first inflorescence initiation (47.77), first inflorescence opening (50.81) and 50% flowering (54.30) were observed at 200 ppm while the maximum number of days (51.15, 54.06 and 57.62 respectively) were observed in control. There was a significant increase in length of flower stalk and inflorescence length by GA<sub>3</sub>. Longest flower stalk (45.91 cm) and inflorescence (38.80 cm) were obtained at 200 ppm while the shortest (38.85 cm and 30.63 cm respectively) were obtained at control. Number of spikes per plant, number of florets per spike and number of fully developed florets obtained maximum value (7.72, 22.35 and 5.17 respectively) at 200 ppm while the lowest (6.38, 14.66 and 3.76 respectively) at control. Duration of flowering and vase life were significantly increased by GA<sub>3</sub>. Maximum duration (52.62 days) and vase life (13.00 days) were obtained at 200 ppm while the minimum (47.59 days and 7.72 days respectively) were obtained by control.

**Spacing:** spacing significantly effected the plant height, number of primary branches and the number of leaves. Higher plant height (85.23 cm) was observed in higher density while low density recorded lower plant height (84.90 cm). Number of primary branches and number of leaves were higher (3.64 and 170.68 respectively) at low density while lower (3.49 and 164.11 respectively) at high density. Days to first inflorescence initiation, first inflorescence opening and 50% flowering were earlier (49.49, 52.33, and 55.42 respectively) in case of high density than at low density (49.91, 52.91 and 56.18 respectively). Length of flower stalk and inflorescence were longer (43.81cm and 37.65 cm respectively) in case of high density than in low density (40.71cm and 33.32 cm respectively). Spacing had insignificant effect on the number of spikes per plant. Number of florets per spike , number of fully developed florets, duration of flowering and vase life were significantly effected by spacing, obtaining higher values (17.64, 4.47, 50.75 days and 10.71 days respectively) at low planting density than those (17.29, 4.23, 49.67 days and 9.96 days respectively) at higher density.

**Nutrient sprays:** Plant height, number of primary branches and number of leaves were significantly increased by the application of nutrient spray obtaining highest values (87.06 cm, 4.26, 182.92 respectively) by 5 sprays while lowest (83.06 cm, 2.90, 154.61 respectively) by control. Days to first inflorescence initiation, days to first inflorescence opening and 50% flowering were minimum (49.05, 51.89, and 55.14 respectively) at 3 sprays while maximum (50.04, 53.14 and 56.49 respectively) at 5 sprays. Length of flower stalk and inflorescence were longest (42.88 cm and 36.18 cm respectively) by 5 sprays while shortest (41.45 cm and 34.78 cm respectively) by 3 sprays. Number of spikes per plant were maximum (7.80) by 5 sprays while minimum (6.97) by 3 sprays. Number of florets per spike and number of fully developed florets were maximum (19.21 and 4.67 respectively) by 5 sprays while minimum (16.05 and 4.03 respectively) by 3 sprays. Duration of flowering and vase life were significantly increased by nutrient sprays. Maximum duration (52.26) and vase life (10.92) were obtained by 5 sprays their minimum value (48.57 and 9.84 respectively) were obtained by 3 sprays.

## **Discussion**

**Gibberellic acid:** GA<sub>3</sub> had a significant effect on all the three growth parameters including plant height, number of primary and number of leaves. Highest plant height, number of primary branches and number of leaves were obtained at 200 ppm. Highest plant height is due to cell elongation caused by GA<sub>3</sub>. Our findings for number of branches and leaves are in consent with those of [9] who advocated that by increasing GA<sub>3</sub> concentration there is increase in branch number in petunia cv. Carnival. Due to increase in the branch number there is increase in the number of leaves. GA<sub>3</sub> had a significant effect on all the floral parameters. GA<sub>3</sub> resulted in early inflorescence, inflorescence opening and 50% flowering. The earliest inflorescence initiation, inflorescence opening and 50% flowering were obtained at 200 ppm. Our findings are in consent with those of [10] in liliun hybrids and [11] in liliun longiforum who advocated the induction of early flowering by GA<sub>3</sub>. There was a significant increase in the length of flower stalk and length of inflorescence. The longest flower stalk and inflorescence were obtained at 200 ppm. This is

due to cell elongation caused by GA<sub>3</sub>. Highest number of spikes per plant were obtained at 200 ppm. This is due increase in the number of branches which increases the number of spikes. Number of florets per spike, number of fully developed florets, duration of flowering and vase life were significantly increased by GA<sub>3</sub> and were maximum at 200 ppm. This might be due to the fact that GA<sub>3</sub> treated plants produced more number of leaves as compared to control, which might have resulted in production and accumulation of more photosynthates that were diverted to flowers. The findings are in agreement with those of [12] who found that GA<sub>3</sub> significantly increased leaf area and number of florets per spike in *Gladiolus*, cv. Friendship and [4] for number of florets per spike and number of fully developed florets, [3] in case of duration of flowering and [6] in case of vase life.

**Table 1: Effect of GA<sub>3</sub>, spacing and nutrient sprays on growth parameters in snapdragon (*Antirrhinum majus* L.) cv. “Rocket Pink”**

Treatment	Plant height (cm)	Number of leaves	Number of primary branches
<b>GA<sub>3</sub> (G)</b>			
<b>Without GA<sub>3</sub> application (G<sub>0</sub>)</b>	<b>79.07</b>	<b>160.58</b>	<b>3.240</b>
<b>GA<sub>3</sub> 100 ppm (G<sub>1</sub>)</b>	<b>85.06</b>	<b>166.99</b>	<b>3.371</b>
<b>GA<sub>3</sub> 200 ppm (G<sub>2</sub>)</b>	<b>91.07</b>	<b>174.598</b>	<b>4.108</b>
<b>C.D (p≤ 0.05)</b>	<b>0.05</b>	<b>2.445</b>	<b>0.066</b>
<b>Spacing (S)</b>			
<b>15cm × 15cm (S<sub>1</sub>)</b>	<b>85.233</b>	<b>164.106</b>	<b>3.488</b>
<b>15cm × 20cm (S<sub>2</sub>)</b>	<b>84.900</b>	<b>170.678</b>	<b>3.658</b>
<b>C.D (p≤ 0.050)</b>	<b>0.041</b>	<b>1.997</b>	<b>0.053</b>

<b>Nutrient sprays (N)</b>			
<b>3 Sprays (N<sub>1</sub>)</b>	<b>83.063</b>	<b>154.609</b>	<b>2.900</b>
<b>4 Sprays (N<sub>2</sub>)</b>	<b>85.066</b>	<b>164.645</b>	<b>3.556</b>
<b>5 Sprays (N<sub>3</sub>)</b>	<b>87.064</b>	<b>182.922</b>	<b>4.264</b>
<b>C.D (p≤ 0.050)</b>	<b>0.050</b>	<b>2.445</b>	<b>0.066</b>

**Spacing:** It was observed that the magnitude of plant height was higher at the closer spacing (15cm x 15 cm). The higher plant height might be due to intra plant competition for light and aeration which promoted the elongation of the main stem. The effect of closer spacing was supported by [13]. Higher number of primary branches and leaves are obtained by wider spacing (15cm x 20cm). This might be due to wider spacing for branch development which then increased the number of leaves as well. Our findings are in consent with [14] who reported number of primary branches were higher at wider spacing. Spacing significantly influenced the number of days to first inflorescence initiation, first inflorescence opening and 50% flowering. First inflorescence initiation, inflorescence opening and 50% flowering were earlier in closer spacing. Length of flower stalk and length of inflorescence obtained higher values in case of closer spacing. This might be due to high competition for light and aeration in closer spacing. Spacing had insignificant effect on the number of spikes per plant. Number of florets per spike, number of fully developed florets, duration of flowering and vase life were significantly influenced by spacing. Broader spacing resulted in higher number of florets per spike, fully developed florets, flowering duration and vase life. This might be due to more photosynthates produced by more number of leaves in case of wider spacing.

**Table 2: Effect of GA<sub>3</sub>, spacing and nutrient sprays on flowering parameters in snapdragon (*Antirrhinum majus* L.) cv. “Rocket Pink”**

<b>Treatment</b>	<b>Days to first inflorescence initiation</b>	<b>Days to first inflorescence opening</b>	<b>Days to 50% flowering</b>	<b>Length of flower stalk (cm)</b>	<b>Length of inflorescence (cm)</b>
<b>GA<sub>3</sub> (G)</b>					
<b>Without GA<sub>3</sub> application (G<sub>0</sub>)</b>	<b>51.148</b>	<b>54.064</b>	<b>57.617</b>	<b>38.850</b>	<b>30.627</b>
<b>GA<sub>3</sub> 100 ppm</b>	<b>50.172</b>	<b>53.003</b>	<b>55.481</b>	<b>42.019</b>	<b>37.030</b>

(G <sub>1</sub> )					
GA <sub>3</sub> 200 ppm (G <sub>2</sub> )	47.768	50.807	54.304	45.908	38.801
C.D (p≤ 0.050)	0.053	0.022	0.033	0.036	0.024
<b>Spacing (S)</b>					
15cm × 15cm (S <sub>1</sub> )	49.487	52.337	55.423	43.813	37.650
15cm × 20cm (S <sub>2</sub> )	49.905	52.911	56.179	40.705	33.322
C.D (p≤ 0.050)	0.043	0.018	0.027	0.029	0.020
<b>Nutrient sprays (N)</b>					
3 Sprays (N <sub>1</sub> )	49.047	51.889	55.135	41.452	34.784
4Sprays (N <sub>2</sub> )	49.999	52.840	55.781	42.441	35.493
5 Sprays (N <sub>3</sub> )	50.042	53.144	56.486	42.884	36.180
C.D (p≤ 0.050)	0.053	0.022	0.033	0.036	0.024

**Nutrient sprays:** Nutrient sprays significantly influenced all the three vegetative parameters. Highest plant height, maximum number of primary branches and number of leaves were obtained by 5 sprays. This might be due to availability of more nutrients in case of 5 sprays. Days to first inflorescence initiation, first inflorescence opening and 50% flowering were significantly influenced by application of nutrient sprays. Minimum number of days to initiation, opening and 50% flowering were observed by 3 sprays while maximum number of days were recorded by 5 sprays. Length of flower stalk and length of inflorescence were maximum in case of 5 sprays while shortest of these were recorded by 3 sprays. This might be due availability of more photosynthates due to more number of leaves in case of 5 sprays. Number of spikes, number of florets per spike and number of fully developed florets were maximum in 5 sprays while the minimum were recorded by 3 sprays. Our results are in consent with [3] and [4]. Duration of flowering and vase life were significantly increased by nutrient sprays. Maximum duration and

vase life were recorded by 5 sprays while 3 sprays recorded the minimum. Our results are in consent with [3] and [6].

**Table 3: Effect of GA<sub>3</sub>, spacing and nutrient sprays on flowering parameters in snapdragon (*Antirrhinum majus* L.) cv. “Rocket Pink”**

Treatment	Number of spikes per plant	Number of florets per spike	Number of fully developed florets	Duration of flowering (days)	Vase life (days)
<b>GA<sub>3</sub> (G)</b>					
<b>Without GA<sub>3</sub> application (G<sub>0</sub>)</b>	<b>6.375</b>	<b>14.657</b>	<b>3.763</b>	<b>47.594</b>	<b>7.724</b>
<b>GA<sub>3</sub> 100 ppm (G<sub>1</sub>)</b>	<b>7.702</b>	<b>15.403</b>	<b>4.122</b>	<b>50.418</b>	<b>10.283</b>
<b>GA<sub>3</sub> 200 ppm (G<sub>2</sub>)</b>	<b>7.724</b>	<b>22.347</b>	<b>5.174</b>	<b>52.619</b>	<b>13.000</b>
<b>C.D (p≤ 0.050)</b>	<b>0.612</b>	<b>0.373</b>	<b>0.050</b>	<b>0.198</b>	<b>0.095</b>
<b>Spacing (S)</b>					
<b>15cm × 15cm (S<sub>1</sub>)</b>	<b>7.126</b>	<b>17.294</b>	<b>4.232</b>	<b>49.671</b>	<b>9.957</b>
<b>15cm × 20cm (S<sub>2</sub>)</b>	<b>7.408</b>	<b>17.644</b>	<b>4.474</b>	<b>50.750</b>	<b>10.714</b>
<b>C.D (p≤ 0.050)</b>	<b>N.S</b>	<b>0.305</b>	<b>0.040</b>	<b>0.162</b>	<b>0.077</b>
<b>Nutrient sprays (N)</b>					
<b>3 Sprays (N<sub>1</sub>)</b>	<b>6.973</b>	<b>16.049</b>	<b>4.083</b>	<b>48.565</b>	<b>9.836</b>
<b>4Sprays (N<sub>2</sub>)</b>	<b>7.031</b>	<b>17.149</b>	<b>4.312</b>	<b>49.810</b>	<b>10.253</b>



<b>5 Sprays (N<sub>3</sub>)</b>	<b>7.796</b>	<b>19.209</b>	<b>4.665</b>	<b>52.257</b>	<b>10.918</b>
<b>C.D (p≤ 0.050)</b>	<b>0.612</b>	<b>0.373</b>	<b>0.050</b>	<b>0.198</b>	<b>0.095</b>

#### REFERENCES:

1. Inaba, Z. and Ohshiro, M. 2005. Effect of planting density and methods of raising seedlings on flowering, yield and quality of cut flowers in snapdragon. *Environment control in Biology*. **43** (3), 201-210.
2. Inaba, Z., Kato, C., Horiuchi, M. and Ohtsuka, H. 2010. Effect of planting density, container size and period of raising seedlings on the flowering and quality of snapdragon cut flowers in non-pinching culture. *Horticultural Research Japan*. **9** (2), 165-170.
3. Sanjeeta, P. and Maiti, R.G. 2008. Study on the response of nitrogen fertilization in field grown Antirrhinum. *Journal of Interacademia*. **12** (2), 176-181.
4. Mjeed, A.J. and Ali, M.A. 2017. Effect of Gytia and Nitrogen applications on growth and flowering of snapdragon plants in two soil depths. *Kurdistan journal for applied research*. Vol. 2, Issue 1.
5. Prakash, V. and Jha, K.K. 1998. Physiology of gladiolus. *Journal of Applied Biology*. **8**: 24-28.
6. Sarhan, A.Z. and El. Sayed, A.A. 1983. Effect of some growth regulators on growth and flowering of snapdragon plants. *Annals of Agricultural Science, Moshtohor*. **19** (2), 419-426.
7. Verzilov, V.F. and Rinkova, L.V. 1972. The effect of gibberellin and chloromequat on ornamental plants. *Izvestiya Akademii Nauk SSSR, Biologicheskaya*. **5**, 761-766.

8. Te-sen, P. and Toop, E.W. 1967. Effect of gibberellic acid on growth of *Antirrhinum majus* 'Utah White' in a carbon dioxide enriched environment. Department of Plant Sciences, university of Alberta, Edmonton.
9. Saleh, M., Safari, W.R., and Hassan, S. 2016. Effect of ascorbic acid, vitamin C, thiamine and GA<sub>3</sub> on growth parameters in petunia, cv. Carnival. *Journal of Horticultural Sciences*. Vol.30. pg141-150.
10. Dhiman, M.R. 1997. Effect of cold storage temperature and plant bio-regulators on growth and flower production in liliun hybrids. M.Sc thesis, Department of Floriculture and Landscape Architecture. Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan, India, pp. 22-55.
11. Pal, A.K. and Das, S.N. 1990. Effect of growth regulators on growth and flowering of *Lilium longiflorum*. Orissa. *Journal of Horticulture*. **18** : pg 18-21.
12. Pal, P. and T. Chowdhury. 1998. Elongation of flowering by GA<sub>3</sub>. *Journal of Horticulture*. Pg 69-77.
13. Singh, K.P. and Sangama. Response of china aster to spacing. *Journal of ornamental Horticulture*. 2001; 4(1):61-2.
14. Chanda S, Roychoudhary N. The effect of planting dated and spacing on growth, flowering and yield of African marigold (*Tageta erecta* L.)cv. Siracole. *The Horticulture Journal*. 1991;4(2):53-56.