

# Multiresidue Method for Determination of 74 Pesticides under Organic and Conventional Olive Farm Soil by QuEChERS and GC-MS/MSTQD

Comment [chpc1]:

## Abstract

The analytical multiresidue method, has been developed and validated for quantification of trace levels of 74 pesticide residues belong to different chemical groups of pesticides in organic and conventional Olive farm soil samples (Old, medial and new olive farms which is 25, 15 and 5 years respectively). Soil samples collected from Al-Jouf Province, Saudi Arabia, and extracted by Quick, Easy, Cheap, Effective, Rugged, and Safe (QuEChERS) and analyzed by Gas Chromatography Mass Spectrometry Triple Quadrupole (GC-MS/MSTQD). The method reveals that experimental results were highly satisfactory in respect of various analytical parameters such as linearity, recovery and precision especially with the tested soil samples which ~~is~~ are complex matrixes, preparation is a critical step, and one that is usually expensive, time-consuming, and labor intensive. The limit of detection (LOD) and limit of quantification (LOQ) for the analyzed pesticides were in the range of 1.01-13.91  $\mu\text{g kg}^{-1}$  and 3.02 - 29.15  $\mu\text{g kg}^{-1}$ , respectively. Pesticide recoveries form spiked soil samples with different pesticides ranged from 65.5 to 111.7 %. The proposed method featured good sensitivity, pesticide quantification limits were low enough, and the precision, expressed as relative standard deviation, ranged from 0.29 to 13.32%. Pesticide residues ~~was being detected~~ by applying the modified QuEChERS and GCMSMSTQD method, ~~the~~ levels ~~wasaswere~~ ranged from 43.00 to 2.00  $\mu\text{g kg}^{-1}$  for 18 different pesticides, 1.99 to 1.00  $\mu\text{g/kg}^{-1}$  for 16 different pesticides, 0.99 to 0.50  $\mu\text{g/kg}^{-1}$  for 12 different pesticides and lower than 0.50  $\mu\text{g kg}^{-1}$  for 28 different pesticide residues. The proposed QuEChERS and GC-MS/MSTQD method ~~were~~ applied successfully for the residues extraction and determination the 74 pesticides.

Formatted: Highlight

Formatted: Highlight

Formatted: Strikethrough

Formatted: Strikethrough, Highlight

Formatted: Strikethrough, Highlight

Formatted: Highlight

Formatted: Not Strikethrough

Formatted: Highlight

## Keywords

Multiresidue, Pesticide residues, Organic farming, Conventional farming, olive farm soil, QuEChERS and GC-MS/MSTQD.

## 31 Introduction

32 Pesticides are widely used in agriculture to protect crops, control the insects, and  
33 improve efficiency of food production. Due to the wide range of pesticides used in  
34 agriculture, the development of fast and simple multi-residue methods that  
35 simultaneously determine a wide range of different pesticides is essential. One of  
36 the most widely used multi-residue methodologies is the Quick, Easy, Cheap,  
37 Effective, Rugged, and Safe (QuEChERS) approach. This offers many advantages  
38 including speed, cost, ease of use, good performance characteristics and wide  
39 applicability range (Pszczolińska, and Michel, 2016).

40 Due to the low concentration levels of soil **pollutants** such as pesticides and other  
41 ~~pollutants~~ ~~substances~~, sample preparation step is needed to determine the type and  
42 quantity of ~~different pollutant~~ **such compounds** present (Caldas, et al., 2011; Wang,  
43 et al., 2012 and AOAC, 20117) ~~and~~ to avoid interferences and improve the  
44 sensitivity of the method. To ~~extract~~ ~~remove~~ contaminants from soil samples, a  
45 technique strong enough to **extract** bound pesticide residues in short time is  
46 necessary (Pinto et., al 2011 and Rashid et al., 2010).

47  
48 The QuEChERS approach is based on a salting-out extraction with a solvent  
49 (mainly acetonitrile, ACN) followed by a dispersive solid phase extraction (d-SPE).  
50 QuEChERS method is very flexible, modifiable, and is growing in popularity due to  
51 all the benefits described by its effectiveness is dependent on the analytic  
52 properties, matrix composition, equipment, and analytical technique availability  
53 (Pinto, et al., 2010; Caldas, et al., 2011; Wang et al., 2012).

54  
55 Soil samples are complex matrixes; therefore, soil sample preparation is a critical  
56 step, and one that is usually expensive, time-consuming, and labor intensive. The  
57 (QuEChERS) method, originally developed for the determination of pesticides in  
58 fruits and vegetables, recently modified and adopted for the analysis of pesticides in  
59 soil (Pszczolińska, and Michel, 2016, Brondi et al., 2011 and Fernandes, et al.,  
60 2013) **was employed in this study**. El-Saeid, et al., (2015) studied the levels of  
61 pesticide residues in two types of farmland soils, sandy and clay soils following  
62 different farming practices conventional and organic were taken from different  
63 depths **of** 10 and 20cm. Samples were prepared for extraction and were extracted

Formatted: Highlight

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough, Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

using acetone: hexane mixture (1:1) and cleanup was performed using florisil column. Clean extracts were subjected to pesticide residues determination of (a total of 86) pesticides belonging to different chemical and action groups using hyphenated GC-MS. Recovery, linearity and experimental limit of detection (LOD) experimental were performed. In case of sandy conventional farmland soil, the detected organochlorines (OCPs) pesticides were 7 or (seven), while the organophosphorus insecticides included four compounds pesticides. For herbicides two compounds were detected i.e. linuron and Amitraz. As for the frequency of the detected pesticide residues, it was found that the most frequent compounds were endosulfan I, chlorpyrifos-methyl, P,P-DDE, amitraz, fenthion, P,P-DDD, linuron, dimethoate, lindane, dieldrin, O,P-DDD, pirimiphos-methyl, alfa-BHC and aldrin. Also, it was observed that the detected pesticides residues were lindane, P,P-DDE, O,P-DDD, P,P-DDD, mirex, dieldrin and aldrin as a OCPs. It is clear that the highest amounts of OCPs groups residues distribution were especially at 20 cm followed by 10 cm (0.273 and 0.235 ppm.), while the numbers of detection pesticide residues at 20 cm depth more than 10cm were 23 and 15 numbers, respectively.

In this study, modified QuEChERS techniques used for the extraction and clean-up procedure followed by GCMSTQD for the analysis of several pesticide residues in soil samples collected from Olive cultivated under conventional and organic farming.

## Material and Methods

### Standards and Reagents

Pesticides internal, calibration and injection standards with declared 99.9% purity, were purchased from Accu Standard, 153 Inc., New Haven, CT, USA as individual or mixture standards at a concentration of 100-200 µg/mL. All internal standards are <sup>13</sup>C 12-labelled, the use of <sup>13</sup>C-labelled compound is preferable because the analysis can be quantified without clean-up. (Maestroni et al., 2000; Maestroni 2002). All solvents (Methanol, dichloromethane and acetonitrile) used for the extraction and analysis procedures of pesticides were residue-analysis grade 99.9 % purity and obtained from Fisher Scientific (Fair Lawn, NJ, USA). QuEChERS kits was were purchased from Phenomenex, Madrid Avenue, Torrance, CA, USA.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Strikethrough, Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Strikethrough, Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Strikethrough

Formatted: Strikethrough, Highlight

Formatted: Highlight

Formatted: Strikethrough, Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Strikethrough

Formatted: Highlight

Formatted: Highlight

Formatted: Strikethrough, Highlight

Formatted: Highlight

## **Samples preparation and Extraction**

First, weigh 10 g soil sample ( $\geq 70\%$  H<sub>2</sub>O content) into a 50 mL centrifuge tube. Alternatively, weigh 3 g air-dried soil sample into a 50 mL tube and add 7 mL H<sub>2</sub>O, vortex briefly, and allow to hydrate for 30 minutes. Then, add 10 mL of acetonitrile to each sample. Shake (manually or mechanically) or vortex samples for 5 minutes to extract pesticides. (In this study a SpexSamplePrep Geno/Grinder 2010 operated at 1500 rpm was used). After that, add the contents of an ECQUEU750CT-MP (citrate salts) Mylar pouch to each centrifuge tube. Immediately shake samples for at least 2 minutes and centrifuge for 5 minutes at  $\geq 3500$  rcf.

## **Sample Cleanup**

Transfer a 1 mL aliquot of supernatant to a 2 mL CUMPSC18CT (MgSO<sub>4</sub>, PSA, C18) dSPE tube. Vortex samples for 0.5–1 min. Centrifuge for 2 min at high rcf (e.g.  $\geq 5000$ ). Filter purified supernatant through a 0.2  $\mu$ m syringe filter directly into a sample vial. Finally, the samples were analyzed by GC-MS/MSTQD.

## **Analysis by GCMSMSTSQ 8000/SRM**

All measurements have been carried out using the latest Thermo Scientific™ TSQ 8000™ triple quadrupole GC-MS/MS system equipped with the Thermo Scientific™ TRACE™ 1310 GC with SSL Instant Connect™ SSL module and Thermo Scientific™ TriPlus™ RSH auto sampler. Injection mode was splitless, Splitless Time 1.0 min GC Column TR™ 5 MS, 30 m  $\times$  0.25 mm  $\times$  0.25  $\mu$ m, carrier gas He (99.999 %, flow rate 1.2 mL/min, constant flow, temperature program 100 °C, 1 min; 10 °C/min to 160 °C, 4 min and 10 °C/min to 250 °C, 2 min, transfer line temperature 280 °C, total analysis time 22.4 min, TriPlus RSH Autosampler Injection volume 1  $\mu$ L. Ionization mode EI, 70 eV, Ion source temperature 250 °C, scan mode SRM using timed SRM SRM transition setup automatically build-up by AutoSRM software. Transitions conditions are shown in (Table 1).

127

128

129 Table 1: GCMSMSTQD 8000 SRM Instrumental conditions

GC Trace Ultra Conditions		TSQ Quantum MS/MS Conditions	
Column	TR-Pesticide 30 m × 0.25 mm × 0.25 µm	Operating mode	Selected Reaction Monitoring (SRM)
Injector	Splitless	Ionization mode	EI
Injected volume	1 µL	Electron energy	70 eV
Injector temperature	225 °C	Emission current	50 µA
Carrier gas	Helium, 1.2mL/min	Q1/Q3 resolution	0.7 u (FWHM)
Oven program	80 °C hold 1 min 15 °C/min to 160 °C hold 1 min 2.2 °C/min to 230 °C hold 1 min 5 °C/min to 290 °C hold 5 min Run Time 57.15 min	Collision gas	Argon
Transfer line temperature	280 °C	Collision gas pressure	1 mTorr
		Polarity	Positive

130

## Method performance

Accuracy and precision of the method: 4 replicates of blank water sample spiked with the pesticide standards. Limit of detection: Instrument Detection Limit (IDL), Sample Detection Limit (SDL), Method Detection Limit, accuracy and precision

## QAQC Strategies

Quality control samples ~~was being~~ prepared and analyzed duplicate sample, blank and spiked, and/or Certified Reference material CRM was prepared for this purpose and processed with each batch (5-10 samples) of sample. ASE and GCMS or GCMSMS TSQ 8000 method limit of detection (LOD) and Limit of Quantification (LQD), repeatability, reproducibility, accuracy and precession also were determined for each compound in the groups of PAHs and Pesticides.

Formatted: Highlight

Formatted: Strikethrough, Highlight

Formatted: Highlight

## Results and Discussion

### QuEChERS and GC MS/MSTQD for analysis of 74 pesticides.

Formatted: Highlight

Simple and rapid method based on QuEChERS extraction and GCMSMSTQD for determination of 74 of different groups of pesticides in soil samples. Retention time, LOD, LOQ, recovery % and target mass of SRM scanning mode was determined as ~~showmen~~ in table (1). The results clearly reflect the developed QuEChERS method offers an efficient, cost effective, and easy sample preparation procedure for the determination of 74 substances or pesticides in soil samples.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Recovery % ranged from 65.5 to 111.7 %, the limit of detection (LOD) and limit of quantification (LOQ) for the analyzed pesticides were in the range of 1.01-13.35  $\mu\text{g kg}^{-1}$  and 3.02 - 29.15  $\mu\text{g kg}^{-1}$ , respectively. The proposed method featured good sensitivity, pesticide quantification limits were low enough, and the precision, expressed as relative standard deviation, ranged from 0.29 to 13.3%.

The calibration curves were linear over wide concentration ranges with correlation coefficients ( $r^2$ ) 0.5092 to 0.9899 for all tested pesticides. In addition, the SRM chromatograms demonstrated high selectivity with no significant interferences observed and an excellent signal/noise ratio ( $> 5:1$ ) for all tested pesticides as ~~showmen~~ in (Figure 1).

Formatted: Highlight

162 Table (2) Parameters of retention time, LOD, LOQ, recovery % and target mass of SRM scanning  
 163 mode.

Name	RT	Mass	Product mass	Collision Energy	LOQ	LOD	$r^2$	Recovery %	SD
Biphenyl	14.82	152	126	23	20.12	6.71	0.7134	108.60	6.32
Methacrifos	17.8	240	180	10	15.20	5.07	0.8379	98.21	6.46
Chloroneb	18.09	206	191	10	19.67	6.56	0.9522	100.42	12.08
Tecnazene	20.86	261	203	13	11.57	3.86	0.9848	104.02	4.98
Propachlor	20.97	120	77	19	8.14	2.71	0.9899	104.94	4.77
Cycloate	21.44	154	83	8	8.00	2.67	0.9905	111.73	3.76
Ethalfuralin	22.01	276	202	15	10.35	3.45	0.9848	113.32	9.52
Trifluralin	22.38	306	160	23	13.92	4.64	0.9760	106.95	7.46
Benfluralin	22.47	292	160	20	11.62	3.87	0.9816	111.04	4.67
Sulfotep	22.59	202	146	10	10.94	3.65	0.9806	110.87	10.89
Diallate	22.78	234	150	18	8.82	2.94	0.9650	86.56	7.30
Alph-BCH	23.04	181	145	13	21.32	10.44	0.8433	104.37	7.45
HCB	23.33	249	214	14	18.70	6.23	0.9382	103.88	0.35
Atrazine	23.94	200	122	10	11.84	3.95	0.9643	108.07	6.52
Terbufos	24.42	231	129	23	15.79	5.26	0.9570	96.64	8.69
Profluraline	24.48	318	199	17	13.67	4.56	0.9688	110.12	7.76
Fenofos	24.56	137	109	6	11.17	3.72	0.9813	110.32	4.49
Diazinone	24.81	137	84	12	21.73	13.91	0.9465	109.28	5.83
Fluchloralin	24.94	264	160	15	16.10	5.37	0.9535	106.12	5.52
Disulfoton	24.99	153	97	12	12.19	4.06	0.9729	87.14	6.75
Tefluthrin	25.14	177	127	15	4.33	1.44	0.9963	100.77	10.39
Triallate	25.28	270	186	18	18.68	6.23	0.9094	89.40	5.74
Endosulfan ether	25.73	272	237	10	24.04	13.35	0.9725	101.44	3.00
Pentachloroaniline	25.92	263	192	20	17.41	5.80	0.9365	102.47	0.29
Alachlor	26.31	146	118	8	17.41	5.80	0.9284	104.52	8.23
Vinclozolin	26.38	285	212	12	15.78	5.26	0.9633	75.14	1.30
Cypermethrin	26.45	163	91	11	15.09	5.03	0.9574	103.94	9.13
Heptachlor	26.62	100	65	12	21.00	11.33	0.8841	102.38	2.75
Acetochlor	26.72	174	146	12	15.56	5.19	0.9497	105.31	7.41


Fenchlorfos	26.84	285	240	23	9.91	3.30	0.9462	77.09	1.62
Primiphos methyl	27.22	290	125	20	17.45	10.82	0.5092	103.68	9.61
Pentachlorothioanisole	27.43	296	246	32	7.59	2.53	0.9922	93.27	1.54
Dichlofluanid	27.48	123	77	16	16.27	5.42	0.9311	69.56	12.45
Aldrin	27.66	263	191	35	22.69	12.23	0.9175	109.07	3.96
Chloropyrifos	27.81	314	258	12	25.88	11.96	0.9256	79.07	7.31
Triadimefon	27.9	208	181	10	12.62	4.21	0.9673	99.28	4.66
Primiphos - ethyl	28.35	318	166	12	5.35	1.78	0.9957	78.98	6.84
Isopropalin	28.44	280	238	8	16.73	5.58	0.9363	87.18	5.81
Isodrin	28.53	261	191	28	23.51	7.84	0.9895	74.27	3.63
Pendimethalin	28.72	252	162	10	23.72	10.24	0.5688	108.86	2.438
Heptachlor epoxide	28.81	353	263	13	21.48	8.49	0.7901	87.79	4.15
Tolyfluanid	28.88	240	137	10	23.91	7.97	0.8218	84.75	4.82
Quinalphos	29.04	298	156	8	8.32	2.77	0.9896	81.76	2.40
Procymidone	29.22	283	67	28	10.94	3.65	0.9655	97.83	1.65
Chlordane- cis	29.45	272	237	12	22.64	10.55	0.6677	95.03	1.24
o, p DDE	29.54	246	176	32	8.96	2.99	0.9908	107.02	2.23
Tetrachlorviphos	29.66	333	109	17	10.77	3.59	0.9805	82.67	8.85
Endosulfan 1	29.79	195	125	19	23.87	10.29	0.9873	96.46	5.35
Chlordane - trans	29.84	272	237	13	14.96	4.99	0.9394	103.58	2.93
Nonachlor- trans	29.95	409	302	22	8.88	2.96	0.9077	108.47	7.16
Pretilachlor	30.24	162	132	18	20.69	6.90	0.9806	82.81	7.75
p, p DDE	30.33	318	248	22	12.72	4.24	0.9478	110.46	8.60
Dieldrin	30.47	279	243	10	15.98	5.33	0.9486	105.50	7.11
o, p DDD	30.57	235	165	22	16.98	5.66	0.9048	97.09	3.34
Endrin	31.04	279	243	8	21.84	10.61	0.9529	105.27	7.74
Chlorobenzilate	31.09	139	111	12	15.03	5.01	0.8409	65.504	2.99
Endosulfan 2	31.26	241	206	10	18.86	6.29	0.9163	89.98	3.90
p, p DDD	31.38	235	165	24	8.43	2.81	0.9876	97.47	11.72
o, p DDT	31.46	235	165	21	10.00	1.61	0.9952	99.12	2.89
Nonachlor - cis	31.54	272	237	10	29.15	11.72	0.7072	111.22	5.03
Endrin - aldehyde	31.73	345	243	17	28.85	9.62	0.6997	104.519	7.11
Carfetraxone ethyl	31.93	340	312	10	16.56	5.52	0.9348	101.10	11.16



Methoxychlor olefin	31.99	308	223	30	20.34	6.78	0.8750	103.19	12.33
Endosulfan sulfate	32.28	274	239	12	19.86	12.29	0.6006	92.73	8.71
o, p Methoxychlor	32.49	227	121	12	19.95	6.65	0.7266	91.57	11.92
Resmethrin 1	32.57	123	81	8	15.18	5.06	0.8423	106.47	10.01
Resmethrin 2	32.67	123	81	8	21.08	7.03	0.7226	100.20	8.12
Nitralin	32.9	274	216	6	21.50	8.50	0.7610	109.44	13.07
Bifenthrin	33.31	181	166	10	12.43	4.14	0.9562	106.05	13.32
Bromopropylate	33.39	183	155	12	25.79	8.60	0.7425	75.06	12.44
Endrin ketone	33.44	215	279	8	19.64	10.55	0.6018	85.30	10.80
Methoxychlor	33.55	227	115	50	23.68	12.56	0.6608	88.07	10.38
Tetradifon	34.1	159	111	20	3.02	1.01	0.9861	102.30	10.86
Leptofos	34.35	171	77	18	20.12	6.71	0.7134	108.60	2.48
Mirex	34.7	272	237	15	15.20	5.07	0.8379	90.21	4.35

164

165



180 Fig. (1) GC-MSMS TQD Chromatogram obtained from 74 pesticides sample

181

182

183

184

## Case Study: Pesticide residues in conventional and organic farming soil.

Pesticide residues ~~was~~<sup>were</sup> detected by applying the modified QuEChERS method: ~~named~~ Chloroneb, Tecnazene, Propachlor, Cycloate, Ethalfluralin, Trifluralin, Beluralin, Sulfotep, Diallate, Alpha BCH, HCB, Atrazine, Terbufos, Profluralin, Fenofos, Diazinon and Fluchloralin and its residue levels ranged from 43.00 to 2.00  $\mu\text{g}/\text{kg}^{-1}$  in organic and conventional olive farm soil (Fig. 2) extracted by QuEChERS and analyzed by GCMSMSTQD.

Meanwhile, the detected pesticide residues levels (Fig. 3) of Endosulfan ether, Pentachloroaniline, Alachlor, Vinclozolin, Cypermethrin, Heptachlor, Acetochlor, Fenchlorfos, **Pirimiphos** methyl, Petachlorothioanisole, Dichlofluanid, Aldrin, Chlropyrifos, Triadimefon, **Pirimiphos** ethyl ranged from 1.99 to 1.00  $\mu\text{g}/\text{kg}^{-1}$  in organic and conventional olive farm soil extracted by QuEChERS and analyzed by GCMSMSTQD.

Also, Isopropalin, Isodrin, Pendimethalin, Heptachlor epoxide, Tolyfluanid, Quinalphos, Procymidone, **cis-Chlorodane**, o,p-DDE, Tetrachlorviphos, Endosulfan I, and **chlordan-trans** was detected in organic and conventional olive farm soil extracted by QuEChERS and analyzed by GCMSMSTQD and ranged from 0.99 to 0.50  $\mu\text{g}/\text{kg}^{-1}$  ( Fig. 4).

Pesticide residues ~~of named~~ Chlordane-trans, Nonachlor-trans, Pretialchlor, **p,p-DDE**, Dieldrin, o,p-DDD, Endrin, Chlorobenzilate, Endosulfan II, **p,p-DDD**, **o,p-DDT**, Nonachlor-cis, Endrin aldehyde, Carfetraxone ethyl, Methoxychlor olefin, Endosulfan sulfate, o,pMethoxychlor, Resmethrin 1, Resmethrin 2, Nitratin, Bifenthrin, Bromopropylate, Endrin\_ketone, Methoxychlor, Tetradifon, Leptofos, and Mirex ~~was~~ ranged as low as 0.50  $\mu\text{g}/\text{kg}^{-1}$ .

Formatted: Strikethrough, Highlight

Formatted: Highlight

Formatted: Strikethrough, Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Strikethrough, Highlight

Formatted: Highlight

Formatted: Strikethrough, Highlight




Fig. (2) Pesticide Residues levels ( $\mu\text{g kg}^{-1}$ ) ranged from 43 to 2 ppb in organic and conventional olive farm soil extracted by QuEChERS and analyzed by GCMSMSTQD.





Fig. (3) Pesticide Residues levels ( $\mu\text{g kg}^{-1}$ ) ranged from 2 to 1 ppb in organic and conventional olive farm soil extracted by QuEChERS and analyzed by GCMSMSTQD.



224

225 Fig. (4) Pesticide Residues levels ( $\mu\text{g kg}^{-1}$ ) ranged from 0.5 to 1 ppb in organic and conventional  
226 olive farm soil extracted by QuEChERS and analyzed by GCMSMSTQD.  
227

227

228 **Conclusions**

229

230 The QuEChERS method is becoming increasingly more popular as a new and  
231 robust procedure. QuEChERS-GC/MS/MS multi-residue method described here is a  
232 simple, rapid and accurate approach suitable for the monitoring of 74 pesticide  
233 residues in old, medial and new olive farms which is 25, 15 and 5 years olive farm  
234 soil samples. The proposed methods have been validated lowing a reliable  
235 determination of the selected compounds with high recoveries. The limit of  
236 detection (LOD) and limit of quantification (LOQ) for the analyzed pesticides were  
237 in the range of  $1.01\text{-}13.91 \mu\text{g kg}^{-1}$  and  $3.02\text{-}29.15 \mu\text{g kg}^{-1}$ , respectively. Pesticide  
238 recoveries form soil samples spiked with pesticides ranged from 65.504 to 111.73  
239 %. The proposed method featured good sensitivity, pesticide quantification limits  
240 were low enough, and the precision, expressed as relative standard deviation,  
241 ranged from 0.29 to 13.32%. Pesticide residues was detected by applying the  
242 modified QuEChERS and GCMSMSTQD method levels was ranged from 43.00 to  
243  $2.00 \mu\text{g kg}^{-1}$  for 18 different pesticides, 1.99 to  $1.00 \mu\text{g kg}^{-1}$  for 16 different  
244 pesticides, 0.99 to  $0.50 \mu\text{g kg}^{-1}$  for 12 different pesticides and lower than  $0.50 \mu\text{g}$

kg<sup>-1</sup> for 28 different pesticide residues. QuEChERS provides high quality results with a high sample throughput. Additionally, there is low solvent and glassware consumption, with low work and cost of analysis per sample.

## References

- AOAC Method. Official Methods of Analysis 2007.01, Pesticide Residues in Foods by Acetonitrile Extraction and Partitioning with Magnesium Sulfate. 2007, 18th: Available from: [http://www.weber.hu/PDFs/QuEChERS/AOAC\\_2007\\_01.pdf](http://www.weber.hu/PDFs/QuEChERS/AOAC_2007_01.pdf), accessed December 2012.
- Caldas S.S., Bolzan C.M., Cerqueira M.B., Tomasini D., Furlong E.B., Fagundes C., and Primel E.G., Evaluation of a Modified QuEChERS Extraction of Multiple Classes of Pesticides from a Rice Paddy Soil by LC-APCI-MS/MS. *J. Agric. Food Chem.*, 2011, 59, 11918-11926.
- El-Saeid, Mohamed H., Mohamed T. Selim, Sherif B. Abdel Ghani. (2015) Monitoring of pesticide residues in organic and conventional farmland soils using GC-MS. *Wulfenia J*, Vol 22, No. 2;Feb 2015, 271-283.
- Khan, S. U. 1982. Studies on bound <sup>14</sup>C-prometryn residues in soil and plants. *Chemosphere*, 11(8):771-795.
- Khan, S. U. 1995. Bound pesticides residues in food products. *AgriScience*. December 1994/January 1995.
- Maestroni, B. 2002. Preparation of Samples and Estimation of Uncertainty of Sample Processing. In Lectures/Uncertainty of sample processing. Training Workshop on Introduction to QC/QA measures in Pesticide Analytical Laboratories, Training and Reference Center for Food and Pesticide Control, Seibersdorf, Vienna, Austria, June 17 - July 26.
- Maestroni, B., A. Ghods, M. El-Bidaoui, N. Rathor, O. P. Jarju, T. Ton and A. Ambrus. 2000. Testing the efficiency and uncertainty of sample processing using <sup>14</sup>C-labelled Chlorpyrifos, Part I. In: Fajgelj A, Ambrus A (eds) Principles of method validation. Royal Society of Chemistry, Cambridge, p 49-58.
- Pinto C.G., Laespada M.E.F., Martín S.H., Ferreira A.M.C., Pavón J.L.P., and Cordero B.M., Simplified QuEChERS approach for the extraction of chlorinated compounds from soil samples. *Talanta*, 2010, 81, 385-391.
- Pinto C.G., Martín S.H., Pavón J.L.P., and Cordero B.M., A simplified Quick, Easy, Cheap, Effective, Rugged and Safe approach for the determination of trihalomethanes and

280 benzene, toluene, ethylbenzene and xylenes in soil matrices by fast gas chromatography  
281 with mass spectrometry detection. Anal. Chim. Acta, 2011, 689,129-136.

282 Powley,C. R. 2004. Extraction efficiency considerations for present and future agrochemical  
283 residue methods. 5<sup>th</sup> European Pesticide Residues Workshop (EPRW), Pesticides in Food  
284 and Drink. Book of Abstracts: 43. June 13-16 2004; Stockholm, Sweden.

285 Pszczolińska, K, and Michel, Monika (2016).The QuEChERS Approach for the Determination  
286 of Pesticide Residues in Soil Samples: An Overview. Journal of AOAC International,  
287 99, (6), 1403-1414(12). DOI: <https://doi.org/10.5740/jaoacint.16-0274>

288 Rashid A., Nawaz S., Barker H., Ahmad I., and Ashraf M., Development of a simple extraction  
289 and clean-up procedure for determination of organochlorine pesticides in soil using gas  
290 chromatography-tandem mass spectrometry. J. Chromatogr. A, 2010, 1217, 2933-2939.

291 Wang Y.H., Du L.W., Zhou X.M., Tan H.H., Bai L.Y., Zeng D.Q., and Tian H., QuEChERS  
292 extraction for high performance liquid chromatographic determination of pyrazosulfuron-  
293 ethyl in soils. J. Chem. Soc. Pak., 2012, 34, 28-32.

294 Brondi S.H.G., de Macedo A.N., Vicente G.H.L., and Nogueira A.R.A., Evaluation of the  
295 QuEChERS method and gas chromatography-mass spectrometry for the analysis pesticide  
296 residues in water and sediment. Bull. Environ. Contam. Toxicol., 2011, 86, 18-22.

297 Fernandes V.C., Domingues V.F., Mateus N., and Delerue-Matos C., Multiresidue pesticides  
298 analysis in soils using modified QuEChERS with disposable pipette extraction and  
299 dispersive solid-phase extraction. J. Sep. Sci., 2013, 36, 376-382.