

Original Research Article

ESTIMATION OF SERUM LIPID PROFILE AND ITS CORRELATION WITH ALVEOLAR BONE HEIGHT IN CHRONIC PERIODONTITIS.

Running title SERUM LIPID PROFILE IN CHRONIC PERIODONTITIS

Abstract

Aim: Numerous studies have been previously carried out to find out any correlation between lipid profile and periodontitis but existence of an association does not establish whether periodontitis causes an elevation in serum lipid level or elevation in serum lipid profile predispose to periodontitis. The aim of ~~the present study~~~~present study~~ was to find any significant correlation between HDL, LDL, VLDL and cholesterol with periodontitis. **Material and methods:** 100 patients with chronic periodontitis (group I) and 50 healthy patients (group II) were selected. The levels of serum ~~lipid, including~~~~lipid including~~ HDL, LDL, VLDL, and TOTAL CHOLESTEROL along with fasting blood glucose were assessed. The relationship between severity of periodontitis based on clinical and radiographic finding with serum lipid correlated. **Results:** ~~There were~~ ~~There was~~ no significant difference found between mean values of total cholesterol, LDL and VLDL among study and control group. Only HDL showed ~~a highly significant difference~~~~highly significant difference~~ ($p < 0.00$) between healthy and patient with chronic periodontitis. **Conclusion:** Estimation of serum lipid ~~profile, especially~~~~profile especially~~ HDL levels in subjects with periodontitis can be considered as a screening method for early diagnosis of atherosclerosis to avoid further progression of cardiovascular changes in early age of the life.

Keywords: Chronic periodontitis, HDL, LDL, Atherosclerosis.

23

24

25 **Introduction**

26 Coronary heart disease (CHD) is one of the leading factors of morbidity and mortality
27 throughout the world being responsible for 16% of death in developing and 50% in developed
28 countries. Atherosclerosis of ~~the coronary arterie e is considered~~ coronary arteries is considered to
29 be the leading cause of premature death among men. The pathological background of the
30 atherosclerosis of coronary arteries is formation of atherosclerotic plaque, which additionally
31 induces other cardiovascular diseases.¹

32 Several bacteria and virus have also been identified as potential etiological factors in
33 cardiovascular diseases (CVD).² Periodontitis is an inflammatory disease of the supporting
34 tissues of teeth caused by specific microorganisms resulting in progressive destruction of the
35 periodontal ligament ~~periodontal ligament~~ and alveolar bone with pocket formation, recession or
36 both.³

37 Perhaps one of the most helpful findings ~~helpful finding~~ about periodontitis is that host response
38 varies between individuals and that either an inadequate host immune response or exaggerated
39 response to bacterial pathogens leads to a more severe form of disease.⁴ The majority of risk
40 factors of cardiovascular diseases are also considered to be risk factors for periodontal diseases.¹

41 Thus, individuals with periodontitis are more likely to have increased d in the level of free fatty
42 acids in the form of cholesterol and triglycerides. These fatty acids are also involved in the
43 athermatnous plaque formation in blood vessels. This increase in atheromas has been associated

44 with thrombotic events in the form of various cardiovascular diseases. Recent studies have also
45 shown isolation of [the](#) periodontal pathogen from this atheromatous plaque.⁶

46 Although numerous studies have been previously carried out to find out any correlation between
47 lipid profile and periodontitis but existence of an association does not establish whether
48 periodontitis causes an elevation in serum lipid level or elevation in serum lipid profile
49 predispose to periodontitis. So the present study is carried out to find any significant correlation
50 between HDL, LDL and cholesterol with periodontitis and also to assess whether periodontitis
51 may be the first step in the etiopathogenesis of cardiovascular disorders in patients whom other
52 obvious risk factors like smoking were absent.

53 **Materials & methods**

54 A hospital based cross sectional study was carried out in the Department of Oral
55 Medicine and Radiology, VSPM Dental College and Research Centre, Nagpur. Subjects were
56 apprised of the purpose of the study and written consent was taken prior to commencement of the
57 study. Ethical clearance was obtained from the ethical committee of the institution.

58 This study consists of 100 patients with chronic periodontitis (group I) and 50 healthy
59 patients (group II). Study consists of patients having age more than 20 yrs without any habits
60 along with no systemic or medical illness like hypertension, myocardial infarction, stroke,
61 asthma, endocrinal diseases.

62 Periodontal condition of the two groups were determined on basis of clinical examination of
63 gingiva and its associated structures including color, contour, surface texture, consistency,
64 bleeding on probing, presence of stippling and pus discharge through gingival crevices. The
65 pocket depth was measured bilaterally in the region of premolar molar region of the mandible

66 with the help of Williams graduated probe. Thus depending upon the attachment loss clinically
67 we divided into three groups like, Mild periodontitis: - < 3mm Moderate periodontitis:-3-6 mm
68 ,severe periodontitis : > 6mm

69 In addition to clinical examination we also carried out radiographic investigation to
70 assess the alveolar bone height with the help of Digora (optime) RVG software. Digital
71 calibrations were made for measurement of alveolar bone height from CEJ upto the bone level in
72 premolar molar region of both the quadrant in mandible.

73 For estimation of lipid profile like total cholesterol, HDL, LDL and VLDL we advised
74 patient to fast at least 8-12 hrs and collected 2 ml of blood sample from antecubital vein by
75 venipuncture method and stored in collecting test tubes with anticoagulant and sent to pathology
76 laboratory for enzymatic analysis.

77 Statistical analysis was done to evaluate the correlation between serum lipid profile and
78 periodontitis along with radiographic bone height secondary to periodontitis to find out the
79 significant difference between these values with the help of chi square, annova and Z test.

80 **Results:**

81 The mean values of Total cholesterol was 171.26 mg/dl, while in control group it was 161.51
82 mg/ dl, and the mean triglyceride in study group was 115.20 mg/dl, while in control group it was
83 99.91mg/dl, the mean LDL cholesterol in study group was 106.40 mg/ dl, while in control group
84 it was 98.59 mg/ dl. The mean of VLDL cholesterol in study group was 23.17mg/ dl, while in
85 control group it was 20.02 mg/ dl. The mean fasting blood glucose level in study group was
86 99.90 mgm/ dl, while in control it was 92.27 mgm/ dl. But there was no statistical significant
87 difference found in Total cholesterol, triglycerides, LDL, VLDL & Fasting blood glucose.

88 Further mean total HDL in study group was 41.20 mg/ dl, while in control group it was
89 42.97mg/dl which found to be statistically significant (P=0.04).

90 Further correlation was done for ~~of~~ Alveolar bone loss in mandibular left & right premolar molar
91 region with Lipid parameters. Among all lipid parameters in mandibular left premolar region,
92 HDL showed negative correlation ($r = - 0.209$) with statistical significant difference (P=0.036),
93 whereas other parameters (LDL, VLDL & triglyceride) showed no significant difference.
94 Further alveolar bone loss relation with lipid parameters in mandibular right premolar region
95 found to be negative and was not significant. In the present study we observed raised levels of
96 lipid parameters in both the groups; among all parameters HDL showed highly significant
97 differences (p-0.0001).

98

99 **Discussion**

100 Periodontitis has been traditionally regarded as a chronic inflammatory oral infection
101 which mainly consists of gram negative anaerobic microflora that leads to gingival
102 inflammation, destruction of periodontal tissues, loss of alveolar bone and exfoliation of the
103 teeth. It is generally accepted that certain organisms within the microbial flora of dental plaque
104 are major etiological agent in periodontitis. These microorganisms particularly P. Gingivalis
105 produces endotoxins in the form of lipopolysacchrides that generates a host mediated tissue
106 destructive immune response. Traditionally it is thought that periodontitis is an oral disease and
107 that the tissue destructive response remains localized within the periodontium, limiting effects of
108 the disease to oral tissues supporting teeth, however recent studies indicate that oral disease may
109 have profound effect on systemic health.⁴⁶

110 A number of studies have reported association between periodontitis and cardiovascular
111 diseases. Most of the risk factors for cardiovascular diseases are also regarded as risk factor for
112 periodontal diseases. Some studies have found no relationship or an inverse relationship between
113 chronic infection and hyperlipidemia.

114 Chronic infections like periodontitis have been demonstrated to induce profound changes
115 in plasma concentration of cytokines like TNF- alpha and IL- beta which can result into elevated
116 levels of free fatty acids, LDL (low density lipoprotein) and triglycerides. These elevations in
117 serum lipids are thought to arise from enhanced hepatic lipogenesis, increased adipose tissue
118 lipolysis, increased synthesis or reduced clearance of LDL due to reduction in lipoprotein lipase
119 activity.⁴⁷

120 Therefore the present study was carried to find out any correlation between chronic infection like
121 periodontitis and increase in serum lipid level in form of cholesterol, triglyceride, HDL (high
122 density lipoprotein), LDL (low density lipoprotein) in otherwise healthy subjects and who did
123 not have any habits

124 In the present study 8% subjects showed increased total cholesterol from study group whereas it
125 was 4% among healthy subjects; this increased TC showed statically significant difference (p
126 <0.05). similar results were also seen by Loesche et al (2000)²¹ & Taleghani F (2005)³⁴

127 These findings give evidence to the theory that periodontitis may be one of the factor that is
128 responsible for increase in total cholesterol levels as in our study the patients did not have any
129 other systemic disorders and no smoking habit, but at the same time other factors such as
130 physical activity, nutrition, stress, socioeconomic status and body mass index (BMI) might have
131 some influence on total cholesterol level which should also be evaluated.

132 Increased triglyceride levels when assessed between study and control group we found three fold
133 increase in subjects with periodontitis but statically it was not significant, which was also seen
134 with Taleghani F (2005)³⁴.

135 In contrast to our study and Taleghani F, Loesche et al (2000)²⁴ found significant correlation
136 when triglyceride levels were assessed. This increase in plasma triglyceride levels could be due
137 to increase in pro-inflammatory cytokines in response to chronic periodontitis. Infection with
138 Gram negative periodontal pathogen can cause rapid release of systemic IL-1 beta and TNF-
139 alpha which are responsible for hyper-triglyceridaemia.

140 When decreased levels of HDL were assessed in both the groups; the frequency of low
141 HDL was higher by 12% in chronic periodontitis as compared to control; with statistical
142 significant difference ($p < 0.0001$). whereas Loesche et al (2000)²⁴, Taleghani F (2005)³⁴, Cristana
143 A (2005)³⁰ did not find any statistical significant difference for HDL.

144 The explanation for relationship between low HDL and periodontitis might be chronic
145 infection in the periodontitis that leads to release of lipopolysaccharide and proinflammatory
146 cytokines.

147 On the other hand HDL also has anti-inflammatory properties that can decrease the
148 adhesion of endothelial cells, thus low plasma concentration of HDL in blood may be a
149 contributory factor to inflammatory process in periodontitis. In this study no statically significant
150 difference was found when for LDL & VLDL. Loesche et al (2000)²⁴ found significantly raised
151 LDL levels in subjects with periodontitis which is contradictory to present study. Thus a cause
152 and effect relationship between HDL and periodontitis needs to be established.

153 Raised blood glucose levels were found in 35% from study group while it was 24% in
154 control group with statistical significant difference (p value < 0.05). Liu et al (1998)⁵⁰, Reimers
155 et al (1998)⁵¹ & Shiba et al (1998)⁵² stated that some cytokines such as TNF- alpha and IL-beta
156 that are produced in response to infection with gram negative bacteria may be responsible for
157 insulin resistance and subsequent poor glyceamic control in periodontitis patients.

158 In our study among all lipid parameters only HDL showed negative correlation with
159 alveolar bone loss and severity of periodontitis which was statistically significant indicating that
160 increase in amount of bone loss in severe periodontitis is associated with low serum level of
161 HDL. Similar results are also seen with Saito T (2004)²⁴²

162 In the present study among all lipid parameters HDL was the most significantly associated with
163 periodontitis. There was significant difference found in plasma levels of HDL in subjects with
164 periodontitis as compared to subjects with healthy periodontium.

165 Unlike previous studies we did not find any statistical difference in total cholesterol,
166 triglyceride and LDL levels in subjects with periodontitis and without it. In the previous studies
167 it has been shown that abnormal levels of total cholesterol and LDL are indicators of
168 atherosclerosis or coronary heart disorders, but HDL can be a better measure especially in
169 individuals less than 60 yrs.

170 **Conclusion**

171 The present study mainly consisted of subjects in third and fourth decades with mean age
172 of 39.2 yrs. HDL levels were found to be statistically significantly lowered in the study group
173 that is in subjects with periodontitis. The abnormal HDL levels also statistically significant with
174 severe form of periodontitis thus it can be concluded that young individuals which are affected

175 with any form of periodontitis should be evaluated for the plasma levels HDL as an early marker
176 for atherogenic lipid profile.

177

178

179 **References**

- 180 1. Valentaviciene G, Paipaliene P, Nedzelskiene I. The relationship between blood serum
181 lipids and periodontal condition. *Stomatologia, Baltic Dental and Maxillofacial Journal*
182 2006; 8(3):96-100.
- 183 2. Saito T, Murakami M, Shimazaki Y. Association between alveolar bone loss and elevated
184 serum C reactive protein in Japanese men. *J Periodontol* 2003; 74:1741-1746.
- 185 3. Newman M, Takei H, et.al. Carranza Clinical Periodontology 10th edition, Elsevier
186 Saunders publication, 2007.
- 187 4. Sanz M, Deanfield J. European workshop in periodontal health and cardiovascular
188 disease- scientific evidence on association between periodontal and cardiovascular
189 diseases: A review of literature. *European Heart Journal Supplement* 2010;12
190 (supplement B), B3-B1
- 191 5. Kauppinen- Makelin R, Nikkila E. Serum lipoproteins in patients with myocardial
192 infarction. *Atherosclerosis* 1998;74:665-674.
- 193 6. DeStefano F, Anda R, Khan H. Dental disease and risk of coronary heart disease and
194 mortality. *Br Dent J* 1993; 306:688-691.

- 195 7. Van der Poll T, Saurwein H. Tumor necrosis factor- alpha: its role in the metabolic
196 response in sepsis. Clin Sci 1993; 84:247-256.
- 197 8. Loesche W, Karapetow F, Pohl C. Plasma lipid and blood glucose level in patients with
198 destructive periodontal diseases. J Clin Periodontol 2000; 27: 537-541.
- 199 9. Taleghani F, Shamaei M, Shamaei M. Association between chronic periodontitis and
200 serum lipid levels. Acta Medica Iranica 2010; 48(1): 47-50.
- 201 10. Cristana A, Rozeli M et al. Relation between chronic periodontal disease and plasmatic
202 levels of triglycerides, total cholesterol and fractions. Braz Oral Res 2005; 19(4):284-289.
- 203 11. Liu L, Spelleken M, Röhrig H. Tumor necrosis factor-alpha acutely inhibits insulin
204 signaling in human adipocytes: Implication of the p80 tumor necrosis factor receptor.
205 Diabetes 1998; 47 (4): 515-522.
- 206 12. Reimers J. Interleukin-1 β induced transient diabetes mellitus in rats. A model of the
207 initial events in the pathogenesis of insulin dependant diabetes mellitus? Danish Medical
208 Bulletin.1998; 45: 157-180.
- 209 13. Shiba T, Higashi N. Hyperglycemia due to insulin resistance caused by interferon γ .
210 Diabetic medicine 1998; 15: 435-436.
- 211 14. Saito T, Murakami M, Shimazaki Y. Association between alveolar bone loss and elevated
212 serum c reactive protein in Japanese men. J Periodontol 2003; 74: 1741-1746.

213 **Table 1: Prevalence of serum lipid level in Study and control groups**

214

	TC	TG	HDL	LDL	VLDL	TC:HDL	LDL:HDL
Study	8	6	74	7	6	19	17
Control	2	1	31	1	1	3	3
χ ² -value	10.00	2.91	60.39	8.00	2.91	3.03	5.07
p-value	0.00	0.23	0.00	0.01	0.31	0.21	0.07

215 P<0.05= significant

216 **Table 2: Correlation of radiographic bone loss (35, 36 & 37 region) with lipid parameters**

217

Lipid Profiles	35 – 36 region		36 – 37 region	
	Correlation ‘r’	p-value	Correlation ‘r’	p-value
Blood Glucose	0.000	0.997	-0.010	0.919
TC	0.021	0.835	-0.059	0.561
TG	0.065	0.523	0.065	0.520
HDL	-0.209	0.036	-0.191	0.057
LDL	0.041	0.686	-0.045	0.658
VLDL	0.062	0.541	0.059	0.558
TC/HDL	0.082	0.415	-0.009	0.930
LDL/HDL	0.154	0.125	0.062	0.542

218 P<0.05=significant

219 **Table 3: Comparison of mean lipid parameters in Study and Control group**

220

(a) Descriptive Statistics

221

Sr. no.	Parameter	Group	N	Mean	Std. Deviation	Std. Error Mean
1	TC	Study	100	171.26	42.57	4.25

		Control	50	161.51	33.22	4.69
2	TG	Study	100	115.20	53.38	5.33
		Control	50	99.91	46.88	6.63
3	HDL	Study	100	41.20	5.65	0.56
		Control	50	42.97	4.84	0.68
4	LDL	Study	100	106.40	39.69	3.96
		Control	50	98.59	30.88	4.36
5	VLDL	Study	100	23.17	10.42	1.04
		Control	50	20.02	9.39	1.32
6	TC/HDL	Study	100	4.10	1.08	0.10
		Control	50	3.69	0.81	0.11
7	LDL/HDL	Study	100	2.61	0.98	0.09
		Control	50	2.31	0.75	0.10

222

223 **Table 4: Prevalence of abnormal serum lipid level in Study and control groups**

	TC	TG	HDL	LDL	VLDL	TC/HDL	LDL/ HDL
Study	8	6	74	7	6	19	17
Control	2	1	31	1	1	3	3
χ^2 -value	10.00	2.91	60.39	8.00	2.91	3.03	5.07
p-value	0.006 S,p<0.05	0.23 NS,p>0.05	P<0.0001 significant	0.018 S,p<0.05	0.31 NS,p>0.05	0.21 NS,p>0.05	0.07 NS,p>0.05

224

225 **Table 5: Correlation of radiographic bone loss (35, 36, 37) with lipid parameters in study**
 226 **group**

227

Pearsons Correlation Coefficient

228


Lipid Profiles	35-36		36-37	
	Correlation 'r'	p-value	Correlation 'r'	p-value
Blood Glucose	0.000	0.997 NS,p>0.05	-0.010	0.919 NS,p>0.05
TC	0.021	0.835 NS,p>0.05	-0.059	0.561 NS,p>0.05
TG	0.065	0.523 NS,p>0.05	0.065	0.520 NS,p>0.05
HDL	-0.209	0.036 S,p<0.05	-0.191	0.057 NS,p>0.05
LDL	0.041	0.686 NS,p>0.05	-0.045	0.658 NS,p>0.05
VLDL	0.062	0.541 NS,p>0.05	0.059	0.558 NS,p>0.05
TC/HDL	0.082	0.415 NS,p>0.05	-0.009	0.930 NS,p>0.05
LDL/HDL	0.154	0.125 NS,p>0.05	0.062	0.542 NS,p>0.05

229

230 **Graph 1-A: Prevalence of abnormal serum lipid level in**

231 **Study and control groups**

232



233