

Allelopathic Effect of *Eucalyptus globulus* on Seed Germination and Seedling Growth of Highland Teff (*Eragrostis tef*) and Barely (*Hordeum vulgare* L.)

ABSTRACT

The phenomenon of allelopathy, whereby a plant species chemically interferes with the germination, growth or development of other plant species has been known and documented for over 2000 years. Allelochemicals are secondary metabolites of plants, released into environment through volatilization, leaching, root exudation and decomposition of residues. This study was aimed to examine the allelopathic effect of *Eucalyptus globulus* on seed germination and early growth of highland teff (*Eragrostis tef*) and barely (*Hordeum vulgare* L.) conducted at Kulumsa Agricultural Research Center. Fresh leaf, juvenile stems, root and mixtures of these aqueous extracts at 0%, 5%, 10%, 15%, 25% and 50% concentrations respectively were used to run the experiment under Laboratory condition. The employed research design was Complete Randomized Design with three replications and two factorials as: plant parts and concentrations. The quantitative data was collected, coded and then subjected to SAS Version 9.1 procedure following two ways ANOVA. The means were compared by using least significance difference test (LSD) at $P \leq 0.05$ probability level. It was noted that aqueous extracts at a concentration of 10%, 15%, 25% and 50% had inhibitory effect on both crops seed germination and seedling growth. The effect was found in much higher than the control treatment. The inhibitory effects were increased as the extract concentration increased and more pronounced in fresh leaf aqueous extract compared to the rest extract sources. These finding indicates allelochemicals in the *Eucalyptus globulus* plant parts adversely affect seed germination and early seedling growth of teff and barely. The Phytotoxicity of *Eucalyptus globulus* fresh leaf aqueous extract showed the highest impact affecting early root growth of barely and shoot growth of teff under the highest level of concentration (50%) of the extracts.

Key Words: Allelopathic, Barely, *Eucalyptus globulus*, Highland teff, Seed Germination, Seedling Growth

INTRODUCTION

The term allelopathy is derived from the Greek-compound words allelo and pathy (meaning “mutual harm” or “suffering”) and was first used in 1937 by Austrian scientist Hans Molisch [1-2]. Allelopathy is defined as the beneficial or harmful influence of chemical substances released by plants that can alter the growth and development of nearby plants or microorganisms [3]. The phenomenon of allelopathy, whereby a plant species chemically interferes with the germination, early growth or development of other plant species has been known and documented for over 2000 years [4]. Allelochemicals or phytochemicals are plant secondary metabolites normally released into the environment through volatilization, leaching root exudation and decomposition of plant residues in the soil [5]. The action of allelochemicals can affect the respiration, photosynthesis, enzyme activity, water relations, stomatal opening, hormone levels, mineral availability, cell division and elongation, and structure and permeability of cell membranes and walls [6-9].

Discharge of allelochemicals into the environment occurs by exudation of volatile chemicals from living plant parts, by leaching of water soluble toxins from aboveground parts in response of action of rain, by exudation of water soluble toxins from below ground parts, by release of toxins from non-living plant parts through leaching of litter decomposition. Many invasive plant species alter natural ecosystems and reduce plant diversity causing plant displacement by allelopathic inhibition of germination or growth via phytotoxic chemical release [9-10]. *Eucalyptus globulus* one of such invasive plant species posing greater challenges to the economic, food security and sustainable development of many developing countries whose livelihood is of totally or partially depend on agriculture [11].

Germination and seedling growth are the screening criteria which are widely used to investigate the effects of allelopathy. Morphological changes, in response to allelochemicals, could be due to effects on cellular or molecular level [12]. Allelochemicals may be present in the leaves, barks, roots, flowers and fruits. Therefore, leaves, juvenile stem, root and mixture of all these parts were selected to accomplish the extract bioassays in this research. Allelochemicals restrict plant growth through negative interactions with some physiological processes

57 such as suppression of cell division, changes in cell wall structure and activity of some enzymes. The effect of
58 allelochemicals action was detected at molecular, structural, physiological, biochemical and ecological levels of
59 plant organization [9].

60
61 Many studies have evaluated the allelopathic activity of *Eucalyptus* species and reported strong inhibitory
62 effects of eucalyptus extracts on germination and growth of various plant crops such as cucumber, sorghum,
63 rice, tomato, eggplant and black gram [13-16]. However, there is no research finding indication on allelopathic
64 effect of *Eucalyptus globules* on highland teff and barely. Therefore, this study is aimed to examine the
65 allelopathic effect of *Eucalyptus globulus* extracts from fresh leaves, juvenile stem, root exudates and mixture
66 extracts of these plant parts on seed germination as well as seedling growth of barely and highland teff.

67 MATERIALS AND METHODS

68 Description of the Study Site

69 The study was conducted at Kulumsa Agricultural Research Center (KARC), central Ethiopia. It is relatively
70 located to the North of Assela town (167 km from Addis Ababa), to the left side of main road from Addis Ababa
71 to Assela. Kulumsa Agricultural Research Center was established in 1966 by government of Ethiopia and the
72 Swedish International Development Agency (SIDA). The research Center is mandated to wheat, malt barley and
73 highland pulse crops research nationally and serves as Wheat Center of Excellence for East Africa (Ethiopia,
74 Kenya, Uganda, Tanzania), regionally. KARC is situated at latitude 08°01'10" N and longitude 39°09'11" E
75 with an annual average maximum and minimum temperature of 23.2°C and 10.5°C respectively, altitude of 2200m.a.s.l.
76 and an average annual rain fall of 823 mm. Soil type clay soil (Luvisols) and soil pH 6.0. Research Center
77 carries out research on crop production, animal breeding, on soil and nutritional issues [17].

80 Experimental Materials

81 *Eucalyptus globulus* plant parts such as fresh leaves; juvenile stem and root were collected from the farmers'
82 farm land around KARC and rinsed thoroughly with sterile distilled water [18]. Seeds of most popular highland
83 teff of variety Felagot (Dz. Cr.442) collected from Bishoftu Agricultural Research Center and seeds of barely
84 variety called Holker were collected from KARC. Bare land local top soil was collected from depth of 0-30cm
85 as of [19] from KARC.

87 Research Procedure

88 Fresh leaves and juvenile stem collected by cutting and root collected through digging and cutting from a tree of
89 10 years old, purposively the mid age of harvesting [20] of *Eucalyptus globulus* and chopped in to pieces, open
90 air dried under shade for fifteen days; the samples then put in oven with a temperature of 50°C for 48 hours [21].
91 All the samples collected prepared powder to fine particles using electric grinder separately. The grinded plant
92 materials were mixed in distilled water at (5%, 10%, 15%, 25%, and 50%) w/v ratio separately [22]. The
93 mixtures then soaked for about 12 hours at room temperature and blended with blender for 24 hours. Then after,
94 the mixtures were centrifuged at 100 × 100 rpm for twenty minutes and the aqueous solutions of each plant part
95 was filtered using What man filter paper Number 1 poured into flat bottomed 250 ml volumetric flasks, well
96 covered and preserved in refrigerator set to -5°C until it was used in a test experiment [10]. For the preparation
97 of mixture extract of the three plant parts, 30ml of each of plant parts with the respective concentrations (5%,
98 10%, 15%, 25%, 50%) were shared equally mixed well and poured into flat bottomed 250 ml volumetric flasks.

100 Bare land local top soil from depth of 0 - 30cm as of [19] collected from three different quadrant of (2x2) meter
101 square from KARC, then mixed well and then equal amount filled to every plastic pot having 3.5 inch depth, 3.5
102 inch top and 3 inch bottom diameters provided with drainage holes; moisten with 80 ml of tap water and left for
103 one day. Seeds of both crops with germination percentage of 85% and purity of 99% were thoroughly washed
104 with distilled water and surface sterilized with 90% ethanol for 2 minutes, followed by soaking for 5 minutes in
105 a solution of 5.25% sodium hypochlorite, then rinsed four times with distilled water [10].

106 Research Design

107 The research has two laboratory based experimental parts. **Experiment I:** For seed germination of teff and
108 barely. **Experiment II:** For seed germination and early seedling growth of both crops. Both experiments were
109 laid down in factorial Randomized Complete Design (RCD), with three replications. The research has also two
110 factors namely plant parts as sources of extract as Factor A: at four levels; juvenile stem (A1), root(A2), fresh
111 leaf (A3) and mixtures of (juvenile stem, root & fresh leaf) (A4) and extract concentration levels as Factor B: at
112 six levels; 0%w/v (B1), 5%w/v (B2), 10% w/v (B3), 15% w/v (B4) 25% w/v (B5) and 50% w/v (B6) were used
113 as of [10].

114
115 **Experiment I:** for clearly monitoring seed germination of both crops, nine seeds of both crops were sown
116 sparsely in a filter paper covered glass Petri dish having 9 cm diameter [23] and 5 ml aqueous extract of

117 eucalyptus plant parts at 5%, 10%, 15%, 25%, and 50%) ratio were applied to each glass Petri dish separately
 118 and 5 ml distilled water applied in the case of control treatment [18]. The seed planted Petri dishes were placed
 119 in green house and arranged in rows for both crops independently and fairly labeled. Seed germination data for
 120 both crops collected on daily basis after planting [22]. The experiment was continued for ten days for seed
 121 germination test and 21 days for seedling growth.

122

123 **Experiment II:** For seed germination and early seedling growth of both crops uniform pots filled with local top
 124 soil from the depth of 0-30 ml prepared were arranged in rows for both crops independently. Nine seeds for both
 125 crops were planted separately in each of the plastic pots and 5 ml aqueous extract of eucalyptus plant parts at
 126 5%, 10%, 15%, 25%, and 50%) ratio were applied to each pot separately and 5 ml distilled water applied in the
 127 case of control treatment [10, 23]. The seed planted pots were placed in green house and fairly labeled.
 128 Moistening seeds with equal amount of tap water and data on seed germination for both crops were collected on
 129 daily basis after planting. After emergence, seedlings thinned to six plants per pot for both crops and the
 130 experiment continued for 21 days in which the final measurement was recorded.

131

132 **Data Collection**

133 Seed germination was considered when radicle emergence (≥ 1 mm) and daily counted for 5 days or until the
 134 last seed germinated [24]. Final data collection for early seedling growth was effective after 21 days of seed
 135 plantation and the data from three randomly selected seedlings of both crop species root and shoot length for
 136 each replicate measured and recorded as of [19]. Shoot length (S.L) measured in cm taken from the stem joint to
 137 the tip of the terminal leaf and Root length (R.L) measured in cm taken from the root joint to the tip of the
 138 tap/main root and data recorded and registered clearly.

139

140 **Statistical data analysis**

141 The quantitative data was collected, recorded clearly, coded and then subjected to two way ANOVA analysis of
 142 variance procedure with SAS Version 9.2 and the means compared by using least Significance difference (LSD)
 143 test at the $P \leq 0.05$ level of probability. The seed germination data recorded and seed germination percentage
 144 (%) was determined using: the formula as of [10, 19, 24-25].

145

$$\text{Germination \%} = \frac{\text{Number of germinated seeds}}{\text{Total number of seeds sown}} \times 100$$

146

147 Percentage phytotoxicity produced by *E.globulus* was by the following formula as used in the research done by
 148 [10].

149

$$\text{phytotoxicity \%} = \frac{\text{radicle length of control} - \text{radicle length of treated sample}}{\text{Radicle length of control}} \times 100$$

150

151 **RESULTS AND DISCUSSION**

152 **Seed germination**

153 The present study provides the evidence that aqueous extracts of *Eucalyptus globulus* plant parts such as
 154 juvenile stem, root, fresh leaf or mixtures of them showed the inhibitory response towards germination
 155 percentage and early seedling growth (shoot length and root length) of teff and barely seeds. Interaction in all
 156 conditions showed that allelopathic effect becomes more pronounced with increasing doses of *Eucalyptus*
 157 *aqueous* extracts. The same result was reported by the study of [10] on seed germination and growth of tomato
 158 plant. Even if seed germination count continued for ten days, there was no seed germination, observed starting
 159 from the seventh day to the last tenth date for both crop species and both on Petri dish and in pot conditions.

160 There was direct relationship between allelopathic effect and extract levels of *Eucalyptus globulus* plant parts.
 161 The seed germination of both crops was delayed for aqueous extract treatments at increasing concentration. At
 162 25% and 50% *Eucalyptus globulus* aqueous extracts of all plant parts, seed germination of both crop species
 163 (Teff & Barely) were inhibited under both condition of germinations (germination on Petri dish & in pot).
 164 However, inhibition of germination becomes more pronounced on Petri dish of both crop species. At 50% level
 165 of all aqueous extracts of *Eucalyptus globulus* plant parts of both crop species resulted in a significant reduction
 166 of germination counts when compared with control. In comparison to all conditions of germination aqueous
 167 extract sources of *Eucalyptus globulus* plant parts, fresh leaf aqueous extract exhibited more germination
 168 inhibition at 25% and 50% level of concentration compared to other extract sources. For example: comparing
 169 *Eucalyptus globulus* plant parts aqueous extracts, it was fresh leaf extract at 50% that highly reduced the
 170 germination percentage of Teff in which only 14.4% of seeds were germinated followed by root extract in which

171 only 18.5% of the same seed germination took place. The highest percentage value of teff seed germination
 172 (36.6%) was obtained by the juvenile stem part of the tree at 50%, Table 1.

173

174 Table 1: Percentage of teff seeds germination after treated with *E. globulus* plant part extracts under Petri dish

Source of extract	Treatment level in (%)					
	0% (Control)	5%	10%	15%	25%	50%
Juvenile stem	100	96.3	88.8	66.6	44.4	36.6
Root	96.3	92.2	81.1	58.8	40.7	18.5
Fresh leaf	100	85.2	66.6	51.8	36.4	14.4
Mixtures*	92.2	88.8	85.2	58.5	40.7	22.2

175 *= mixture of juvenile stem, root & fresh leaf each 33%

176

177 The result of analysis of variance, ANOVA is also presented in Table 2. Significant differences among the mean
 178 values were found based on least significance difference, LSD, test. Seed germination of teff was significantly
 179 (at $p \leq 0.05$) responded to the increasing aqueous extracts concentrations for all *Eucalyptus globules* plant parts
 180 both in petri dish and on pot conditions, Table2&3.

181

182 Table 2: Mean comparison of teff seeds germinated per Petri dish after treated by *Eucalyptus globulus* extract.

Treatment concentration level in (%)	Source of extract			
	Juvenile stem	Root	Fresh leaf	Mixtures*
0% (Control)	9.00 ^a	8.67 ^a	9.00 ^a	8.33 ^a
5%	8.67 ^a	8.33 ^a	7.67 ^{ab}	8.00 ^a
10%	8.00 ^a	7.31 ^{ab}	6.00 ^b	7.67 ^{ab}
15%	6.00 ^b	5.33 ^b	4.67 ^{bc}	5.31 ^b
25%	4.00 ^{bc}	3.67 ^c	3.33 ^c	3.67 ^c
50%	3.33 ^c	1.67 ^c	1.32 ^c	2.00 ^c
Mean	6.51	5.82	5.32	5.82
CV%	37.53	54.44	58.03	39.31
LSD	4.32	3.86	4.56	3.86

183

184 Note: means with the same letter in the same column are not statistically significant different at alpha level of
 185 0.05 as evaluated by ANOVA. Significant at $p \leq 0.05$, *= mixture of juvenile stem, root & fresh leaf each 33%,
 186 CV%= coefficient variation in percentage, LSD= least significant difference

187

188 The lowest mean value of teff seeds germination (1.32) in the petri dish, was observed under 50% concentration
 189 of fresh leaf extracts (Table 2), whereas on the pot it was stem juvenile extract the least mean value of
 190 germination, 3.33, (Table 3) obtained for the maximum concentration. There was direct relationship between
 191 seed germination of both crop species and allelopathic effects of extract level. The seed germination of both
 192 crops was delayed for aqueous extract treatments at increasing concentration. Inhibition of teff seed germination
 193 more pronounced under fresh leaf extract at 50% level of concentration, followed by root aqueous extract under
 194 the same level of concentration .Where about 1.3 and 1.67 teff seeds germinated per Petri dish for fresh leaf and
 195 root aqueous extract respectively (Table 2).

196

197 From Table 3, it can be understood that the most allelopathic effect was exhibited at highest level of
 198 concentration (50%) of *Eucalyptus globulus* on teff seed germination under pot condition. Comparing the
 199 extract sources under pot condition, juvenile stem aqueous extract at 50% treatment showed the ultimate
 200 allelopathic effect on teff seed germination where only 3.33 germinated seeds were obtained as least mean value
 201 of the three replications.

202

203

204 Table 3: Mean comparison of teff seeds germinated per pot after treated by *Eucalyptus globulus* extract

	Source of extract			
	Juvenile stem	root	Fresh leaf	Mixtures)
0%(control)	8.67 ^a	9.00 ^a	9.00 ^a	8.67 ^a
5%	8.67 ^a	8.67 ^a	8.32 ^{ab}	8.67 ^a
10%	8.00 ^a	7.67 ^{ab}	7.31 ^{ab}	8.00 ^a
15%	6.67 ^b	6.67 ^b	6.3 ^b	6.67 ^b

25%	6.01 ^b	5.31 ^b	5.00 ^b	5.67 ^b
50%	3.33 ^c	4.33 ^{bc}	4.00 ^c	4.31 ^c
Mean	6.89	6.94	6.65	6.99
CV%	66.64	27.72	33.50	61.32
LSD	3.47	5.06	9.03	11.21

205 Note: means with the same letter in the same column are not statistically significant different at alpha level of
 206 0.05 as evaluated by ANOVA test. Significant at $p \leq 0.05$, *= mixture of juvenile stem, root & fresh leaf each
 207 33%, CV%= coefficient variation in percentage, LSD= least significant difference

208

209

Table 4: Mean comparison of barely seeds germinated per Petri dish after treated by *Eucalyptus globulus* extract

Treatment concentration level in (%)	Source of extract			
	Juvenile stem	root	Fresh leaf	Mixtures
0%	8.67 ^a	9.00 ^a	8.67 ^a	8.67 ^a
5%	8.33 ^a	8.33 ^a	8.10 ^a	8.33 ^a
10%	7.67 ^{ab}	7.33 ^{ab}	7.00 ^{ab}	8.33 ^a
15%	7 ^{ab}	7 ^b	6 ^b	7.3 ^{ab}
25%	5.67 ^b	5.67 ^{bc}	5 ^b	5.3 ^b
50%	3.33 ^c	4.67 ^b	2.67 ^c	3.3 ^c
Mean	6.76	6.99	6.22	6.86
CV%	66.26	52.07	61.06	59.86
LSD	4.01	6.13	4.81	4.37

210

211

212

213

214

Note: means with the same letter in the same column are not statistically significant different at alpha level of 0.05 and 0.01 as evaluated by ANOVA test. Significant at $p \leq 0.05$ *= mixture of juvenile stem, root & fresh leaf each 33%, CV%= coefficient variation in percentage, LSD= least significant difference

215

216

217

218

219

220

221

Table 5: Mean comparison of barely seeds germinated per pot after treated by *Eucalyptus globulus* extract

Treatment concentration level in (%)	Source of extract			
	Juvenile stem	root	Fresh leaf	Mixture
0%	9.00 ^a	8.67 ^a	8.67 ^a	9.00 ^a
5%	8.67 ^a	7.67 ^{ab}	8.67 ^a	8.68 ^a
10%	8.00 ^{ab}	7.30 ^{ab}	8.00 ^{ab}	7.32 ^b
15%	7.00 ^b	7.31 ^{ab}	6.31 ^b	6.30 ^{bc}
25%	6.33 ^{bc}	5.30 ^b	4.67 ^{bc}	5.00 ^b
50%	4.31 ^c	4.00	3.30 ^c	3.67 ^c
Mean	7.21	6.70	6.60	6.66
CV%	24.17	29.41	33.04	37.46
LSD	1.86	3.40	3.87	4.33

222

223

224

225

Note: means with the same letter in the same column are not statistically significant different at alpha level of 0.05 as evaluated by ANOVA test. Significant at $p \leq 0.05$, *= mixture of juvenile stem, root & fresh leaf each 33%, CV%= coefficient variation in percentage, LSD= least significant difference.

226

227

228

229

230

231


232

233

Shoot length and Root length

The allelopathic potential of eucalyptus plant part aqueous extracts on the inhibition of both shoot and root lengths of the target crop species increased gradually with the increase of the different extracts concentration. The intensity of root length reduction increases with the increasing concentrations of aqueous extracts of all extract sources as that of shoot length. Significant reduction in root length was recorded at all concentrations compared to control. The length of root values varied from 1.6cm to 5.2cm for Teff and 2cm to 10.1cm for Barely. The minimum values 1.6 cm and 2cm were observed for 50% treatment of fresh leaf for teff and barely respectively (Figure 2&4).


234 Teff seedling root length inhibition was more pronounced under 50% of all aqueous extract of *Eucalyptus*
 235 *globulus* plant parts; however more teff seedling root length inhibition noticed under fresh leaf aqueous extract
 236 at 50% level of extract treatment. Where only about 1.6cm teff root length recorded compared to control
 237 treatment that accounts 5.3cm in length, Figure 2. Several studies of [10, 15] on (sorghum, rice, and black
 238 gram), tomato and eggplant had evaluated the allelopathic activity of *Eucalyptus* species and reported strong
 239 inhibitory effects of its extracts on germination and growth of various crop plants.
 240



241
 242
 243
 244
 245

Figure 1: Effect of *Eucalyptus globulus* plant parts aqueous extract on shoot length of Teff


246 Shoot length of both crops significantly responded to aqueous extracts of *Eucalyptus globulus* fresh leaf,
 247 juvenile stem, root and mixtures of them (fresh leaf, juvenile stem & root). However, the response depends on
 248 the concentrations and plant parts considered in this particular study. For instance, teff seedling shoot length
 249 inhibition is directly proportional to the increase in concentration of all *Eucalyptus globulus* plant parts juvenile
 250 stem, root, fresh leaf and Mixtures of aqueous extract. As it can be observed on Figure 1, shoot length of teff
 251 seedling inhibition was more pronounced under the treatment mixtures (juvenile stem, root & fresh leaf) of
 252 extract at 50% level of treatment exhibited, which accounts 8.5 cm (the lowest mean value) compared to control
 253 teff seedling shoot length of 18.5 cm long.
 254
 255



256
 257
 258
 259
 260
 261
 262
 263


Figure 2: Effect of *Eucalyptus globulus* plant parts aqueous extract on root length of Teff

The intensity of seedling shoot length reduction increases with the increasing concentration of aqueous extracts of all extract sources. The study indicated that the concentration of all plant parts fresh leaf, juvenile stem, root and mixtures of (fresh leaf, juvenile stem & root) starting from the lowest level of concentration inhibited shoot length of both target crop species (Figure 1 & 3). This finding is in agreement with findings of [26-27]



264
265
266
267

Figure 3: Effect of *Eucalyptus globulus* plant parts aqueous extract on shoot length of Barely




268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284

Figure 4: Effect of *Eucalyptus globulus* plant parts aqueous extract on root length of Barely

Phytotoxicity

The study shows that the phytotoxicity potential of eucalyptus plant parts aqueous extract increases with the increasing concentrations of aqueous extracts of all extract sources.

285
286
287
288
289




290
291
292
293
294

Figure 5: Inhibitory effect of eucalyptus extract on shoot length of teff. More phytotoxicity effect pronounced under the highest concentration level at 50% of all *Eucalyptus globulus* plant parts aqueous extract compared to the lower level and control treatment. Compared to both crops plant


295 parts, phytotoxicity effect was more pronounced in root length of both crops. *Eucalyptus globulus* fresh leaf
 296 aqueous extract was recognized as the highest impact affecting early root growth of teff and barely at about
 297 69.20% (which is almost the same effect) under the highest level of concentration (50%) of the extract (Figures
 298 6 & 8), however the highest inhibition of teff shoot length at about 54% and shoot length of barely at about
 299 34.5% pronounced under the treatment of mixed extract and fresh leaf extract respectively (Figures 5 & 7) under
 300 the highest level of concentration (50%).

301
 302 There was direct relationship between teff shoot length and inhibitory effects of *Eucalyptus globulus* plant parts
 303 aqueous extract. As the level of *Eucalyptus globulus* plant parts aqueous extract increases, teff shoot inhibition
 304 percentage also increases. Teff shoot length inhibition more pronounced under the treatment of mixed extract
 305 which accounts about 54% compared to control treatment, where there was no teff shoot length inhibition
 306 exhibited. Similar result was obtained by the authors [15, 27]. The intensity of teff root inhibition increases as
 307 the level of *Eucalyptus globulus* plant parts aqueous extract increases, almost all plant parts aqueous extract
 308 inhibited teff root length in similar manner under the same level of concentration.
 309



325
 326 Figure 6: Inhibitory effect of eucalyptus extract on root length of teff.
 327


328 It was fresh leaf aqueous extract that pronounced the highest inhibitory (phytotoxicity) effect on early root
 329 growth of teff at about 69.20% under the highest level of concentration (50%) of the extract, compared to
 330 control treatment which almost do not show teff root inhibition.



331
 332
 333 Figure 7: Inhibitory effect of eucalyptus extract on shoot length of barley.
 334

335 The study of [26] also explained that, aqueous leaf extract of *Eucalyptus globulus* at various levels (doses)
336 inhibited seed germination and early growth of wheat seeds.

337 There was direct relationship between barely shoot length and phytotoxicity (inhibitory) effects of aqueous
338 extract level of *Eucalyptus globulus* plant parts. Barely shoot length inhibition increases as aqueous extract
339 treatments level increase under all treatment level, however barely shoot length inhibition more pronounced
340 under fresh leaf extract at 50% level of concentration. Compared to control which accounts barely root length
341 inhibition at about 34.5% of fresh leaf aqueous extract at 50% level of concentration was exhibited. The study
342 carried out by [27] stated that phytotoxic substances can suppress the growth of many plants by inhibiting seed
343 germination and seedling growth. Barely root length inhibition showed direct proportionality to *Eucalyptus*
344 *globulus* all plant parts aqueous extract level of concentration, Figure 8. More barely seedling shoot length
345 inhibition pronounced under 50% of all aqueous extract of *Eucalyptus globulus* plant parts; however compared
346 to all other extract sources more seedling root length inhibition noticed under fresh leaf aqueous extract at 50%
347 level of extract treatment, that exhibited 69.2% root length inhibition (which is almost the same effect root
348 length inhibition to that of teff root length) compared to control treatment that accounts no barely root length
349 inhibition.
350



369
370 Figure 8: Inhibitory effect of eucalyptus extract on root length of barely.
371

372 Conclusion

373 The result of this study indicates that, higher the concentration of *Eucalyptus globulus* plant extracts, the higher
374 the influences on the germination of barely and teff plant species under laboratory condition. Based on the result
375 of this study, the interaction among various levels of *Eucalyptus globulus* extract treatment depicted that all
376 extracts at lower level (5%) has lower inhibitory effect as compared to higher level (50%) on both crop species.
377 This gradual decrease in germination (%), shoot and root length was due to allelopathic effects of *Eucalyptus*
378 *globulus* aqueous extract from lower level (5%) to higher level (50%) as compared to control. Compared to all
379 aqueous extracts of *Eucalyptus globulus* plant parts of juvenile stem, root, fresh leaf and mixtures of (juvenile
380 stem, root & fresh leaf), fresh leaf aqueous extract pronounced more inhibitory effect on germination percentage
381 (%), shoot length and root length of target crops species. This suggested that there is a variation in phytotoxicity
382 effect or allelochemical concentration in different plant parts. On the overall findings it can also be concluded
383 that allelopathy is a concentration-dependent phenomenon whereby its effect increases as the concentration of
384 the extracts increases. Compared with the control (0%), higher concentrations reduced the germination
385 percentage, shoot length and root length in the given average time.

386 387 REFERENCES

- 388 [1]. Reigosa M.J., Pedrol N & González L. Allelopathy A Physiological Process with Ecological Implications.
389 2006; 1-637
390 [2]. Ferguson J, J, Rathinasabapathi B., and Chase C.A. Allelopathy: How Plants Suppress Other Plants.
391 Horticultural Sciences Department, UF/IFAS Extension, Gainesville, FL 32611. 2016; 1-5

- 392 [3]. Gatti A., Ferreira A., Arduin M., Gualtieri S., & Perez. A. Allelopathic effect of aqueous extract of
 393 *Artistolochiaesperanzae* O.Kuntze on development of *Sesamum indicum* L. seedlings Acta bot. bras. 2010;
 394 24(2): 454-461.
- 395 [4]. Sodaeizadeh H. and Hosseini Z. Allelopathy an Environmentally Friendly Method for Weed Control,
 396 International Conference on Applied Life Sciences. 2012; 387-392.
- 397 [5]. Zhang, Xiaoxi; Liu, Zengwen; Tian, Nan; Luc, Nhu Trung; Zhu, Bochao; Bing, Yuanhao. Allelopathic
 398 effects of decomposed leaf litter from intercropped trees on rape. Turkish Journal of Agriculture and
 399 Forestry. 2015; 39(6): 898-908.
- 400 [6]. Lisanework N. and Michelsen A. Allelopathy in agroforestry systems: the effects of leaf extracts of
 401 *Cupressus lusitanica* and three Eucalyptus spp. on four Ethiopian crops. Agroforestry Systems. 1993; 21:
 402 63-74.
- 403 [7]. Chou, C.H. Roles of allelopathy in plant biodiversity and sustainable agriculture. Critical Reviews in Plant
 404 Science. 1999; 18: 609-636.
- 405 [8]. Getachew Admassu. The impact of eucalyptus plantation expansion on food security in Bambasi Woreda,
 406 Benishangul Gumuz Regional State, Western Ethiopia, Master's Thesis. 2016; 1-45
- 407 [9]. Morsi M. and Abdelmigid, H. M. Allelopathic activity of *Eucalyptus globulus* leaf aqueous extract on
 408 *Hordeum vulgare* growth and cytogenetic behavior. AJCS. 2016; 10(11): 1551-1556
- 409 [10]. Sorecha E.M. and Bayissa B. Allelopathic Effect of *Parthenium hysterophorus* L. on Germination and
 410 Growth of Peanut and Soybean in Ethiopia; School of Natural Resources Management and Environmental
 411 Sciences. *Advances in Crop Science and Technology an open access journal* . 2017; 5 (3): 1-4
- 412 [11]. Gessesse Dessie & Teklu Erkossa. Eucalyptus in East Africa Socio-economic and environmental issues.
 413 Forestry Department Planted Forests and Trees Working Papers. Food and Agriculture Organization of the
 414 United Nations FAO, Rome, Italy. 2011; 1-30
- 415 [12]. Dragoeva AP, Koleva VP, Nanova ZD, Georgiev BP. Allelopathic effect of *Adonis vernalis* L.: Root
 416 growth inhibition and cytogenetic alterations. J Agri Chem Environ. 2015; 4: 48-55.
- 417 [13]. Djanaguiraman M, Vaidyanathan R, Annie sheeba J, Durgadevi D, Bangarusamy U. Physiological responses
 418 of *Eucalyptus globulus* leaf leachate on seedling physiology of rice, sorghum and blackgram. Intl J Agri
 419 Biol. 2005; 7(1):35-38.
- 420 [14]. Fikreyesus S, Kebebew Z, Nebiyu A, Zeleke N, Bogale S. Allelopathic effects of *Eucalyptus camaldulensis*
 421 Dehnh on germination and growth of tomato. Am-Eurasian J Agri Environ Sci. 2011; 11 (5): 600-608.
- 422 [15]. Dejam M, Khaleghi SS, Ataollahi R. Allelopathic effect of *Eucalyptus globulus* Labill. on seed
 423 germination and seedling growth of eggplant (*Solanum melongena* L.). Intl J Farm Alli Sci. 2014; 3 (1):
 424 81-86.
- 425 [16]. Gliessman SR. Allelopathic Effects of Crops. Technology & Engineering, Santa Cruz. 2007; 384.
- 426 [17]. Basazan fantahun and Getachew Belay. Correlation and Path Coefficient Analyses of Some Breeding
 427 Lines of the Man-Made Crop, Triticale (*X. Triticosecale* Wittmak). JAD. 2014; 4(2) 2014.
- 428 [18]. Ameer M, and Rekaby L.S.A. Allelopathic effects of aqueous leaf leachates and root exudations of
 429 *Conocarpus erectus* L. trees against the germination and early growth of some ornamental plants. *Research*
 430 *Journal of Pharmaceutical, Biological and Chemical Sciences*. 2017; 8(4): 741-749.
- 431 [19]. Gurmu W.R. Effects of Aqueous Eucalyptus Extracts on Seed Germination and Seedling Growth of
 432 *Phaseolus vulgaris* L. and *Zea mays* L. *Open Access Library Journal*. 2015; (2): 1-8
- 433 [20]. Meaza Kidane. Extraction and characterization of essential oil from Eucalyptus leaves using steam
 434 distillation. Master's Thesis. 2016; 1-64
- 435 [21]. El-Khawas S.A. & Shehata M.M. The allelopathic potentialities of *Acacia nilotica* and *Eucalyptus rostrata*
 436 on monocot *Zea mays* L and Dicot *Phaseolus vulgaris* L. J. Plant. Biotechnology. 2005; 4(1): 23-34.
- 437 [22]. Shikha R and Jha A. Evaluation of effect of leaf extract of *Parthenium hysterophorus* L. on seed
 438 germination, seedling growth and fresh weight of *Phaseolus mungo*. *American Journal of Research*
 439 *Communication* Volume 4(2). Department of Botany J. P. University, Bihar, India. 2016; 86-103
- 440 [23]. Mujawar I., Kanade M. and Murumkar C. Investigation of allelopathic effect of *Pascaliaglaucaortega* on
 441 seed germination and seedling Growth of wheat. *Indian Journal of Fundamental and Applied Life*. 2016; 6
 442 (3):50-55.
- 443 [24]. Mohammed. M. Hassan, Hussien M. Daffalla, Samia O. Yagoub, Magdoleen G. Osman, Migdam E. Abdel
 444 Gani and Abdel El Gabar E. Babiker. Allelopathic effects of some botanical extracts on germination and
 445 seedling growth of *Sorghum bicolor* L. *Journal of Agricultural Technology*. 2012; 8(4): 1423-1469.
- 446 [25]. Mona. H. Soliman Ahlam, H. Hamad, Hamdah, Al-Gohny and Shroug, S. Al-gohny. Allelopathic Effect of
 447 Moringa oleifera Leaves Extract on Seed Germination and Early Seedling Growth of Faba Bean
 448 (*Vicia faba* L.). *International Journal of Agricultural Technology*. 2017; 13(1): 105-117
- 449 [26]. Solomon W. Senbet. Effects of eucalyptus globulus leaf extract on Germination response and seedling
 450 growth of five Wheat species, Msc Thesis. Adis Ababa, Ethiopia, 2016.

- 451 [27]. Yu DG, Liu GD, Bai CJ. A study on evaluation of *Graminaceous grass* adaptability under Eucalyptus.
452 Chinese Agricultural Science Bulletin. 2008; 24(10): 521-527