

Original Research Article

Assessment of diversity among tropical and subtropical maize inbreds based on morphological traits, grain yield and carotenoid content

ABSTRACT

The available maize germplasm contains sufficient genetic variability for pro-vitamin A content which could be utilized for development of pro-vitamin enriched varieties aimed at reducing vitamin A deficiency among millions of resource-limited Africans. Fifty-one maize inbred lines from Nigeria and CIMMYT (Zimbabwe) comprising of thirty-seven yellow and fourteen white endosperm types were evaluated in replicated trials at the Institute of Agricultural Research and Training, Moor Plantation, Ikenne and the Teaching and Research farm, University of Ilorin. The objective was to assess the extent of diversity among the inbreds using morphological traits, grain yield and carotenoid content. The results showed wide variability ($P = .05$) among the inbred lines for grain yield and other traits as well as total carotenoids. Differences in weather factors in the two locations significantly affected the expression of the traits investigated especially grain yield which recorded the highest deficit in Ilorin due to prolonged drought-stress that occurred at anthesis-flowering period. Although many of the inbred lines exhibited differential performances for grain yield across the two locations, three inbred lines- BD74-49, BD74-68 and BD74-81 showed consistency in their performance for this trait across the two locations. Significant variation in kernel carotenoid content was also obtained among the 40 maize genotypes investigated. Total carotenoid ranged from as low as 0.03 μ g/g in the white kernel inbred TZEI 65 to 56.52 μ g/g in a light orange coloured inbred line BD74-89. The UPGMA dendrogram constructed from the morphological traits and grain yield clustered the 51 inbred lines into two groups based essentially on their geographical origin, while the dendrogram generated based on total carotenoid content clustered the 40 inbred lines into two major groups with clusters of at least three inbred lines in a sub-cluster. Clustering patterns revealed that lines with high carotenoid content were found in different sub-clusters indicating the feasibility of developing high carotenoid maize varieties from the currently available gene pool.

Key words: *Zea mays L., Pro-vitamin A, morphological traits, carotenoid content, inbred lines.*

1. INTRODUCTION

Carotenoids which are natural plant pigments in the endosperm and fruits of crop plants are important source of vitamin A and antioxidants that are essential for human consumption. Dietary carotenoids are essential precursors of vitamin A and retinoid compounds needed in animal morphogenesis (Bendich and Olson, 1989), which plays an important role in the proper functioning of the eye, growth of bone, reproduction, cell division, and cell differentiation. It also regulates the immune system, which helps prevent or fight off infections by making white blood cells that destroy harmful bacteria and viruses. Maize (*Zea mays* L.) kernel contains tremendous variation in content and composition of several coloured pigments collectively known as carotenoids and is known for its genetic diversity of carotenoid content and profiles (Menkir *et al.*, 2008). Consequently, it is one of the potential crops for bio-fortification aimed at increasing the bio-available micronutrient content of food crops through genetic selection

Comment [ss1]: Not consistent with the information provided in the materials and method section, CIMMYT Kenya.

Comment [ss2]: Flowering period – this contains both anthesis and silking periods.

Comment [ss3]: State its full meaning

through plant breeding. Pfeiffer and McClafferty (2007) as well as Wurtzel *et al.*, (2012) hypothesized that exploitation of the natural genetic diversity of maize for carotenoids through bio-fortification will increase pro-vitamin A concentration in maize endosperm, which will be beneficial for public health. Secondly, the process also holds promise for sustainable and cost-effective food-based solutions to combat micronutrient deficiencies in Africa where maize is a major security and staple crop for more than 300 million people (Smale *et al.*, 2013). However, pro-vitamin A usually constitutes only 10 to 20% of the total carotenoids in maize kernel, and the study conducted by Pixley *et al.*, (2013) revealed that the commonly cultivated and consumed yellow maize cultivars have less than 2µg/g dry weight pro-vitamin A.

Diversity study is important in assessing the extent, pattern and association among breeding populations. Genetic diversity assessment has also been used in the identification of parental lines for evolving superior genotypes that could be utilized in crosses and in the establishment of heterotic groups that could yield superior hybrids in maize. In recent years, molecular breeding techniques including SSR-based markers (Smith *et al.*, 1997; Warburton *et al.*, 2002) have been utilized in the assessment of genetic diversity among breeding populations for different maize attributes including carotenoid contents. The importance of developing pro-vitamin A enriched maize variety underscores the necessity to have information on the extent of diversity among inbred parents with respect to pro-vitamin contents. Such varieties could be used in the development of yellow endosperm maize varieties (Open Pollinated Varieties and hybrids) which possess genes for adaptation and superior agronomic performance and thus represent a good genetic base to breed tropical maize for high levels of pro-vitamin A carotenoids (Russell *et al.*, 1997). In two separate studies, Adeyemo *et al.*, (2012) as well as Adeyemo and Omidiji (2014) assessed the nature and extent of the genetic diversity among tropically adapted yellow endosperm maize inbred lines ~~in~~ at the International Institute ~~for~~ of Tropical Agriculture (IITA) maize breeding programme. In the first study, Adeyemo *et al.*, (2012), noted the diversity of carotenoid contents among 38 tropically adapted yellow endosperm maize inbred lines while the second report (Adeyemo and Omidiji, 2014) contained information on the nature of variability for different carotenoid contents among the newly developed 122 inbred lines and grouped into three major groups based on carotenoid contents. In the study reported herein, 51 maize inbred lines collected from two major geographical regions (Nigeria and Kenya), were evaluated in two locations using 12 morphological traits (including grain yield) and carotenoid content. The objectives were to (i) assess the extent of genetic diversity among white and yellow endosperm maize inbred lines, (ii) identify lines with high carotenoids content, and (iii) delineate the genotypes for future breeding purposes.

2. MATERIALS AND METHODS

2.1 Source of Genetic Materials

Fifty-one maize inbred lines were received from three different sources viz., CIMMYT (Nairobi, Kenya), International Institute of Tropical Agriculture (IITA), Ibadan and Institute of Agricultural Research and Training (IAR&T), Ibadan were used for this study. The details of the inbred lines including their grain colour are presented in Table 1.

2.2 Description of Experimental Sites

Ikenne (61 m above sea level, Latitude 6° 52'N and Longitude 3° 42' E) is in the rain forest ecology with annual rainfall and temperature ranges of 1000-1200 mm ~~and with temperature range of 22°C – 32.2°C, respectively.~~ Ilorin (370 m above sea level and Latitude 8° 30' N and,

Longitude 4° 32'E) is in the southern guinea savannah ecology with annual rainfall ~~about of~~ 1000 mm and temperature range of 24°C – 35°C ~~respectively~~. However, while rainfall distribution at both locations is bimodal, mid-season drought is likely to occur at any stage of crop growth. This is a regular feature of weather pattern at Ilorin which makes the zone a drought-prone ecology.

2.3 Field Evaluation

The inbred lines were evaluated at the IAR&T sub-station, Ikenne during the growing season of 2012 and at the Teaching and Research Farm, University of Ilorin between July-September of 2013. Seeds were sown at each location using a randomized complete block design with three replicates. Each plot consisted of single row with spacing of 75cm between the rows and plant to plant spacing of 25cm, giving a plant population of 53,333 plants/ha. Pre-emergence herbicide (Atrazine 223 g/L + Metolachlor 277 g/L) was applied at planting at both sites followed by supplementary hand weeding as required. The recommended fertilizer dose and other cultural practices in each zone were adopted.

2.4 Morphological study

Data were recorded from five random plants in a plot on twelve traits namely; days to mid-anthesis and silking, anthesis-silking interval (ASI), plant and ear heights (cm), leaf length and width (cm), ear length and diameter (cm), number of rows per ear, number of kernels per row and grain yield (kg/plot). Days to mid-anthesis and silking were estimated as the number of days from planting to when 50% of the plants in a plot have shed pollen and had silk emergence respectively. ASI was estimated as the difference between ~~to the~~ mid-anthesis and silking. Plant height was measured from the soil surface to the tip of the central axis while ear height was recorded in centimeters from the distance ~~from of~~ the ground to the base of the uppermost developed ear. Grain yield was obtained by weighing ears in each plot in kg/plot and later converted to tons/ha after adjusting the moisture level to 12% and assuming a shelling percentage of 80%.

2.5 Data Analyses

Data collected were subjected to analyses of variance (ANOVA) ~~first~~ on individual location ~~basis before a prior to combined ANOVA across locations analysis~~. The quantitative traits were also subjected to cluster analysis. Sequential Agglomerative Hierarchical Non-overlapping (SAHN) clustering was performed on similarity matrices utilizing the Unweighted Pair Group Method with Arithmetic Averages (UPGMA) method. Data analysis was done using NTSYSpc software, version 2.02i (Rohlf, 1998).

2.6 Carotenoid analysis

Forty of the inbred lines were further analyzed for their carotenoid contents. Spectrophotometry was used to determine optical density (OD) values of each sample at 440nm and subsequently, using the Lambert Beer equation, carotenoid content of each sample was determined as follows. The data on kernel carotenoids were analyzed in a completely randomized design (CRD) to test the statistical validity of variance among different genotypes for maize kernel carotenoids content. The total carotenoid content of maize kernel was determined spectrophotometrically using Lamber-Beer equation;

$$E = \epsilon \times c \times d$$

Where,

Comment [ss4]: Is it husked or dehusked ear weights? Please specify.

Comment [ss5]: Why 40 of the inbred lines (out of the 51 inbreds) were used for the carotenoid analysis? And how can the selected 40 inbreds be identified in case one wants to use them for future work?

What was your sample size per inbred line (i.e. number of ears or kernels for each inbred line) used for the carotenoid analysis?

E = extinction (photometer reading)

ϵ = molar extinction coefficient

c = concentration

d = distance = 1, to determine concentration

$c = E/\epsilon$

Lutein, Zeaxanthin and beta-carotene are the major carotenoids in maize. Therefore, an average value for ϵ and for the molecular mass was used.

ϵ lutein = 122,688 l/mol cm; M = 568 g/mol

ϵ zeaxanthin = 133,480 l/mol cm; M = 568 g/mol

ϵ β -carotene = 134,000 l/mol cm; M = 537 g/mol

ϵ average = 130,056 l/mol cm;

M average = 557.7 g/mol.

Hence, $c = OD/130056 \times 557.7 \times 1000/0.5$

Where

C = concentration of total carotenoids content ($\mu\text{g/g}$) in a given sample on dry weight basis;

OD = optical density taken at 450nm wavelength using a spectrophotometer;

W = weight of sample (0.5g).

Each sample was analyzed in three replicates and the data was subjected to statistical analyses using completely randomized design (CRD).

Table 1: Maize inbred lines used for genetic diversity study

S/N	Genotype	Pedigree	Source	Grain Colour
1	Eruwa-3-1	Unknown	IAR&T	White
2	Ijebu-Ode W-2	Unknown	IAR&T	White
3	Ikole 1-W	Unknown	IAR&T	White
4	Kishi 4-1	Unknown	IAR&T	White
5	Isara 1-W	Unknown	IAR&T	White
6	Bdg-1	Unknown	IAR&T	White
7	Bodija Shgari	Unknown	IAR&T	White
8	TFE 20ABI-1	Unknown	IAR&T	Yellow
9	TZM 228Y	Unknown	IAR&T	Yellow
10	TZM-1327-A	Unknown	IAR&T	Yellow
11	Isara-1-Y	Unknown	IAR&T	Yellow
12	Ijebu-Igbo-Y	Unknown	IAR&T	Yellow
13	Eruwa Local5-1	Unknown	IAR&T	Yellow
14	TZEI 135	TZE-Y Pop STR Co S6 Inbred 17-2-3	IITA IB	Yellow
15	TZEI 157	TZE-Y Pop STR Co S6 Inbred 102-1-2	IITA IB	Yellow
16	TZEI 17	TZE Comp5-Y C6 S6 Inbred 35	IITA IB	Yellow
17	TZEI 10	TZE-Y Pop STR Co S6 Inbred 152	IITA IB	Yellow
18	TZEI 16	TZE Comp5-Y C6 S6 Inbred 31	IITA IB	Yellow
19	TZEEI 74	TZEF-Y SR BC1 x 9450 STR S6 Inb 5A	IITA IB	Yellow
20	TZEEI 72	TZEF-Y SR BC1 x 9450 STR S6 Inb 2C	IITA IB	Yellow
21	TZEEI 68	TZEE-Y SR BC1 x 9450 STR S6 Inb 11	IITA IB	Yellow
22	TZEEI 64	TZEE-Y SR BC1 x 9450 STR S6 Inb 8A	IITA IB	Yellow
23	TZEEI 82	TZEF-Y POP STR COS6 Inb44	IITA IB	Yellow
24	TZEEI 75	TZEF-Y SR BC1 x 9450 STR S6 Inb 7B	IITA IB	Yellow
25	TZEEI 73	TZEF-Y SR BC1 x 9450 STR S6 Inb 3A	IITA IB	Yellow

26	TZEEI 59	TZEE-Y SR BC1 x 9450 STR S6 Inb 3A	IITA IB	Yellow
27	TZEEI 76	TZEE-Y SR BC1 x 9450 STR S6 Inb 8B	IITA IB	Yellow
28	TZEI 65	TZE-W Pop STR Co S6 Inbred 141-1-2	IITA IB	White
29	TZEI 165	TZE Comp5-Y C6 S6 Inbred 8A	IITA IB	Yellow
30	TZEI 146	TZE-Y Pop STR Co S7 Inbred 49-3-3	IITA IB	Yellow
31	TZEI 129	TZE-Y Pop STR Co S6 Inbred 16-1-3	IITA IB	Yellow
32	BD74-43	CML12-B	CYMMIT	Yellow
33	BD74-44	CML130-B	CYMMIT	Yellow
34	BD74-48	CML161-B	CYMMIT	Yellow
35	BD74-49	CML165-B	CYMMIT	Yellow
36	BD74-68	CML451-B	CYMMIT	Yellow
27	BD74-81	CLYN242-B	CYMMIT	Yellow
38	BD74-88	CLYN243-B	CYMMIT	Yellow
39	BD74-89	CLYN244-B	CYMMIT	Yellow
40	BD74-90	CLYN226-B	CYMMIT	Yellow
41	BD74-91	CLYN246-B	CYMMIT	Yellow
42	BD74-92	DTPYC9-F114-2-4-1-1-B	CYMMIT	Yellow
43	BD74-107	CLRCY031-B	CYMMIT	Yellow
44	BD74-109	CLQRCYQ14-B	CYMMIT	Yellow
45	BD74-112	CLYN208-B	CYMMIT	Yellow
46	BD74-155	CLYN247-B	CYMMIT	White
47	BD74-156	(DTPYC9-F46-1-2-1-2-B)-B	CYMMIT	White
48	BD74-157	(DTPYC9-F38-4-3-1-1-B)-B	CYMMIT	White
49	BD74-158	(DTPYC9-F15-3-4-1-1-B)-B	CYMMIT	White
50	BD74-159	(DTPYC9-F143-5-4-1-2-B)-B	CYMMIT	White
51	BD74-160	(DTPYC9-F13-2-3-1-2-B)-B	CYMMIT	White

3. RESULTS AND DISCUSSION

3.1 Genotypic performance for grain yield and related characters

The maize inbreds differed significantly ($P = .01$) in the expression of the characters evaluated from one location to the other except kernel rows/ear (Table 3), indicating differences in the environmental factors at each location. Means for all the characters were higher at Ikenne compared to Ilorin with mean plant and ear heights >100cm and accompanied by a difference of 1.97 t/ha in grain yield. Majority of the inbred lines also yielded less 1.0 t/ha at Ilorin which may be due to prolonged drought stress towards the flowering period which supports earlier reports of differential performances of maize genotypes occasioned by drought and air temperature stresses (Megyes *et al.*, 2005; Halof and Savari, 2007; Kovacevic *et al.*, 2009; Randjelovic *et al.*, 2011; Hao *et al.*, 2016; Gheysari *et al.*, 2017). The genotypes also differed significantly ($P = .01$) for all the characters with the magnitude of the mean squares due to genotype being larger than the genotype by location interaction mean squares for all the characters except for anthesis-silking-interval and number of rows/kernels. The genotype by location interaction effects also differed significantly ($P = .01$) for all the characters which suggests differences in the genotypes response to the prevailing environmental conditions in both locations. Consequently, the differences in weather factors at the two locations contributed largely to the observed inconsistency in the inbred lines performance especially for grain yield.

The ranges in the means among the inbred lines were very large for most of the traits studied except for the flowering traits, leaf width and ear diameter (Table 3). For example, differences between the highest and poorest yielding inbred was > 3 t/ha. However, plant and ear heights were the most variable among the inbred lines with a difference of > 200cm between the tallest and shortest genotype and > 100cm between the highest and lower ear placements.

Comment [ss6]: This should be Table 2, not Table 3.

Mean performances of the maize inbreds for characters across the two locations showed less variation for flowering traits (Table 43). The inbred lines were highly variable for the characters studied with different genotypes exhibiting superiority for each of the characters. For example, inbred line BD74-49 was the highest yielding genotype with a superiority of 1.32 t/ha over TZEEI 76 which was the poorest yielding inbred. Another inbred line (BD74-157) had the largest ear diameter while inbred BD74-49 exhibited superiority of 38.98% over TZEI 68 for ear length. The highest number of kernel rows per ear was recorded in genotype BD74-157 while inbred line BD74-49 had maximum number of kernels per row. Inbred BD74-158 was the tallest and also had the highest ear placement while inbred TZEI 17 was the shortest with the corresponding lowest ear placement. Values obtained for leaf parameters showed that inbred BD74-92 had the longest leaf while inbred BD74-48 had the broadest leaf. The inbred lines also changed rank for grain yield from one location to another with Inbreds BD74-49, and BD74-89, Kishi 4-1, BD74-68 and BD74-159 in that order being the top five yielding genotypes at Ikenne while inbreds BD7-49, BD7-81, Eruwa 3-1, BD7-68 and BD74-107 were the five top yielding hybrids at Ilorin. Therefore, the alteration in ranking of the inbred lines in their performances for grain yield and other traits observed, must have been due to response to the factors of environment in each location which justifies the use of the multi-trait selection method to identify the best genotypes for further varietal (OPVs and Hybrid) development programme.

3.2 Cluster analysis of genotypes based on morphological traits and grain yield

The dendrogram from the data generated based on the 12 attributes and revealed by UPGMA, clustered the inbred populations into two major groups with similarity coefficient of 1.40 (Figure 1). The first major group comprised 21 inbred lines and all were of Nigerian origin (i.e. IITA and IAR&T) while the second major group comprising 30 inbred lines were all from CYMMIT except Eruwa 3-1 which is of Nigerian origin. The genotypes in the first major group were further sub-divided into four sub-clusters of 10, 4, 2 and 5 genotypes respectively. The second major group was also further sub-divided into eight sub-clusters. Sub-clusters I and II in the second group comprised two and four genotypes respectively with a similarity coefficient of 0.68 while Sub-clusters III, IV, V and VI respectively comprised five, two, two, and four genotypes. Sub-clusters VII and VIII had seven and two genotypes respectively.

3.3 Genotypic performance for Estimation of total carotenoid in genotypes

The quantitative estimation of the total carotenoid of the 40 inbred lines selected for the investigation revealed significant differences ($p < 0.01$) which indicated that they were genetically different for this character (Table 54), which is the basic need to take up any breeding programme for improvement of any trait. For example, the mean value for kernel carotenoids content varied from minimum of $0.03\mu\text{g/g}$ dry weight in inbred line TZEI 65 to as high as $56.52\mu\text{g/g}$ dry weight in inbred BD74-89 with population mean of $22.37\mu\text{g/g}$. Sivaranjani *et al.*, (2013) in their own study also reported wide range variability ($6.5 - 67.3\mu\text{g/g}$) for carotenoid contents in maize. Several studies have earlier reported carotenoid content ranging from $0.94\mu\text{g/g}$ to $66.0\mu\text{g/g}$ in different maize populations which also was dependent on the population being investigated, region where the study was conducted and stage of improvement of the

population in maize (Daood *et al.*, 2003; Egesel *et al.*, 2003; Hulshof *et al.*, 2007; Harjes *et al.*, 2008; Mishra and Singh, 2010; Cardoso *et al.*, 2009; Kumar and Singh, 2012).

The accumulation of carotenoids the biosynthesis of which occur during seed development (Li *et al.*, 2008), impact a yellow-orange colour to the endosperm and are an easily scored phenotype. Consequently, the higher mean values for total carotenoid associated with coloured endosperm in the inbred lines is an indication of considerable high concentration of total carotenoids (Kumar and Singh, 2012). When coloured endosperm was taken into consideration, the range of kernel carotenoids content varied from a minimum of 7.7µg/g to a maximum of 56.52µg/g dry weight in maize inbred BD74-89. Buckner *et al.*, (1996) hypothesized that the dominant *Yellow1* (*Y1*) allele in maize controls the first rate-limiting step in the carotenoid biosynthesis by coding for the enzyme phytoene synthase, while the homozygous recessive allelic state (*y1y1*) results in almost no synthesis of carotenoids in the endosperm. In the present study, white kernel inbred lines (for example, TZEI 65: 0.03 µg/g) having *y1* allele in the endosperm produced almost no carotenoid in the endosperm. Presence of similar negligible amount of total carotenoids (1.14µg/g) in the white maize endosperm (*y1y1y1*) was also reported by Buckner *et al.*, (1996). The significant range for the total carotenoids (7.71 - 56.52µg/g) in the yellow/orange inbred lines could be due to the control of carotenoid biosynthesis by other genes, besides the *Y1*.

The inbred lines with kernels of different shades of yellow comprised genotypes with relatively high and low total kernel carotenoids. Thus, the impact of kernel colour on total carotenoid content could not be ascertained in this study. Harjes *et al.*, (2008) and Mishra and Singh (2009) also observed poor association between these two parameters in their own studies. However, inbred lines- BD74-89, TZEEI 82, Bodija-Shagari-Y, PVA SYN 13 (9), TZEEI 75, TZEI 146, TZEI 165 and Eruwa Local 5-1-Y had high kernel carotenoid content and can be advanced for breeding programmes geared towards improved carotenoid content in future maize varieties (OPVs and hybrids) of maize.

3.4 Cluster Analysis of the Genotypes based on Carotenoid Content

The dendrogram of the data generated from total carotenoid content separated the 40 maize inbred lines into two major clusters (Figure 2) with the larger cluster (Cluster I) having 31 genotypes and Cluster II with nine (9) genotypes. However, the clustering of the inbred lines for carotenoid content was not based on geographical origin. Cluster I was further subdivided into five sub-clusters of 11, 6, 2, 7 and 5 genotypes respectively while cluster II was also sub-divided into two sub-clusters of 3 inbred lines each. The least genetic similarity was observed in genotypes TZEI 65, TZEEI 82 and BD74-89. Due to wider level of diversity based on carotenoid content, nine (9) genotypes were clearly separated and grouped in a distinct cluster.

4. Conclusion

Hybrid maize breeding programmes depended on selection of diverse inbred lines in order to utilize hybrid vigour as a means of improving crop productivity. Consequently, understanding the genetic diversity among important breeding materials is generally considered a critical first step to achieve success in breeding programmes. The selection process of good performing and stable genotypes is usually complicated by the intangible forces of environment resulting in the phenomenon of genotype by environment (G x E) interaction. Consequently, Prolonged drought stress which coincided with the grain filling period at Ilorin significantly reduced grain yield in that location compared to Ikenne. The difference between the two locations was 1.97 t/ha

representing >80 percent yield advantage. However, three inbred lines – BD74-49, BD74-68 and BD74-81 still showed consistency in their performance for this grain yield across the two locations, indicating their stability of performance for grain yield.

The carotenoid analysis using spectrophotometer results show that nine (9) inbred lines had the highest carotenoid content. Inbred line – BD74-89 was identified as one of the potential sources of carotenoids and hopefully pro-vitamin A as it exhibited the highest carotenoid content among the 40 inbred lines. The 31 inbred lines included in group 1 of the dendrogram had low concentration of carotenoid except for TZEI 165 and Eruwa 5-1-Y while 9 inbred lines included in group II had increased concentration of carotenoid.

Comment [ss7]: Also outline/mark out here in the inbred lines that you identified for future breeding purposes.

Table 2: Weather Data for Teaching and Research (T & R) Farm, Unilorin (2013).

Month	Rainfall (mm)	Temperature		Evaporation (ml)	Wet Bulb reading (°C)	Dry bulb reading (°C)
		Minimum (°C)	Maximum (°C)			
April	92.10	22.60	29.96	9.79	25.20	31.50
May	56.20	24.90	30.10	6.30	27.70	32.70
June	87.00	24.90	30.00	6.04	24.50	29.50
July	108.10	25.10	26.60	5.80	25.00	32.00
August	5.50	25.70	25.90	4.90	24.00	27.50
September	169.30	24.80	29.60	1.57	28.00	30.00
October	136.80	24.00	30.00	2.70	25.00	30.50
November	0.0	25.03	29.70	5.40	25.50	31.00
December	0.0	23.50	30.06	6.33	26.00	31.30

Comment [ss8]: Remove this Table or otherwise provide such information for the Ikenne site used in the study.

The information here for rainfall is not tallying with the one provided in the sites description in the materials and methods sections.

Table 32: Performance of the maize inbreds across locations for morphological at Ikenne and Ilorin and combined analysis of variance for morphological traits, grain yield and carotenoid attributes

Location	Days to		Anthesis Silking Interval (days)	Plant height (cm)	Ear height (cm)	Leaf length (cm)	Leaf width (cm)	Ear diameter (cm)	Ear length (cm)	No. of rows per ear (number)	No. of Kernel per row (number)	Grain yield (t/ha)
	50% Anthesis (days)	50% Silking (days)										
Ikenne	56.51	59.33	2.96	196.87	105.68	60.97	6.14	4.24	17.32	12.10	18.81	2.20
Ilorin	57.10	60.45	3.41	94.70	36.68	50.22	5.86	1.44	11.05	11.17	18.81	0.23
Mean	56.81	59.99	3.20	145.78	71.08	55.58	6.0	2.84	14.18	11.63	18.81	1.21
Range	7.00	8.00	4.60	123.00	93.00	64.00	6.50	7.48	21.50	10.00	36.00	3.57
SE±	0.68	0.69	0.30	10.74	8.86	5.21	0.61	0.75	2.63	1.85	4.95	0.25
F - Test												
Replication	1.95	3.26	0.42	132.46	26.80	45.31	0.76	0.95	38.45	26.00	124.51	0.16
Location (L)	28.26**	64.97**	10.62**	789509.13**	366234.08**	8767.04**	6.29**	600.34**	3006.74**	66.82**	0.00	295.75**
Genotype (G)	5.99**	6.40**	0.39**	7479.06**	4703.57**	788.46**	9.37**	1.61**	19.71**	5.23**	75.64**	0.54**
G x L	2.88**	2.87**	0.56**	795.51**	1139.83**	99.94**	0.84**	0.51**	9.98**	5.84**	29.18**	0.29**
Pooled Error	0.46	0.48	0.09	115.28	78.57	27.16	0.32	0.56	6.94	3.42	24.50	0.06
%CV	1.20	1.15	9.60	7.36	12.46	9.37	10.23	26.44	18.58	15.90	26.31	21.21

** , Significant F-Test at 0.01 level of probability

Table 43: Agronomic performance of maize inbreds at Ikenne and Ilorin combined, yield and Genotypic parameters of both locations of maize inbred lines

Genotype	Days to anthesis (days)	Days to silking (days)	ASI (days)	Plant height (cm)	Ear height (cm)	Leaf length (cm)	Leaf width (cm)	Ear diameter (cm)	Ear length (cm)	No. of row per ear (no)	No. kernel per row (no)	Grain yield (t/ha)
BDG-1	57	60	3	113.50	46.27	49.98	4.97	2.65	13.03	11.67	17.67	1.00
Bodija Shagari	58	62	3	125.88	54.88	50.70	5.62	2.71	13.83	10.50	17.83	1.25
BD74-109	58	61	4	179.00	104.00	64.03	6.35	2.98	15.00	11.83	16.67	1.35
BD74-107	57	60	3	192.17	109.50	71.02	7.47	3.19	17.02	12.00	17.00	1.53
BD74-112	57	61	4	182.17	110.33	56.62	6.23	3.19	15.95	12.67	18.00	1.37
BD74-90	57	61	4	162.50	91.17	56.62	4.95	2.79	13.18	12.00	18.17	1.42
BD74-81	56	59	3	185.33	102.83	70.67	8.10	3.71	17.62	12.50	27.17	1.55
BD74-88	58	61	4	179.50	98.17	62.55	6.78	3.39	15.13	10.67	14.33	1.26
BD74-89	57	60	3	183.50	97.83	74.02	7.68	3.05	14.70	11.83	19.83	1.61
BD74-91	58	61	3	192.17	103.50	71.28	7.83	3.29	17.75	12.67	24.33	1.45
BD74-155	59	62	3	195.67	112.00	67.65	7.02	3.24	17.17	12.67	18.00	1.39
BD74-43	57	60	3	189.17	101.50	69.80	7.83	2.95	12.88	11.17	18.33	1.55
BD74-44	56	59	3	167.50	93.67	62.98	6.90	3.23	14.23	12.83	21.83	1.45
BD74-48	58	61	3	198.33	106.00	72.05	8.38	3.41	16.88	12.00	22.33	1.54
BD74-49	58	61	4	200.83	107.50	76.88	8.10	3.97	19.55	13.17	29.33	2.04
BD74-68	56	59	3	186.67	101.33	74.95	8.05	3.48	16.65	12.67	23.83	1.70
BD74-157	59	63	4	197.67	111.67	71.75	6.82	3.78	14.45	14.67	24.00	1.39
BD74-158	58	61	4	206.67	113.75	64.67	7.40	3.09	16.98	11.83	23.67	1.37
BD74-159	59	62	3	193.50	105.91	64.00	7.33	3.43	14.77	11.67	20.67	1.51
BD74-160	59	62	3	188.67	106.50	65.97	7.02	3.20	14.52	12.33	22.50	1.43
BD74-92	57	60	3	180.17	107.00	77.93	8.05	3.14	15.07	11.67	16.17	1.47
BD74-156	59	62	3	177.50	102.17	60.83	6.78	3.66	15.75	11.17	19.83	1.49
Eruwa-3-1	57	60	3	145.67	60.01	60.83	6.53	3.62	16.65	14.50	25.33	1.37
Eruwa Local5-1	58	61	3	104.43	43.97	51.27	5.27	2.57	13.45	11.00	18.00	1.25
Ijebu-Igbo-Y	57	60	3	142.50	68.73	55.80	5.27	2.48	13.55	11.67	15.33	1.22
Ijebu-Ode-W-2	57	60	3	141.08	71.73	49.50	6.20	3.07	12.10	11.33	16.00	1.32
Ikole-1-W	56	60	3	131.33	63.75	53.48	5.90	3.46	12.55	11.00	15.83	1.47

Isara-1-Y	58	61	3	114.85	57.08	48.93	5.52	3.07	13.15	11.33	25.33	1.40
Isara-I-W	56	60	4	110.11	48.08	45.92	6.75	3.46	13.40	11.17	15.83	1.24
Kishi-4-1	57	60	3	129.48	56.67	51.87	5.60	2.54	14.22	11.33	15.33	1.56
TFE 20ABI-1	57	60	3	139.81	56.43	58.53	5.63	2.67	13.20	11.50	18.67	0.94
TZEEI 129	55	58	3	115.17	43.25	46.12	4.48	2.56	14.40	11.00	19.00	0.79
TZEEI 146	55	59	4	113.25	47.27	51.67	4.67	2.44	12.88	10.83	16.83	1.03
TZEEI 165	56	58	3	110.08	43.83	44.43	4.43	2.56	11.93	11.00	17.00	0.93
TZEEI 59	56	59	4	116.67	45.97	41.27	4.80	1.96	13.28	11.83	18.00	0.90
TZEEI 65	56	59	3	109.17	43.17	43.47	5.72	2.18	12.25	11.33	18.00	0.87
TZEEI 76	56	59	3	116.33	46.50	43.45	5.30	2.19	13.67	11.17	16.67	0.72
TZEI 10	56	59	3	123.67	46.50	44.67	5.10	2.27	14.07	11.17	18.33	0.84
TZEI 135	56	59	3	114.83	44.50	40.33	4.73	2.77	11.77	12.50	20.33	0.98
TZEI 157	56	60	4	109.50	43.17	50.50	4.30	2.34	13.38	11.33	16.83	0.82
TZEI 16	57	60	4	119.17	44.00	46.83	5.53	2.14	12.70	11.33	16.67	0.75
TZEI 17	56	59	3	100.83	36.83	44.00	4.17	2.54	12.67	10.67	19.17	0.89
TZEI 64	56	59	3	110.00	42.83	46.83	5.03	1.96	13.52	11.00	13.33	0.99
TZEI 68	56	59	3	106.00	39.62	36.83	4.87	2.17	11.93	10.83	15.83	0.86
TZEI 172	55	58	3	108.67	44.00	42.58	5.17	2.03	13.42	10.00	17.00	0.93
TZEI 173	57	60	3	118.00	50.67	43.90	5.33	2.65	12.97	12.00	17.67	0.81
TZEI 174	56	59	4	119.50	49.33	38.67	4.05	2.29	11.83	9.83	12.50	1.00
TZEI 175	56	59	3	109.17	40.17	48.60	4.08	2.74	14.40	10.67	15.67	0.93
TZEI 182	57	60	4	115.00	47.67	46.00	4.77	2.34	12.45	11.17	14.83	0.94
TZM-1327-A	57	61	4	126.98	53.17	54.00	5.68	2.56	12.30	11.33	19.17	1.02
TZM 228-Y	56	60	4	135.76	58.91	47.67	5.63	2.39	13.17	11.67	17.33	1.19
SE±	0.15	0.16	0.06	4.94	3.92	1.61	0.17	0.07	0.25	0.13	0.50	0.04

Comment [ss9]: Make provision for means of traits.




Figure 1: Dendrogram of 51 maize inbred lines revealed by UPGMA cluster analysis based on 12 different morphological and yield traits.

Table 54: Kernel Colour and Total Carotenoids in 40 maize inbred lines

S/N	Genotype	Kernel Colour	Total Carotenoids (µg/g) dry weight
1	Bodija Shagari	Orange	33.93
2	BD74-107	Yellow	21.97
3	BD74-90	Yellow	24.46
4	BD74-81	Orange	10.07
5	BD74-88	Orange	26.42
6	BD74-89	Light orange	56.52
7	BD74-91	Orange	25.81
8	BD74-43	Orange	14.43
9	BD74-44	Light orange	16.97
10	BD74-48	Orange	14.56
11	BD74-49	Orange	19.31
12	BD74-68	Orange	7.71
13	BD74-92	Yellow	13.91
14	Eruwa Local 5-1	Orange	31.38
15	Ijebu-Igbo-Y	Yellow	19.04
16	Ijebu-Ode-Y	Yellow	16.32
17	Isara-1-Y	Light orange	25.77
18	TZEI 129	Yellow	16.37
19	TZEI 146	Yellow	33.46
20	TZEI 165	Yellow	31.23
21	TZEEI 59	Yellow	19.58
22	TZEI 65	White	0.03
23	TZEEI 76	Light Orange	12.84
24	TZEI 135	Light orange	15.20
25	TZEI 157	Light orange	27.70
26	TZEI 16	Light orange	19.52
27	TZEI 17	Light orange	15.95
28	TZEEI 68	Yellow	26.42
29	TZEEI 72	Yellow	22.90
30	TZEEI 73	Orange	29.39
31	TZEEI 74	Orange	9.78
32	TZEEI 75	Yellow	30.16
33	TZEEI 82	Yellow	43.97
34	TZEI 64	Light orange	29.57
35	TZEI 181	Yellow	20.58
36	TZEI 128	Light orange	16.98
37	TZEI 67	Light orange	25.86
38	TZM 228-Y	Yellow	16.53
39	PVA SYN 13 (9)	Light orange	34.64
40	PVA SYN 16 (9)	Light orange	18.07
Mean			22.37
SE ±			1.64
%CV			7.35

REFERENCES

- Adeyemo O, Omidiji O. SSR-based and carotenoid diversity assessment of tropical yellow endosperm maize inbred lines. *Plant Genetic Resources: Characterization and Utilization*. 2014;12(1); 67–73
- Adeyemo O, Menkir A, Melaku G, Omidiji O. Genetic diversity assessment and relationship among tropical yellow endosperm maize inbred lines using SSR markers. *Maydica*. 2012;56: 17.
- Bendich A, Olson JA. Biological actions of carotenoids. *FASEB Journal*. 1989; 3:1927-1932.
- Buckner B, Miguel, PS Buckner, DJ Bennettzen JL. The *y1* gene of maize codes for phytoene synthase. *Genetics*. 1996;143: 479– 88.
- Cardoso WS, Peas MCD, Galvao JCC, Rios SD, Guimaraes PED, Schaffert RE, Borem A. Variability of maize genotypes for grain carotenoid composition. *Brazilian Journal of Agricultural Research*. 2009;44:164–72.
- Daood HGM, Nagy-Gasztonyi G, Vorosvary L, Holly L, Horvath M, Nyerges. Analysis of carotenoid in some Hungarian maize landraces. *Maize Breeding and Utilization*. 2003;1–2.
- Egesel CO, Wong JC, Lambert RJ, Rocheford TR. Combining ability of maize inbreds for carotenoid and tocopherols. *Crop Science*. 2003;43: 818–23.
- Gheysari M, Sadeghi SH, Loescher HW, Amiri S, Zareian MJ, Majidi MM, Asgarinia P, Payero JO. Comparison of deficit irrigation management strategies on root, plant growth and biomass productivity of silage maize. *Agricultural Water Management*. 2017;182: 126–138.
- Halof EN, Sarvari M. Relationship between yield quality and quantity of maize hybrids and fertilizer. *Cereals Research Communications*. 2007;35:369-372.
- Hao B, Xue Q, Marek TH, Jessup KE, Hou X, Xu W, Bynum ED, Bean BW. Radiation-use efficiency, biomass production, and grain yield in two maize hybrids differing in drought tolerance. *Journal of Agronomy and Crop Science*. 2016;202: 269–280.
- Harjes CE, Rocheford TR, Bai L, Brutnell TP, Kandianis CB, Sowinski SG, Stapleton AE, Vallabhaneni R, Williams M, Wurtzel ET, Yan J, Buckler ES. Natural genetic variation in lycopene epsilon cyclase tapped for maize biofortification. *Science*. 2008;319:330–333
- Hulshof PJM, Schuil TK, West CE, Hollman PCH. Quick screening of maize kernels for pro

- vitamin A content. *Journal of Food Comparison and Analysis*. 2007;20: 655–61.
- Kovacevic V, Sostaric J, Josipovic M, Iljkic D, Markovic M. Precipitation and temperature regime impact on maize yields in eastern Croatia. *Journal of Agricultural Science Research*. 2009;41:49-53.
- Kumar AD, Singh NK. Carotenoid and SSR marker-based assessment among short duration maize (*Zea mays* L.) genotypes. *Maydica*. 2012;57:106-113.
- Li FQ, Vallabhaneni R, Yu J, Rocheford T, Wurtzel ET. The maize phytoene synthase gene family: Overlapping roles for carotenogenesis in endosperm, photomorphogenesis, and thermal stress tolerance. *Plant Physiology*. 2008;147: 1334–1346
- Megyes A, Nagy J, Ratonyi T, Huzsvai L. Irrigation of maize (*Zea mays* L.) in relation to fertilization in a long-term field experiment. *Acta Agronomica, Hungarica*. 2005;53: 41-46
- Menkir A, Liu W, White WS, Maziya-Dixon B, Rocheford T (2008). Carotenoid diversity in tropical-adapted yellow maize inbred lines. *Journal of Food Chemistry*. 2008;109:521–529.
- Mishra P, Singh NK. Spectrophotometric and TLC based characterization of kernel carotenoids in short duration maize. *Maydica*. 2009;55:95-100
- Mishra P, Singh NK. Spectrophotometric and TLC based characterization of kernel carotenoids in short duration maize. *Maydica*. 2010;55:95-100
- Pfeiffer WH, McClafferty B. Biofortification: breeding micronutrient-dense crops Chapter 3. In: Kang MS, Priyadarshan PM. *Breeding major food staples for the 21st century*. 2007;61-91.
- Pixley K, Rojas NP, Babu R, Mutale R, Surles R, Simpungwe E. Biofortification of maize with pro-vitamin A carotenoid. In: S.A. Tanumihardjo (Editorial). *Carotenoids and human health* SE 17. Humana Press, New York, NY. 2013;271–292.
- Randjelovic V, Prodanovic S, Tomic Z, Bijelic Z, Simic A. Genotype and year effect on grain yield and nutritive values of maize (*Zea mays* L.) *Journal of Animal and veterinary Advances*. 2011;10(7):835-840.
- Rohlf FJ. NTSYSpc, Numerical taxonomy and multivariate analysis system, version 2.0, Exeter Software, New York. 1998
- Russell JR, Fuller JD, Macaulay M, Hatz BG, Jahoor A, Powell W, Waugh R. Direct comparison

of levels of genetic variation among barley accessions detected by RFLPs, SSRs, and RAPD. *Journal of Theoretical and Applied Genetics*. 1997;95: 714-722

Sivaranjani R, Prasanna BM, Hossain F, Santha IM. Genetic variability for total carotenoid concentration in selected tropical maize (*Zea mays*) inbred lines. *Indian Journal of Agricultural Sciences*. 2013;83(4):431-6

Smale M, Byerlee D, Jayne T. Maize revolutions in sub-Saharan Africa. In. *An African green revolution SE - 8*. Otsuka K, Larson DF (Eds.). Amsterdam, Netherlands: Springer. 2013;165-195.

Wurtzel ET, Cuttriss A, Vallabhaneni R. Maize pro-vitamin A carotenoids, current resources, and future metabolic engineering challenges. *Journal of Frontiers in Plant Science*. 2012;3:29.

Smith JSC, Chin ECL, Shu H, Smith OS, Wall SJ, Senior ML, Mitchell SE, Kresovich S, Ziegler J. An evaluation of the utility of SSR loci as molecular markers in maize (*Zea mays* L): comparisons with data from RFLPs and pedigree. *Journal of Theoretical and Applied Genetics*. 1997;95:163–173.

Warburton ML, Xianchun X, Crossa J, Franco J, Melchinger AE, Frisch M, Bohn M, Hoesington D. Genetic characterization of CIMMYT inbred maize lines and open pollinated populations using large scale fingerprinting methods. *Crop Science*. 2002;42:1832–1840.