

Original Research Article

FLORISTIC CHARACTERIZATION AND PHYTOSOCIOLOGY OF A VEGETATION IN A CAATINGA AREA

ABSTRACT

The present study aimed to characterize the floristic and phytosociology in Caatinga area in the ~~p~~Paraíba semi-arid in Brazil. For the phytosociological survey of the area, was used sample units with dimensions of 20 m x 20 m (400 m²), which were distributed randomly in order to collect the name of the species, the circumference at ground level (CFL), the circumference at breast height (CBH), the total height of the individuals and the state trees. The total surface of the area was 323.65 ~~he-ha~~ and 255.85 ~~he-ha~~ of shrub by arboreal vegetation what represent 79% of the area with vegetation, the remaining 21%, were classified as clean field, reservoirs, courses of water and highways, totaling 67.8 hectares. To the floristic composition 2.362 individuals were observed, which ones, 22 species belonging to 14 families. Considering the habit of the found species, 68% can be considered as arboreal and 32% as shrubby. The vegetation can be classified as a closed shrub-arboreal Caatinga. The most representative species in the area were *Poincianella pyramidalis*, *Mimosa tenuiflora*, *Aspidosperma pyrifolium* and *Anadenanthera colubrina*.

Keywords: Measurement; forest inventory; forest management; steppic savannah biome

1. INTRODUCTION

The Caatinga biome is found in semi-arid region of northeastern states of Brazil, extending to the northeast of the state of Minas Gerais. It is estimated that the total area covered by this Biome is about 1.000.000 km² [1]. The annual precipitation in the region is less than 1000 mm a year, with rains distributed irregularly. Moreover, the solar radiation is extremely high, as the annual average temperature, while the relative levels of moisture and cloudiness are the lowest in the region of Brazil [2].

The Caatinga has an inappropriate usage history of their land, with the transformation of native forests in arable fields, large areas for livestock, among other uses. Due to this fact, 45% of the Caatinga has been modified by human activities [3]. However, studies indicate that this value is underestimated [4]. Despite the ~~immense~~ lack of knowledge about the Biome [5], the Caatinga has been systematically devastated.

Thus, this type of exploration on such a complex and unknown environment may take even an irreversible process of degradation and consequent desertification. The native forest remnants, is located almost exclusively within semi-arid, playing an important role in the state's socio-economy [6]. However, the forest cover of the semi-arid region has been drastically reduced by the lack of proper management and the type of exploration adopted.

Comment [J1]: Agroecosystems?

34 As seen, the Caatinga lost part of its native vegetation as a result of inadequate
35 management. Due to this fact, there is damage to the soil and water bodies of the region [7].

36 Much of the environmental degradation problems is related to the absence of a proper
37 planning of the occupation of the land, respecting the characteristics of the various
38 ecosystems, particularly its richness and diversity. Environmental degradation always had
39 the impulse to economic enterprises that do not consider the environmental changes in
40 costs [2].

41 Despite the existence of some phytosociological work of the Caatinga vegetation
42 composition, there is still much to knowledge of this Biome as a whole, determining their
43 distribution patterns, abundance and relationship with environmental factors, to be
44 established based on quantitative data, the different faces of the Caatinga and its floristic
45 connections [8].

46 In this sense, the aim of this study was to characterize and elaborate a phytosociological
47 and floristic diagnosis of an area located in a representative Caatinga Biome.

48
49

50 2. MATERIAL AND METHODS

51

52 2.1 Characterization of the study area

53 The area where were developed the research is part of Tamandua Farm, owned by the
54 MocoAgropecuaria Ltda. company, located in Santa Terezinha (PB). The area is located in
55 the micro-region of low Hinterland Piranhas, with altitude ranging between 250 and 310
56 meters, in the coordinates 07° 00 'S and 37° 23' W.

57 According to the Köppen climate classification, the region of the study is characterized as
58 having type climate Bsh, semiarid region, marked by rainy and dry season [9].

59 The area is part of a geomorphology unit Country Depression, an extended low plain,
60 smooth-rolling predominant relief, sometimes wavy, with residual elevations scattered in the
61 landscape, in which the granite rock shows exposed or minimal soil capping and vegetation
62 [10]. The vegetation in which the study was conducted is characterized as Steppic savannah
63 [11].

64 2.2 Environmental characterization of the area

65 The first step to characterize the environmental area was conducting a topographic survey
66 using a GPS navigation. In this phase were used an aerial photograph with high resolution of
67 the whole farm, provided by the owner. This procedure defined the perimeter of the studied
68 area. For making the maps was used the computer program TrackMaker® professional
69 version 4.7.

70 After defining the perimeter, it was raised areas with vegetation, grassland, water reservoirs,
71 waterways and roads. The representation of terrain elevation through colors, i.e. the
72 hypsometry area was made using the computer program Surfer® version 10.

73 The assessment of the ground truth was done through several visits to the area during the
74 various surveys.

Comment [J2]: To describe richer and more abundant families already sampled and reported in previous studies.

Comment [J3]: Insert a map

Comment [J4]: Use up-to-date literature. See Alvarez et al., 2013

Comment [J5]: Insert reference

Comment [J6]: Insert reference

Comment [J7]: Describe how many visits and how many samples.

75 **2.3 Floristic survey**

76 The floristic survey of the shrub-tree layer was carried out through periodic campaigns of
77 field, in order to go through the greatest extensions of the study area, with a view to observe
78 the greatest possible number of established plant species. In addition to the campaigns, they
79 were held collections in the plots used for phytosociological. To collect the plant material
80 was used pruning shears. The collected material was properly prepared and forwarded to
81 the Herbarium of the Academic Unit of Forestry (CSTR-UFCEG) for identification.

82 For the classification of the species adopted the [12]. As for the floristic comparisons, we
83 used different work carried out in Caatinga vegetation.

84 **2.4 Phytosociological survey**


85 For the phytosociological survey of the area, was used sample units with dimensions of 20
86 m x 20 m (400 m²), which were distributed randomly. In each unit has been noted, the
87 common name, the circumference at ground level (CFL), the circumference at breast height
88 (CBH), the total height of the individuals and the state trees, alive or dead. Were considered
89 for data collection purposes, all living or dead, still standing, and as inclusion criteria, total
90 height greater than 1.0 meters, and the largest circumference at chest level or equal to 6.0
91 cm, following the recommendation of the Forest Management Network Measurement
92 Protocol of Caatinga [13].

93 Circumferences were measured with the aid of tape and to the heights was used retractable
94 metal rod, graduated in meters with divisions of 50 cm.

95 **3. RESULTS AND DISCUSSION**

96 **3.1 Characterization of the study area**

97
98
99 | The total area of the studied was 323.65 ~~he~~ha. It was also found that the total study area,
100 255.85 he showed shrub and tree vegetation, which represents 79% of the area with
101 vegetation. After the 25 sampling units used in the characterization of vegetation in the
102 experimental area the definition of areas with vegetation were distributed. The remaining
103 21% after aerial photography analysis, were classified as areas of grassland, reservoirs,
104 | water ways and roads, totaling 67.8 ~~he~~ha (Fig. 1).




106
107 **Fig. 1. Map of the study area with vegetation (white), Grassland (yellow), bodies of**
108 **water (blue) and distribution of sample units.**
109

Comment [J8]: Inform: if the areas sampled have a history of deforestation and, if so, at what stage of regeration they are; b) how far from the border with cultivated areas, highways, dwellings ... the areas sampled meet


Comment [J9]: Move to the Material and Methods section

110 The hypsometric chart (Fig. 2) shows that the altitude, ranging from 230 m to 320 m. It can
 111 be observed that these altitudes are commonly found throughout the region.
 112



113
 114
 115
 116 **Fig. 2. Hypsometric chart map of the study area.**
 117

118 The sampling sufficiency was verified by the collector curve (Fig. 3). There was an initial
 119 increase of trend and, to the extent that the sampled area increased, tended to stabilize
 120 indicating that sampling of the species in the experimental area was enough. The curve
 121 tended to stabilize with 9600-10000 m² of area sampled, indicating that sampling was
 122 satisfactory for the species in the study area.
 123



124
 125
 126 **Fig. 3. Adequate-sSampling effort of the species in the study area.**
 127

128 Regarding the floristic composition, were found a total of 2.362 individuals, including 22
 129 species belonging to 14 families (Table 1). However, one of the observed species was not
 130 identified botanically, and that at the time of collection of botanical material, it had no fertile
 131 material. The percentage of dead individuals was 4%. Whereas the habit of the species
 132 found, 68% can be considered as a tree and shrub 32%.
 133
 134
 135
 136

Comment [J10]: Move to the Material and Methods section ver para a seção de material e métodos

Comment [J11]: Method, subsection data analysis!

Comment [J12]: Format figure: Increase font size; increase image resolution; exclude internal lines;

137 **Table 1. Floristic ratio of woody species in the study area.**

| Family/Specie | Common name | Habit |
|---------------------------------|------------------|----------|
| Anacardiaceae | | |
| <i>Myracrodrum urundeuva</i> | Aroeira | Arboreal |
| Apocynaceae | | |
| <i>Aspidosperma pyrifolium</i> | Pereiro | Arboreal |
| Bignoniaceae | | |
| <i>Tabebuia impetiginosa</i> | Pau d'arco | Arboreal |
| Bombacaceae | | |
| <i>Pseudobombax marginatum</i> | Embiratanha | Arboreal |
| Burseraceae | | |
| <i>Commiphora leptophloeos</i> | Imburana | Arboreal |
| Capparaceae | | |
| <i>Capparis flexuosa</i> | Feijao bravo | Shrub |
| Combretaceae | | |
| <i>Combretum leprosum</i> | Mofumbo | Shrub |
| Erythroxylaceae | | |
| <i>Erythroxylum pungens</i> | Rompegibao | Shrub |
| Indeterminada | | |
| Indeterminada | Cipo de vaqueiro | Shrub |
| Fabaceae | | |
| <i>Amburana cearensis</i> | Cumaru | Arboreal |
| <i>Bauhinia cheilantha</i> | Mororo | Arboreal |
| <i>Poincianella pyramidalis</i> | Catingueira | Arboreal |
| <i>Libidibia ferrea</i> | Pau ferro | Arboreal |
| <i>Senna macranthera</i> | Sao Joao | Shrub |
| Mimosaceae | | |
| <i>Anadenanthera colubrina</i> | Angico | Arboreal |
| <i>Mimosa tenuiflora</i> | Jurema preta | Arboreal |
| <i>Piptadenia stipulacea</i> | Jurema branca | Arboreal |
| Rhamnaceae | | |
| <i>Ziziphus joazeiro</i> | Juazeiro | Arboreal |

Comment [J13]: Add author

Comment [J14]: Add author

Comment [J15]: Add author

Comment [J16]: Add author

Comment [J17]: Add author

Comment [J18]: Add author

Comment [J19]: Add author

Comment [J20]: Add author

Formatted: Spanish (Colombia)

Comment [J21]: Add author

Comment [J22]: Add author

Comment [J23]: Add author


Comment [J24]: Add author

138

139 Among the ten species that stood out in relation to the importance value, we can highlight
 140 the catingueira (*Poincianella P. pyramidalis*), in which it had the highest importance value
 141 (Fig. 4), which was mainly due to the high number of individuals of species in the
 142 experimental area [14-15].

Comment [J25]: Abbreviate gender after first citation. In all cases below

143



Comment [J26]: Format picture. Increase the font size; increase the resolution

144

145

146

Fig. 4. Importance value (IV) of the ten species that showed the highest.

147

148 | Considering the horizontal structure, it was observed that once again the *Poincianella-P.*
 149 *pyramidalis* specie stands when comparing phytosociological parameters with other species
 150 (Table 2).

151 **Table 2. Values of phytosociological parameters for the species sampled.**

| Cientific name | DR | FA | FR | DoR | VI |
|---------------------------------|------|-------|------|-------|------|
| <i>Amburana cearensis</i> | 0.08 | 8.00 | 0.87 | 0.04 | 0.33 |
| <i>Anadenanthera colubrina</i> | 3.30 | 64.00 | 6.99 | 10.14 | 6.81 |
| <i>Aspidosperma pyrifolium</i> | 8.98 | 76.00 | 8.30 | 10.11 | 9.13 |
| <i>Bauhinia cheilantha</i> | 2.03 | 36.00 | 3.93 | 0.59 | 2.19 |
| <i>Capparis flexuosa</i> | 0.08 | 8.00 | 0.87 | 0.03 | 0.33 |
| <i>Cnidocolus quercifolius</i> | 0.68 | 32.00 | 3.49 | 1.40 | 1.86 |
| <i>Combretum leprosum</i> | 7.11 | 76.00 | 8.30 | 2.47 | 5.96 |
| <i>Commiphora leptophloeos</i> | 2.24 | 52.00 | 5.68 | 5.02 | 4.31 |
| <i>Croton blanchetianus</i> | 26.7 | 96.00 | 10.5 | 6.58 | 14.5 |
| <i>Croton cf. alagoensis</i> | 0.55 | 12.00 | 1.31 | 0.15 | 0.67 |
| <i>Erythroxylum pungens</i> | 1.65 | 44.00 | 4.80 | 1.70 | 2.72 |
| <i>Indeterminada</i> | 0.34 | 12.00 | 1.31 | 0.13 | 0.59 |
| <i>Libidibia ferrea</i> | 0.34 | 4.00 | 0.44 | 0.15 | 0.31 |
| <i>Mimosa tenuiflora</i> | 8.93 | 76.00 | 8.30 | 11.29 | 9.51 |
| <i>Morta</i> | 4.02 | 84.00 | 9.17 | 4.97 | 6.05 |
| <i>Myracrodrum urundeuva</i> | 0.13 | 8.00 | 0.87 | 0.12 | 0.37 |
| <i>Piptadenia stipulacea</i> | 2.92 | 76.00 | 8.30 | 1.73 | 4.32 |
| <i>Poincianella pyramidalis</i> | 29.1 | 100.0 | 10.9 | 40.89 | 26.9 |
| <i>Pseudobombax marginatum</i> | 0.25 | 24.00 | 2.62 | 0.85 | 1.24 |
| <i>Sebastiania sp.</i> | 0.08 | 4.00 | 0.44 | 0.05 | 0.19 |
| <i>Senna macranthera</i> | 0.17 | 8.00 | 0.87 | 0.02 | 0.36 |
| <i>Tabebuia impetiginosa</i> | 0.08 | 8.00 | 0.87 | 0.19 | 0.38 |
| <i>Ziziphus joazeiro</i> | 0.17 | 8.00 | 0.87 | 1.38 | 0.81 |

Comment [J27]: Only scientific names in italics. In this case translate into English

Comment [J28]: Same before comment

152 DR = relative density (%); FR = relative frequency; DoR = relative dominance (%); VI = importance
 153 value (%).

154 It was observed that the ordering of the 22 sampled species by their importance values
 155 followed, mainly, the relative dominance (Table 2), indicating that the basal area of the
 156 individuals was essential to the determination of IV of the species, as was the case
 157 *Poincianella-P.pyramidalis*, *Mimosa-M.tenuiflora*, *Anadenanthera-A.colubrina* and
 158 *Aspidosperma-A.pyrifolium* decreasingly.

159 Of the 2.362 sampled individuals, 2.232 individuals are distributed in the first three classes,
 160 representing approximately 94.5% of all samples with diameters less than 15 cm. (Table 3).
 161 The estimated cylindrical volume was approximately 52.1 m³/heha.


162 **Table 3. Diametric distribution of individuals (N), percentage of individuals, basal area**
 163 **(BA) and volume per hectare (V/he), according to the DAB (diameter at the base).**

| Class (cm) | N | % | BA m ³ /heha | V/heha |
|------------|------|-------|-------------------------|--------|
| 0.0 – 5.0 | 1148 | 48.60 | 1.2327 | 3.5908 |
| 5.0 – 10.0 | 844 | 35.73 | 3.2361 | 11.997 |

| | | | | |
|--------------|-------------|------------|---------------|---------------|
| 10.0 – 15.0 | 240 | 10.16 | 2.8042 | 12.169 |
| 15.0 – 20.0 | 82 | 3.47 | 1.8899 | 9.6911 |
| 20.0 – 25.0 | 31 | 1.31 | 1.2252 | 6.2648 |
| 25.0 – 30.0 | 11 | 0.47 | 0.6681 | 3.9533 |
| 30.0 – 35.0 | 3 | 0.13 | 0.2519 | 1.7177 |
| 35.0 – 40.0 | 2 | 0.08 | 0.2274 | 1.6271 |
| 40.0 – 45.0 | 1 | 0.04 | 0.1450 | 1.0877 |
| Total | 2362 | 100 | 11.680 | 52.099 |

164

165 The basal area average was found to be 11.68 m²/ha and is considered a low value in
 166 relation to other studies conducted in semiarid region. The species that had higher basal
 167 area were *Poincianella*—*P.pyramidalis*, *Mimosa*—*M.tenuiflora*, *Anadenanthera*
 168 *A.colubrina* and *Aspidosperma*—*A.pyrifolium* (Fig. 5).




169

170 **Fig. 5. Relation of the ten species that showed higher basal area (m²).**

171 Structurally, the following vegetation characteristics were evident: the majority of individuals
 172 in the class 3.0 to 4.5 m high, with some emerging species, reaching about 10 m or more,
 173 with relevant importance in community structure, as equal among the most IV, as
 174 *Poincianella*—*P.pyramidalis*, *Mimosa*—*M.tenuiflora* and *Anadenanthera*—*A.colubrina* due to
 175 significant numbers of individuals and basal area (Fig. 6). It was also clearly visible two
 176 strata, one tree and other shrubs. In the tree layer, with a predominance of individuals with
 177 height greater than 4.5 m.

178



179

180 **Fig. 6. Percentage of individuals in classes (cm).**

181 **4. DISCUSSION**

Comment [J29]: Format figure: increase font size and resolution

Comment [J30]: Where and when? Add!

Comment [J31]: Format figure: increase font size and resolution

Comment [J32]: 48/5000
Individuals of what? Where and when? It is not clear

182 The results of this study show that there is considerable species richness in the area, even if
183 compared with inventories in other areas of Caatinga [15-17].

184 In the arboreal component occurred 15 species, distributed in nine families, especially
185 Mimosaceae, Fabaceae and Euphorbiaceae, the first family with four species and the last
186 two with three and two species, respectively, and the other families with only one specie.

187 In the shrub component occurred seven species, distributed in six families. Euphorbiaceae,
188 with two species of shrubs, being considered the most rich family ~~wealth~~ in this component.

189 Without considering the habit of the species is observed that families with highest species
190 richness were Fabaceae with five species, followed by Euphorbiaceae families with four
191 species and Mimosaceae with three species. The other families had only one specie. Similar
192 results were obtained in studies performed in the Caatinga of the Serido region [18-19].
193 Other research carried out within the same farm was noted that families with the highest
194 number of species were Euphorbiaceae, Caesalpinaceae and Mimosaceae [20].

195 The high species richness of Fabaceae in the study area indicates their status as the main
196 family [21]. Leguminous are constantly seen in inventories in areas of savannah, as in
197 crystalline soffits [22,16], sandy soils of sedimentary basins [23], or in areas with a
198 predominance of rocky outcrops [24], or surveys restricted to the woody component [25]
199 which include other habits [15].

200 The species that have held the highest number of individuals sampled were
201 *Poincianellapyramidalis*, *Croton blanchetianus* and *Aspidospermapyrifolium*. The
202 *Poincianellapyramidalis* species and *Croton sonderianus* are considered those that stood
203 out most in the number of individuals in several works in areas of Caatinga [26-27].

204 The five species that had higher relative density were *Poincianellapyramidalis*, *Croton*
205 *blanchetianus*, *Aspidospermapyrifolium*, *Mimosa tenuiflora* and *Combretumleprosum*. The
206 forest specie *Poincianellapyramidalis* showed higher relative dominance. Other studies, find
207 in an area of Caatinga in good condition, the *P. pyramidalis* has the second highest
208 importance value [28]. The prevalence of *P. pyramidalis* on other species is due to the large
209 number of individuals (688) and also by the way these individuals are distributed throughout
210 the sampled area. The species were present in 100% of sampled plots, thus affecting
211 substantially the highest importance value.

212 With regard to the relative frequency, the five species that stood out were:
213 *Poincianellapyramidalis*, dead trees, *Aspidospermapyrifolium*, *Mimosa tenuiflora* and
214 *Combretumleprosum*, and the last three species showed similar percentages (8.3%).

215 It can be seen that the majority of individuals (94%) were positioned in the first three
216 diameter classes, numerically decreasing with increasing the diameter, to form a graphic
217 format of inverted "J". Thus, in the event of a disturbance on vegetation, older individuals
218 (larger diameter class), in a less quantity, may die younger individuals, along with
219 regenerating quickly repopulate the disturbed area. The presence of many individuals with
220 stem diameter in the initial diameter classes shows an initial stage of secondary feature by
221 the studied vegetation [29].

222 The high diversity of species found in the study area may have several causes, such as the
223 presence of large flagstones, microhabitats in the region, numerous banks of rivers and
224 streams. Furthermore, several studies indicate that these environments serve as refuges,
225 and to minimize the effects of drought, it enables a higher probability of survival of various

226 species. Moreover, the vegetation conservation state may have contributed to the high
227 richness [26, 30].

228 5. CONCLUSIONS

229
230 The most representative species in the area were *Poincianellapyramidalis*, *Mimosa*
231 *tenuiflora*, *Aspidospermapyrifolium* and *Anadenanthera colubrina*.

232 The most important families with the number of species sampled were Mimosaceae,
233 Fabaceae and Euphorbiaceae.

234 The species *Poincianellapyramidalis* showed the highest rates recorded for the horizontal
235 structure, vertical, basal area and volume.

236 Regarding the characterization of vegetation, it is clear that it can be classified as a closed
237 shrub-arboreal Caatinga.

238 COMPETING INTERESTS

239 Authors declare that is no competing interests.
240
241

242 REFERENCES

- 243
244 1. Queiroz MA. Caatinga Plant Genetic Resources for the Development of the Brazilian
245 Semi-Arid (Caatinga Plant Genetic Resources for the Development of the Brazilian
246 Semiarid). *Revista Brasileira de Geografia Física*. 2012; 4 (6): 1135-1150.
247
248 2. Souza BI, Menezes R, Artigas RC. Effects of desertification on species composition of the
249 Caatinga biome, Paraíba / Brazil. *Geographical Research*. 2015; 88 (1): 45-59. DOI:
250 <https://doi.org/10.14350/rig.44092>. English.
251
252 3. MMA - Ministry of the Environment. Monitoring of Brazilian biomes: Caatinga biome.
253 Brasília: MMA; 2010. English.
254 4. Castelletti CHM. How much is left of the caatinga? A preliminary estimate. In: LEAL, IR,
255 TABARELLI M, SILVA JMC. Ecology and Conservation of the Caatinga. Recife, UFPE
256 University Ed.; 2003. English.
257
258 5. Moro MF, Araújo FS, Rodal MJN, Martins FR. Synthesis of the floristic and
259 phytosociological studies carried out in the Brazilian semiarid region. In: Eisenlohr PV, Felfili
260 JM, Melo MMRF, and Meira Neto JAA. Phytosociology in Brazil: Methods and case studies.
261 Vol. II. Publisher of the Federal University of Viçosa, Viçosa; 2015. English.
262
263 6. Silva KA, Santos JMFF, Santos DM, Ferraz EMN and Araújo EL Spatial variation in the
264 structure and composition of the herbaceous community in the semiarid region of
265 northeastern Brazil. *Brazilian Journal of Biology*. 2013; 73 (1): 135-148. DOI:
266 <http://dx.doi.org/10.1590/S1519-69842013000100015>
267
268 7. Junior LRP, Alberici PA, Kallianna DA. Floristic composition and phytosociology of a
269 caatinga fragment in Monteiro, Paraíba. *Holos*. 2013; 6 (1): 73-87. English.
270
271 8. JTC Jr., Drumond MA. Comparative study of the phytosociological structure of two
272 Caatinga fragments at different levels of conservation. *Brazilian Forest Research*. 2014; 34
273 (80): 345-355.

Comment [J33]: After all, what is that for? It is necessary to say if the areas serve or not for the conservation of the biodiversity; whether or not they can be exploited; whether or not the study serves as a reference for management and conservation studies in the region. The results should be presented in the light of their regional meanings for conservation and science.

Formatted: Font: 11 pt, Bold, English (United States), All caps

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

- 274
275 9. Insa, National Institute of the Semi-Arid. Accessed 8 January 2013. Available:
276 <[http://www.insa.gov.br/grupodepesquisala_vouraxerofila / index.php? Option = com_cont](http://www.insa.gov.br/grupodepesquisala_vouraxerofila/index.php?Option=com_cont)
277 and nteview = articleid = 56&temid = 62&lang = en>. English.
278
279 10. Sudema. Update of the forest diagnosis of the State of Paraíba. João Pessoa; 2004.
280 Portuguese.
281
282 11. Ibge. Map of vegetation of Brazil. IBGE, Rio de Janeiro; 1993. English.
283
284 12. Cronquist A. An integrated system of classification of flowering plants. New York,
285 Columbia University Press; 1981.
286
287 13. Forest Management Network of the Caatinga: Protocol of measurements of permanent
288 plots / Scientific Technical Committee. Recife: Associação Plantas do Nordeste; 2005.
289 English.
290
291 14. Costa GM, Cardoso D, Queiroz LP, Conceição AA. Local variations in floristic richness in
292 two caatinga core regions. Rodriguésia, Botanical Garden Research Institute of Rio de
293 Janeiro. 2015; 66 (3): 685-709. DOI: 10.1590 / 2175-7860201566303. English.
294
295 15. Pinto MS, Sampaio EV, Nascimento LM. Floristic and vegetation structure of an altitude
296 brejo in fishing, PE, Brazil. Northeastern Journal of Biology. 2012; 21 (1): 47-79. English.
297
298 16. WJ Ax, Silver APN, Mello AA. Floristic composition in areas of Caatinga and Brejo de
299 Altitude in Sergipe state, Brazil. Check List, 2012; 8 (1): 1089-1101. DOI:
300 <http://dx.doi.org/10.15560/8.6.1089>
301
302 17. Ferreira EVR. Floristic composition, community structure and seed dispersal syndrome
303 of a Caatinga remnant in Poço Verde - Sergipe. Dissertation Master. São Cristóvão: Federal
304 University of Sergipe, 2011. 81p. English.
305
306 18. Oliveira DG, Silver AP, Ferreira RA. Herbaceous of the Caatinga: floristic composition,
307 phytosociology and strategies of survival in a plant community. Brazilian Journal of
308 Agricultural Sciences. 2013; 8 (4): 623-633. DOI:
309 <http://dx.doi.org/10.15628/holos.2012.1188>. English.
310
311 19. Andrade-Lima D. The caatinga dominium. Brazilian Journal of Botany, 1981; 4 (1): 149-
312 153.
313
314 20. Araújo LVC, Paulo MCS, Paes JB. Dendrometric characteristics and basic density of the
315 Jurema-preta (*Mimosa tenuiflora* (Willd.) Poir.) From two regions of the state of
316 Paraíba. Caatinga Magazine. 2007; 20 (1): 89-96. Portuguese.
317
318 21. Cardoso DBOS, Queiroz LP. Caatinga in the context of a community goal: evidence of
319 biogeography, phylogenetic patterns and abundance of species in Leguminosae. In:
320 Carvalho CJB.; Almeida EAB. (eds.). Biogeography of South America: patterns and
321 processes. Roca, São Paulo; 2011. English.
322
323 22. Lima IB, Barbosa VMR. Floristic composition of RPPN farms farm, in Paraíba, Paraíba,
324 Brazil. Northeastern Journal of Biology. 2014; 23 (1): 49-67. English.
325

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

- 326 23. Rocha PLB, Queiroz LP, Pirani JR. Plant species and habitat structure in a sand dune
327 field in the Brazilian Caatinga: A homogenous habitat harbouring an endemic biota. *Brazilian*
328 *Journal of Botany*. 2004; 27 (1): 739-755. DOI: [http://dx.doi.org/10.1590/S0100-](http://dx.doi.org/10.1590/S0100-84042004000400013)
329 [84042004000400013](http://dx.doi.org/10.1590/S0100-84042004000400013)
330
- 331 24. Gomes P, Alves M. Floristic diversity of two crystalline rocky outcrops in the Brazilian
332 northeast semi-arid region. *Brazilian Journal of Botany*. 2010; 33 (1): 661-676. DOI:
333 <http://dx.doi.org/10.1590/S0100-84042010000400014>
334
- 335 25. We read JR, Rodal MJN. Phytosociology of the woody component of a section of the
336 caatinga vegetation in Serra da Capivara National Park, Piauí, Brazil. *Acta Botânica Brasília*.
337 2002; 16 (1): 23-42. English.
338
- 339 26. Lima IB, Barbosa VMR. Floristic composition of RPPN farms farm, in Paraíba, Paraíba,
340 Brazil. *Northeastern Journal of Biology*. 2014; 23 (1): 49-67. English.
341
- 342 27. AraújoFilho JA, Barbosa TML. Agroforestry management of the caatinga: a proposal of a
343 production system. In OLIVEIRA, TS., ROMERO, RE., ASSIS JUNIOR, RN, SILVA JRCS.
344 (Eds.). *Agriculture, sustainability and the semi-arid*. Fortaleza: Brazilian Society of Soil
345 Science; Federal University of Ceara; 2000. English.
346
- 347 28. Andrade LA, Pereira IM, Leite UT, Barbosa MRV. Analysis of the cover of two
348 caatingaphytophysognomies, with different use histories, in the city of. State, Brazil. *Cerne*
349 *Magazine*. 2005; 11 (3): 253-262. English.
350
- 351 29. Junior LRP, Alberício PA, Kallianna DA. Floristic composition and phytosociology of a
352 caatinga fragment in monteiro, paraíba. *Holos*. 2013; 6 (1): 73-87. English.
353
- 354 30. Albuquerque UP, Araújo EL, El-Deir ACA, Lima ALA, Souto A, Bezerra BM, Ferraz EMN,
355 Freire EMX, Sampaio VSB, Las-Casas FMG, Moura GJB, Pereira GA, Melo JG, RAMOS
356 MA, Rodal MJN, Schiel N, Lyra NRM, Alves RRN, Azevedo-Júnior SM, Telino-Júnior WR,
357 Severi W. Caatinga Revisited: Ecology and Conservation of an important seasonal dry
358 forest. *The Scientific World Journal*. 2012; 205182(1):1-18. DOI:
359 <http://dx.doi.org/10.1100/2012/205182>

Formatted: English (United States)