

Original Research Article
**Development Of Young Plants Of Pumpkin
(*Cucurbita moschata*) In Different Sources And
Dosages Of Phosphorus**

ABSTRACT

The research aimed to analyze the effect of different dosage and sources of phosphorus in growth and development of young plants of pumpkins (*Cucurbita moschata*) 'jacarezinho' cultivar. The experimental design utilized was the completely randomized with the factorial scheme 5x3 constituted by five dosages of P₂O₅ (0; 22.5; 45; 67.5 e 90 kg.ha⁻¹) and three phosphate sources (Simple superphosphate, Triple superphosphate and Natural phosphate of ARAD) with three repetitions. The experiment was performed in a greenhouse at field area at Federal Rural University of Amazônia in the Capitão Poço City, Pará State. The research was carried out from March to April, year of 2016. The analyzed variables were plant height (PH), stem diameter (SD), number of leaves (NL), fresh root mass (FRM), fresh aerial mass (FAM), dry aerial mass (DAM) and dry root mass (DRM) at 30 days after sowing. The average of dosages x sources interactions were submitted the regression analyze (P<0.05). There was a significant response to the interaction dosage x source for the PH, FAM and DAM. However, for the SD, there was a significant response only to the dose factor, with linear response. The dosage at 0 t ha⁻¹ showed the **worst/least** response in all variables. ARAD was the source that had the **worst** performance in the analyzed variables in comparison to others sources. Young plants of pumpkins were significantly influenced by phosphate fertilization.

Keywords: *Agriculture, Cucurbita moschata; Phosphate fertilization; vegetables.*

1. INTRODUCTION

Between the cultivated species of the Cucurbitaceae's family, the pumpkin (*Cucurbita moschata*) occupies a relevant position in agribusiness, being one of the most consumed vegetables in Brazil [1]. This vegetable presents a fruit of globular shape, with average weight of 2 kg to 3 kg, herbaceous stem, large dark green leaves. Its pulp is yellow, rich in nutrients as vitamin A and B complex, including calcium, iron magnesium, potassium and zinc [2]. Also presents good quality for consumption, conservation and flavor [3].

According to [4] the world production of pumpkins has grown from 21.4 million tons in 2006 to 25.2 million tons in 2014 with an average yield of $13.4 \text{ t}\cdot\text{ha}^{-1}$. In Brazil, there are few data about pumpkin's commercialization wherein the latest information available is from 2006 that showed more than 127,000 farms harvested 84 thousand hectares of pumpkins (5.24% of the world harvested area), they produced 398 thousand tons (1.86% of world production), with average yield of $4.5 \text{ t}\cdot\text{ha}^{-1}$, below the world average (12.5 to $13.5 \text{ t}\cdot\text{ha}^{-1}$) [5].

Vegetables require supply of nutrients in quantities satisfactory for their perfect growth and development that result in adequate yields. One of the factors that cause a decrease in productivity is unbalanced mineral nutrition that directly influences the production and the final quality of the harvest.

Phosphorus (P) is one of nutrients more demanded by plants due to its performance in plant metabolism. This element plays a role in root growth [6-7], metabolic processes such as ATP's and nucleic acids' formations and photosynthesis [8-9] that result in high quality of the crops. However, the majority of Brazilian soils have low natural fertility and high phosphorus fixation capacity [10-11].

There are a few researches about/on the effect of phosphorus dosage and source of this nutrient in development and yield of vegetables. [12] when applied P_2O_5 dosage between 245 and 284,6 $\text{mg}\cdot\text{dm}^{-3}$ of Simple/single superphosphate (SS) in radish crops cultivated at Red Latosol verified higher plant height and dry aerial mass. [13] have realized a research with phosphorous omission at nutritive solution in beet crop and they found high purplish on the leaves and roots, low plants growth and paralyze of roots growth. [14] have observed that dosage of $340 \text{ mg}\cdot\text{dm}^{-3}$ from Triple Superphosphate (TSP) at Dystrophic Red Latosol has caused increase of P in the leaves and fresh root mass in beet crop.

According to [15] phosphorus limitations at the beginning of vegetative cycle can result in limitations in roots and aerial part development that the plants may not recover even increasing the P supply. Although, the high importance about phosphate fertilizing there is a lack of knowledge in relation to phosphorus necessities in young plants of pumpkins in

Brazilian Amazon region. Therefore, the determination of ideal dosage for this crop is very important to seedling production system.

Another important aspect is the best choice of phosphate sources to be used because the inadequate choice can be promote yield reduction. It knows there are many phosphate sources can be used in agriculture, such as, Simple and Triple superphosphate. [16] verified those soluble phosphates, as Simple? Single and Triple superphosphate show more performance at phosphorus supply at annual crops.

Besides soluble sources, there are natural phosphates that are insoluble in water, show slow liberation of P in soil solution and give more P supply to the plants overtime. [17] report that organic agriculture principles do not allowed the use of soluble fertilizers made of industrial processes. The natural phosphates and thermophosphate? thiophosphate are options that may be used in phosphate fertilizers in agriculture systems.

Thus, it is important news performance about this theme with objective to find news viable strategies for the farmers. The research aimed to analyze the effect of different dosage and sources of phosphorus in growth and development of young plants of pumpkins (*Cucurbita moschata*) 'jacarezinho' cultivar.

2. MATERIAL AND METHODS

The experiment was performed in a greenhouse at field area at Federal Rural University of Amazônia, Capitão Poço Campus, in the Capitão Poço City, Pará State (1°44'39" S; 47°3'26" W, altitude 73 m). The research was carried out from March, 24 to April, 24, year of 2016. The soil used in this research is classified as Dystrophic Yellow Latosol [18] and showed the following chemical characteristics: pH (water) = 4.9, M.O = 7.86 g.kg⁻¹, P = 3,0 mg.dm⁻³, K = 15,0 mg.dm⁻³, Na = 8.0 mg.dm⁻³, Ca = 0.6 cmol.dm⁻³, Ca + Mg = 0.8 cmol.dm⁻³, Al = 0.8 cmol.dm⁻³, H + Al = 3.96 cmol.dm⁻³. According to the Köppen classification, the regional climate is classified as the Am type (tropical altitude).

We used pumpkins seeds of 'jacarezinho' cultivar and to overcome their dormancy were used soaking in distilled water for a period of 24 hours. After this step, five seeds per pot with 2 L capacity were sown.

The experimental design utilized was the completely randomized design (CRD) with the factorial scheme 5x3 constituted by five dosages of P₂O₅ and three phosphate sources with three repetitions those results in 45 experimental units. The dosages tested were 0; 22.5; 45; 67.5 e 90 kg.ha⁻¹ and the sources were Simple superphosphate (SS) with 18% of P₂O₅, Triple superphosphate (TSP) with 44% of P₂O₅ and Natural Phosphate of ARAD (ARAD) with 33% of P₂O₅. The irrigation was applied once a day to keep the soil moisture nearby the field capacity, by use of manual watering can. The choice of P₂O₅ dosages was realized/obtained according to Fertilization Manual of Pará State that indicates the dosage of 90 kg.ha⁻¹ de P₂O₅ for pumpkins [19].

Sowing was done on 24th March, after 10 days of sowing, the plants that showed less vigor were cut off, allowing the ones that are only uniform in aerial growth leaving one plant per pot. In March 24, was performed the sowing and in the tenth day was realized the cutting of young plants showed less vigorous, following a criterion of uniformity of aerial part, leaving one plant per pot. The following parameters were carried out 30days after sowing

(DAS). At 30 days after sowing were carried out the following growth and development variables: plant height (PH), was measured from the surface of the soil to the top of the plant with a millimeter ruler, stem diameter (SD), was measured with use of a pachymeter at 5 cm above the ground, number of leaves (NL), were performed by simple count, fresh root mass (FRM), obtained by weighing the roots in a precision balance and fresh aerial mass (FAM), obtained by weighing leaves + stem, in a precision balance. Also we analyzed the dry aerial mass (stem + leaves) (DAM) and dry root mass (DRM) by using a forced air circulation oven at a temperature of 65 °C until a constant mass and then the material was weighed on precision digital balance.

The data were evaluated using the Shapiro-Wilks ($P > 0.05$) and Levene ($P > 0.05$) tests to verify the normality and homoscedasticity assumption. Next, they were submitted at variance analyze and the average were compared by F test ($P < 0.05$) by using SISVAR software [20]. Finally, the average of dosages x sources interactions were submitted the regression analyze ($P < 0.05$) [21].

3. RESULTS AND DISCUSSION

The variance analyze shows that there was significant effect of dosage x sources interaction for the plant height (PH) ($P < 0.05$). But for number of leaves (NL) and stem diameter (SD) there were significant effect just for the isolate factors (Table 1).

Table 1. Summary table of variance analysis of the mean squares (MS) for plant height (PH) number of leaves (NL), stem diameter (SD) of young plants of pumpkins in function of dosages and sources of phosphorus

F.V	DF	Mean Squares (MS)		
		PH	NL	SD
DOSAGES	4	29.02**	17.41**	0.01*
SOURCES	2	17.25**	12.35**	0.00 ^{NS}

DOSAGES*SOURCES	8	6.36*	2.24 ^{NS}	0.01 ^{NS}
Error	30	2.60	1.24	0.00
CV (%)		16.24	19.61	11.44

*Significant at the 5% level of error probability, **Significant at the 1% probability level of the error and NS Not significant at the 0.05 level of probability by the F test. F.V: sources of variations; DF: degrees of freedom.

In relation to plant height (PH) the quadratic model was verified to Simple superphosphate source with the worse at 0 kg.ha⁻¹ de P₂O₅ that resulted in plants height between 2 and 2.5 cm. While the best dosage was at 57.25 kg.ha⁻¹ de P₂O₅ that provided plant height up to 3.6 cm. Plants submitted dosages more than 57.25 kg.ha⁻¹ showed quickly decrease of this variable (Figure 1).




Figure 1. Height plant of young plants of pumpkins in function of dosages and sources of phosphorus

This response can be due to chemical composition of fertilizers, because the SS contain 18% soluble P_2O_5 , 26% CaO and 12% S. So, this source give to soil, beyond P and Calcium, the Sulphur, that allow the formation of agricultural plaster that may realized soil correction that results in improving of root systems [22].

Similarly [12] observed significant response to height plant with increase of dosage of Simple? Superphosphate at Dystrophic Yellow Latosol in radish. The authors found an increase of 61.7% ($251.32 \text{ mg.dm}^{-3}$) in height plant in comparison to plants without phosphorus supply. We found adjustment linear to Triple superphosphate that means increase of PH in different level of P_2O_5 . On the other hand, we did not observe statistical difference in ARAD sources for this variable (Figure 1).

The absence of a response from the ARAD source, may be related to its main characteristic is to have slow release of P_2O_5 . Therefore, its use is more frequent in perennial species such as fruit trees [23].

However, [24] argue that in recent years, the use of alternative sources, such as phosphate rocks, has increased, although this product has low solubility in water when compared to soluble phosphates. What justifies the use of this source in the present research, it is possible to have different results if there had been more research time.

The response of number of leaves (NL) was adjusted to the quadratic model, where the ideal dosage was 62.5 kg.ha^{-1} , obtaining 6.78 leaves per plant. For the stem diameter (SD), the linear model was adjusted that showed the increase of SD was directly proportional to the level of P_2O_5 (Figures 2A and 2B).




Figure 2. Number of Leaves (A) and stem diameter (B) of young plants of pumpkins in function of dosages and sources of phosphorus

The lower results observed in low doses of phosphorus can be resulted the phosphorus deficiency that affects the plant growth, because it causes less leaf emission, which reduces the leaf area, consequently, limits the capture of the solar radiation and results in lower production of photoassimilates [25].

According to [26], the low availability of P in the soil is a limiting factor for the development of the crop, because this nutrient, besides being a constituent of the nucleic acids, also has great importance in the storage and transfer of energy, being indispensable for good development of crops.

There was no significant response of the interaction of the treatments to fresh root mass and dry root mass. However, there was statistic interaction for fresh and dry aerial mass (Table 2).


Table 2. Summary table of variance analysis of the mean squares (MS) for fresh root mass (FRM), fresh aerial mass (FAM), dry root mass (DRM) e dry aerial mass (DAM) of young plants of pumpkins in function of dosages and sources of phosphorus

F.V	DF	Mean Squares			
		FRM	FAM	DRM	DAM
DOSAGES	4	57.31**	50.23**	0.35 ^{NS}	4.05**

SOURCES	2	16.99 ^{NS}	49.46 ^{**}	0.08 ^{NS}	5.62 ^{**}
DOSAGES*SOURCES	8	5.16 ^{NS}	6.92 ^{**}	0.28 ^{NS}	0.60 ^{**}
Error	30	6.81	1.88	0.28	0.19
CV (%)		38.86	18.70	12.19	25.61

**Significant at the 5% level of error probability, **Significant at the 1% probability level of the error and NS Not significant at the 0.05 level of probability by the F test. F.V: sources of variations; DF: degrees of freedom.*

It can be noticed that the production of fresh root mass (FRM) was influenced by the dosages of phosphorus (P_2O_5), and presented a quadratic adjustment. This results shows the absence of the phosphate fertilization had severe consequences for the development of the root system. The best result was obtained with the $56.12 \text{ kg} \cdot \text{ha}^{-1}$ dose of P_2O_5 , obtaining $8.88 \text{ g plant}^{-1}$ (Figure 3A).






Figure 3. Fresh root mass (FRM) (A), dry root mass (DRM) (B), fresh aerial mass (FAM) (C) and dry aerial mass (DAM) (D) of young plants of pumpkins in function of dosages and sources of phosphorus.

The high decrease of fresh root mass can be related toxicity by phosphorus. According to [27], elevated concentrations of phosphorus can reduce photosynthesis due to the excessive exportation of triose-P from mitochondria to the cytosol, which prejudice the Rubisco regeneration and, therefore, the fixation of CO₂ in the photosynthetic process. Thus, phosphorus plays an important role in plant nutrition, as it performs metabolic functions such as the synthesis of proteins and nucleic acids [28]. Ideal dosage of phosphorus favors development of the root system, absorption of water and nutrients, increasing the quality and yield of harvested fruits.

On the other hand, the dry root mass (DRM) did not show statistical difference in function of P₂O₅ levels (Figure 3B). In relation to fresh aerial mass (FAM), all sources showed adjustment quadratic in function of different dosages, but the higher response was the TSP at 67.87 kg.ha⁻¹ dosage of P₂O₅. The second source with better performance was the SS, at 86.68 kg.ha⁻¹ dosage of P₂O₅ and the ARAD demonstrated the worst result.

As for the dry aerial mass (DAM), it is visible that all the sources were fitted to the quadratic regression model, where again the best of them was the SS with 71.10 kg.ha⁻¹ of P₂O₅, followed by TSP with 56.33 kg.ha⁻¹ of P₂O₅ and finally the ARAD with 55 kg.ha⁻¹ of P₂O₅ (Figure 3D).

However, for DAM the best performance was obtained by the SS source, this source has the advantage of containing sulfur in its composition which contributes directly to the plant growth. Sulfur (S) is absorbed by plants in the form of sulfate anionic, presents in organic matter, and a small proportion in the atmosphere, in the form of sulfuric gas. When there is deficiency of this secondary macronutrient, the protein synthesis is inhibited because the S is a participant of two essential amino acids (Cystine and Methionine), as a consequence the plants have a lower content of chlorophyll and less developed roots [29]. Many studies has showed the higher efficiency with TSP than natural phosphate, where the crops have more dry biomass production [30-32].

According to (Table 3), there was no significant difference between the sources of phosphorus for stem diameter (SD), fresh root mass (FRM) and dry root mass (DRM). In relation to NF, the most soluble sources (SS and ST) were better compared to ARAD.

Table 3. Tukey test for number of leaves (NL), stem diameter (SD), fresh root mass (FRM) and dry root mass (DRM) of young plants of pumpkins in function of phosphate sources

Sources of Phosphorus	NL	SD	FRM	DRM
Simple Superphosphate (SS)	6,40a	0,478a	7,03a	0,80a
Triple Superphosphate (TSP)	6,00a	0,468a	7,57a	0,91a
Natural Phosphate of Arad (ARAD)	4,66 b	0,460a	5,52a	0,94a

** Average followed by distinct letters differs in the column (lowercase) by the Tukey test at 5% probability.*

[26] found a similar result when testing the natural reactive phosphate doses of ARAD and Triple superphosphate, and there was no significant difference for the stem diameter in sugarcane. Kano et al. [33] observed a linear increase in the number of leaves in lettuce in function of TSP dosages (0, 200, 400, 600 and 800 kg ha⁻¹ of P₂O₅). In general, the

fertilization with the TSP source provided a greater number of leaves, corroborating with the results obtained by [34] and [35] in lettuce.

4. CONCLUSION

Young plants of pumpkins were significantly influenced by phosphate fertilization. As the superphosphate and triple superphosphate sources provided better development and accumulation of dry matter. The natural reactive phosphate of ARAD was lower in all analyzed variables.

REFERENCES

1. Araújo HS, from Quadros BR, Cardoso I, Ismael A, Corrêa CV. Potassium doses covered in pumpkin culture. *Tropical Agriculture Research*, 2012, 469-475.
2. Yavuz D, Yavuz, N, Seymen M, Türkmen Ö. Evapotranspiration, crop coefficient and seed yield of drip irrigated pumpkin under semi-arid conditions. *Scientia Horticulturae*, 2015, 197, 33-40. <http://dx.doi.org/10.1016/j.scienta.2015.11.010>.
3. Branches SRR, Lima NRS, Anjos JL, Carvalho HWL, Oliveira IR, Sobral LF, Curado FF. Technical aspects of pumpkin cultivation in the Northeast region of Brazil. Aracaju: Embrapa Coastal Trails, 2010. 36 p.
4. FAO. Agency of the United Nations for Food and Agriculture. FAOSTAT, 2017. Available at: <<http://www.fao.org/faostat/en/#data/QC>>. Accessed on: January 27, 2019.
5. IBGE. Brazilian Institute of Geography and Statistics. IBGE Automatic Recovery System (SIDRA). Agricultural Census 2006, 2016. Available at:

<https://sidra.ibge.gov.br/pesquisa/censo-agropecuario/censo-agropecuario_2006_segundaapuracao>. Accessed on: January 27, 2019.

6. Filgueira FAR. *New Olericultura Manual: Modern agro-technology in the production and commercialization of vegetables*. 3. ed. Viçosa-MG: UFV, 2012. 421 p.
7. Prates FBS, Lucas CDSG, Sampaio RA, Junior DDSB, Fernandes LA, and June GRZ. Growth of jatropha seedlings in response to fertilization with simple superphosphate and rock dust. *Revista Agrônômica*, 2012; 43 (2): 207-213.
8. Taiz L, zeiger E. *Plant physiology*. 5. ed. Porto Alegre, Artmed, 2013. 918p.
9. Oliveira RAD, Comin JJ, Tiecher T, Piccin R, Somavilla LM, Loss A, and Brunetto G. Release of Phosphorus Forms from Cover Crop Residues in Agroecological No-Till Onion Production. *Revista Brasileira de Ciência do Solo*, 2017; 41. <http://dx.doi.org/10.1590/18069657rbc20160272>.
10. Novais RF, Smyth TH, Nunes FN. Phosphor. In: Novais et al. (Eds). *Soil fertility*. Brazilian Society of Soil Science. Viçosa-MG, 2007. P471-552.
11. Mendes FF. Genetic control of phosphorus efficiency in tropical maize. Lavras: UFLA, 2012. 134p.
12. Nunes JAS, Bonfim-Silva EM, Moreira JCF. Production of radish subjected to phosphate fertilization. *Journal of the University Center of Patos de Minas*. 2014; 5: 33-44.
13. Alves L, Silva Filho GN. 2009. Production of lettuce seedlings (*Lactuca sativa* L.) in presence of different phosphate sources and phosphate solubilizing microorganisms. *Semina*, 30: 557-562.
14. Avalhães CC, Mello Prado R, Gondim ARO, Alves AU, Correia MAR. Yield and growth of beet due to fertilization with phosphorus. *Scientia Agraria*, 2009; 10 (1), 075-080.
15. Grant CA, Flaten DN, Tomasiewicz DJ, Sheppard SC. The importance of phosphorus in the initial development of the plant. *Agronomic Information*, Piracicaba, 2001; 95.

16. Resende AV, Furtini Neto AE, Alves VMC, Muniz JA, Curi N, Faquin V. et al. Sources and methods of phosphorus application to maize in cultivated soil of the Cerrado region. R. Bras. Ci. Solo, 2006.30: 453-466.
17. Faria CMB, Silva, DJ, Pinto JM, Gomes TCA. Effect of natural phosphates on melon plants cultivated in pots. R. Bras. Ci. Solo, 30: 1083-1091, 2006.
18. BRAZILIAN AGRICULTURAL RESEARCH COMPANY - EMBRAPA. National Soil Research Center. Brazilian system of soil classification. 3.ed. Brasília, 2013. 353p.
19. Cravo MS, Viegas IJM, Brazil EC. Recommendations of Fertilization and Liming for the State of Pará 1st Edition. Belém: Embrapa Amazônia Oriental, 2010. 262p.
20. Ferreira DF. Sisvar: a computer statistical analysis system. Science and agro-technology, 2011; 35 (6), 1039-1042.
21. Banzatto DA, Kronka SN. Agricultural experimentation. 4 ed. Jaboticabal: Funep, 2006. 237p.
22. Vitti GC, Wit A, Fernandes BEP. Agronomic efficiency of natural reactive thermophosphates and phosphates. In: SYMPOSIUM ON PHOSPHORUS IN BRAZILIAN AGRICULTURE. Anais ... Piracicaba: POTAFOS, 2004. p.690-694.
23. Oliveira RLL, Alves JDN, Abade MTR, Saldanha ECM, Okumura RS. Response of soybean to phosphate fertilization on yellow latosol in northeastern Para. Encyclopedia Biosphere, Goiania, 2014; 10 (18): 2682-2689.
24. Horowitz N, and Meurer EJ. Efficiency of two natural phosphates as a function of particle size. Rural science. Santa Maria. 2003; 33 (1) 41-47.
25. Bonfim-Silva EM, Silva TJA, Cabral CEA, Gonçalves JM, Pereira MTJ. Production and morphology of the java legume submitted to phosphate fertilization. Encyclopedia Biosphere, 2011; 7 (12), 1-10.
26. Relative TL, Caioni S, Lange A, Caioni C, Silva ACS, Yamashita OM, and Neto AL. Residual phosphorus in cane soca for forage production in northern Mato Grosso. Revista de Ciências Agroambientais, 2016; 14 (1).

27. MARSCHNER, H. Mineral Nutrition of higher plants. 2. Ed. London: A.P, 1995. 889 p.
28. Niu F, Zhang D, Li Z, Van Iersel MW, and Alem P. Morphological response of eucalypts seedlings to phosphorus supply through hydroponic system. *Scientia Horticulturae*, 2015;194, 295-303.: <http://dx.doi.org/10.1016/j.scienta.2015.08.029>.
29. Raij BV. Soil fertility and fertilization. Piracicaba: Ceres / Potafos, 1991, 343p.
30. Frandoloso JF, Lana MC, Fontaniva S, and Czyzca RV. Efficiency of phosphate fertilizers associated with elemental sulfur in corn crop. *Journal of Ceres*, 2010; 57 (5).
31. Fontoura SMV, Vieira RCB, Bayer C, Ernani PR, and Moraes RPD. Technical efficiency of phosphate fertilizers in Oxisol under no-tillage. *Brazilian journal of soil science*. Campinas. 2010; 34 (6): 1907-1914.
32. Silva EFL, de Araújo ASF, dos Santos VB, Nunes LAPL, Carneiro RFV. Biological fixation of N₂ in cowpea under different doses and sources of soluble phosphorus. *Bioscience Journal*, 2010.26 (3).
33. Kano C, Cardoso All, Villas Boas RL. Nutrient accumulation and response of lettuce to phosphate fertilization. *Revista Biotemas* 2012. 25: 39-47.
34. Mantovani JR, Oliveira IAC, Marques DJ, Silva AB, and Landgraf PR. Phosphorus content in the soil and production of crisp lettuce as a function of phosphate fertilization. *Semina: Agrarian Sciences*, 2014. 35 (4).
35. Effect of different doses, forms of application and sources of P in the conservation of seedless watermelon. *Horticultura Brasileira* 2016; 34: 529-536