

1  
2  
3  
4  
5  
6  
7  
8  
9

**Original Research Article**  
**Physiological quality in *Leucaena leucocephala***  
**seeds conditioned with potassium nitrate**  
**submitted to saline and water stresses**

---

10  
11  
12  
13  
14

**ABSTRACT**

*Leucaena leucocephala* is a species that occurs in semiarid regions capable of developing physiological and biochemical strategies in adverse environmental situations, such as saline soils and water deficiency. Even so, there are still losses in germination and initial development of plants mainly due effects the abiotic stresses, with this we have looked for alternatives that diminish the impact on the seeds with the use of potassium nitrate. The objective of this work was to evaluate the physiological quality in *L. leucocephala* seeds conditioned in potassium nitrate solution and submitted to saline and water stresses. After scarification, the unconditioned leucine seeds were separated and identified, and the conditioners were immersed in the 1.0 Mmol/L solution of potassium nitrate, 24 hours. The treatments were NaCl: 0; 30; 60; 90; 120 and 150 Mmol/L and the water potential: 0,0; -0.2; -0.4; -0.6, and -0.8 MPa. The seeds were distributed over two leaves of Gemitest<sup>®</sup>, moistened with distilled water 2.5 times the dry weight and incubated in B.O.D. at 25 °C for 10 days. Then, normal seedlings, germination speed index, seedling length and seedling mass were measured. Eleven treatments with four replicates of 50 seeds were used. The design was completely randomized, and the statistical analyzes were done in SAS. The use of potassium nitrate (KNO<sub>3</sub>) contributed to the maintenance of the physiological quality of *L. leucocephala* seeds under conditions of saline stress and water restriction.

15  
16  
17  
18  
19

Keywords: *Leucena*; water potencial; salinity

20  
21  
22  
23  
24  
25

**1. INTRODUCTION**

The *Leucaena leucocephala* (Lam.) Belongs to the family Fabaceae (Subfamily Mimosidae), is fast growing, is nitrogen fixer and has been highlighted as a promising alternative for the recovery vegetation cover and rehabilitation of degraded areas, since it has symbiosis with nitrogen-fixing bacteria, improving soil fertility [1].

Its use has met a wide variety of demands, including reforestation disturbed areas for erosion control, animal feed, green manuring, fence posts, poles and cellulose. **Planting/ Sowing** can be done either by seedlings or directly on the site [2], however, sowing saline soils in with a low frequency of rainfall can restrict soil moisture, resulting in slow and irregular emergence, with direct reflexes in the development of seedling [3].

26  
27  
28

The effects of salinity on germination can be noticed both by interference of salts in the cellular metabolism and by reduction the osmotic potential of seed, causing water stress and making water absorption difficult [4]. Salinity may also cause injury, such as chlorophyll

29 degradation and changes in protein metabolism, important amino acid levels linked to the  
30 germination process, and seedling initial development [5].

31 The presence of water is essential, and its scarcity is one of the adversities encountered by  
32 species of plants that develop in saline or dry environments, since availability of water to the  
33 seed, as well the speed of absorption, is directly influenced of the water potential difference  
34 between seed and soil [6]. Therefore, seeds of species the Fabaceae family even in arid and  
35 semiarid regions develop physiological and biochemical strategies capable of developing in  
36 adverse environmental situations, such as, **for example**, in soils with high salt concentration  
37 or affected by water deficiency [7,8].

38  
39 Due to the severity of losses caused by the effects of saline and water stress on germination  
40 and initial plant development, alternatives and treatments have been sought to reduce their  
41 impact on seeds. Among the substances with the potential to induce a resistance to these  
42 types of stress in the seeds, nitric oxide (NO), inorganic free radical and an extremely  
43 versatile biological messenger can be supplied exogenously the seeds through their  
44 conditioning in potassium nitrate solutions [9].

45 Research has demonstrated the involvement of nitric oxide in hormonal signaling [10] and in  
46 numerous physiological plant processes, including: protective function against oxidative  
47 stress [11]. In seeds, nitric oxide stimulates germination, both under normal conditions and  
48 under stress, besides favoring seed dormancy of some species [12] and promoting the  
49 elongation and formation of adventitious roots [13].

50 The objective of this work was to evaluate physiological quality in *Leucaena leucocephala*  
51 seeds conditioned in potassium nitrate solution and submitted to saline and water stresses.

52

## 53 **2. MATERIAL AND METHODS**

54

55 Seeds of *L. leucocephala* were collected from mature fruits with brown staining from ten  
56 matrices, spaced at a distance of 6 meters, located in the vicinity of the municipality of  
57 Macaíba - RN (Latitude: -5.86677376, Longitude: -35.33632815) in the period from July to  
58 August 2017. The region presents Aw tropical rainy climate, with an average temperature of  
59 26 °C, relative air humidity of 74 % and precipitation of 1,134 mm annually, according to  
60 Köppen and Geiger [14].

61 After collection, the seeds were manually removed from the pods, selected and mechanically  
62 scarified, using N°. 100 sandpaper, on side opposite the thread, only to break the tegument.  
63 After scarification, the water content of seeds was determined by greenhouse method at 105  
64 ± 3 °C for 24 hours, as recommended by Brazil [15]. And the results were expressed in  
65 average percentage of humidity (wet basis).

66 Then, the unconditioned seeds were separated and deposited in identified glass containers.  
67 The conditioned seeds were immersed in glass containers containing the 1.0 Mmol/L  
68 solution of potassium nitrate for 24 hours. To simulate the stress conditions, following  
69 concentrations of NaCl: 0 were used; 30; 60; 90; 120 and 150 Mmol/L [16], and also the  
70 water potentials: 0; -0.2; -0.4; -0.6 and -0.8 MPa, calculated according to the equation  
71 proposed by Michel and Kaufmann [17]. After conditioning and under conditions of saline  
72 and water stress, the seeds had their physiological potential evaluated through the tests  
73 described below.

74

75 For germination, four replicates of 50 seeds were used for each treatment, distributed on two  
76 sheets of Germitest® type paper towels and covered with a third sheet of the same paper,

77 then rolled into a roller. The substrate was moistened with distilled water (AD) in an amount  
78 2.5 times the dry substrate weight, according to Brazil [15].

79 Subsequently, paper rolls with seeds that were not conditioned and conditioned with  
80 potassium nitrate were subjected to saline stress and water stress separately and incubated  
81 in a type germination chamber (Biochemical Oxygen Demand) at 25 °C for 10 days, when  
82 final seedlings were counted. Results were expressed as percentage of germination.

83 The germination speed was carried out in conjunction with the germination test, with daily  
84 counts of seeds that emitted a primary root of at least 2 mm from the first to the tenth day  
85 after sowing. Determined from the germination velocity index (GVI), calculated according to  
86 the formula proposed by Maguire [18].

87  
88 The dry mass of seedlings was performed with total of ten normal seedlings per  
89 experimental unit, randomly taken at the end of germination test, dried for 24 hours in a  
90 greenhouse at 105 ± 2 °C, weighed in precision scale and the results expressed in grams of  
91 dry mass of seedlings.

92 The experiment was carried out in a completely randomized design with four replicates of 50  
93 seeds, using the factorial arrangement scheme: 2 (unconditioned and conditioned with  
94 KNO<sub>3</sub>) x 6 (NaCl concentrations), for salt stress and 2 conditioned and conditioned with  
95 KNO<sub>3</sub>) x 5 (osmotic potentials), for water stress. The results were submitted to analysis of  
96 variance and, when there was significance, by the F test, submitted to regression analysis.  
97 Statistical analyzes were performed using SAS<sup>®</sup> software [19].

98

### 99 3. RESULTS AND DISCUSSION

100

101 According to the analysis of variance, when applying the F test, statistical differences were  
102 verified for the variables in which *L. leucocephala* seeds were submitted, both in relation to  
103 the conditioning and concentrations in the saline stress (NaCl) and different potentials  
104 osmotic (PEG). Significance was also observed for the interaction between the factors,  
105 observing different behaviors for physiological analyzes, except for the germination variable,  
106 when subjected to saline stress conditioning and interaction, for which there was no  
107 significant effect on the evaluated treatments. When the germination of seeds submitted to  
108 water stress was evaluated, no difference was found between the means of the evaluated  
109 treatments (Table 1).

110

111 **Table 1. Summary of variance analysis for the variables germination (G), germination**  
112 **velocity index (GVI), seedling length (SL) and seedling dry mass (SDM) from**  
113 **unconditioned seeds and conditioned with potassium nitrate subjected to saline and**  
114 **water stresses.**

F values – Saline stress				
Source of variation	G (%)	GVI	SL (cm)	SDM (g)
Conditioning (C)	175,91**	334,67**	387,11**	0,16**
Saline concentrations (S)	21,10 <sup>ns</sup>	34,08**	4,01**	0,08**
(C) x (S)	74,83 <sup>ns</sup>	13,73**	25,97**	0,06**
CV (%)	7,82	14,83	11,83	12,15


F values – Hydrical stress				
Source of variation	G (%)	GVI	SL (cm)	SDM (g)
Conditioning (C)	22,94 <sup>ns</sup>	14,88**	14,21**	0,0016**
Osmotic potentials (O)	26,56 <sup>ns</sup>	51,37**	9,08**	0,0012**
(C) x (O)	28,02 <sup>ns</sup>	55,43**	120,70**	0,0010**
CV (%)	7,53	3,90	1,94	7,45

115 <sup>ns</sup> not significant, \*\* significant at 1% by the F test.


116

117 The results of the germination test on conditioned or non-saline seeds (Figure 1 A) show  
 118 that, despite salinity, there was a positive effect of conditioning on this variable. Although the  
 119 seeds did not present significant variation with the progressive increase of the salinity of the  
 120 substrate, it was observed effect of conditioning on the germination of seeds until the  
 121 concentration of 90 Mmol/L of NaCl, which presented higher germination than the  
 122 unconditioned seeds. In general, the seeds presented high germination, even when not  
 123 conditioned. A similar result was verified by Amaro [20], which evaluated *Copaifera*  
 124 *langsdorffii* Desf. (Fabaceae) seeds on a substrate moistened with NaCl solution, with a high  
 125 percentage of germination, around 95 %. The *Leucaena leucocephala* (Lam.) De Wit seeds  
 126 Cavalcante and Perez [21] added sodium chloride (NaCl) to substrate and observed high  
 127 germination percentages of the seeds, which reaffirmed ability of the seeds this species to  
 128 present a high germinative performance even under saline stress conditions.

129



130



131

132 **Figure 1. (A) Germination, (B) Germination velocity index, (C) Seedling length, and (D)**  
 133 **Dry mass of *L. leucocephala* seedlings from not conditioned seeds conditioned with**  
 134 **potassium nitrate to saline stress).**

135 Thus, *L. leucocephala* seeds, as well as those of several other species of the Fabaceae  
 136 family, show resistance or some kind of defense system, capable of synthesizing and  
 137 accumulating different compounds of small molecular mass such as sugar alcohols, proline  
 138 and glycine betaine, which are referred to as compatible osmopolymers, osmoprotectants or  
 139 solutes. And these compounds have an exact function in the plants, being able to be related  
 140 protection of the plants to abiotic stresses, functioning as a tool for the cellular osmotic  
 141 adjustment, and protection against oxidative damages under adverse conditions, as in  
 142 drought and salinity tolerance [22].

143 However, it was verified that potassium nitrate (KNO<sub>3</sub>) did not promote seed germination  
 144 improvement, since these already present high viability even under conditions of salt stress.

145 It was observed that the rate of germination increased as the substrate salt concentrations  
 146 increased, evidencing that potassium nitrate (KNO<sub>3</sub>) conditioning had a positive influence the  
 147 vigor of *L. leucocephala* seeds (Figure 1 B). However, the not conditioned seeds presented  
 148 a tendency to reduce the germination speed. The increase in the germination speed for  
 149 conditioned seeds (Figure 1 B) can be explained by the presence of compounds that act on  
 150 potassium nitrate (KNO<sub>3</sub>) in the seeds, since they can stimulate and influence the  
 151 physiological quality of the seeds, its vigor. This is because the mechanism of action of  
 152 potassium nitrate acts on the reception of electrons, reducing the nitrite form inside the  
 153 seeds, reoxidizing the NADPH and increasing the availability of NADP for the reduction of  
 154 the dehydrogenases of the cycle of pentose phosphate, helping the overcoming of seed  
 155 dormancy [3]. In addition to being an excellent growth promoter of seedlings, potassium  
 156 nitrate can also act to maintain balance in plant cells, promoting respiration and metabolism  
 157 of carbohydrates [23].

158

159 Regarding the unconditioned seeds, lower performance was observed in all salinity  
 160 concentrations when the germination rate, length and dry mass of seedlings were evaluated.  
 161 As for the variable germination, the vigor of the unconditioned seeds was more affected by  
 162 the salinity. Evaluating the physiological quality of seeds of *Crambe abyssinica* Hochst. ex R.


163 E. Fr., Nunes et al. [24] found that the increase in saline concentration of the substrate also  
 164 reduced the rate of germination. This decrease in germination speed was also observed by  
 165 Sousa Filho [25] in *Erythrina mulungu* Mart ex Benth. (Fabaceae) seeds under saline stress.  
 166 These results confirm the negative effect of soil salinity on the vigor of seeds of different  
 167 species, reducing the emergence speed and the growth of the seedlings, generating  
 168 damages to agricultural production.


169 Salinity probably interferes with the physiological quality of seeds as well, in terms of  
 170 reservoir energy expenditure to absorb water and subsequently not reserving the reservoir  
 171 for other processes, inducing changes in the activities of catalase, polyphenoloxidase and  
 172 peroxidase enzymes [26]. In studies carried out by Marques et. [27], the effect of saline  
 173 stress on emergence and establishment of seedlings seems to have been responsible for  
 174 the inhibition of cotyledonary reserve depletion and embryonic axis growth, since in the  
 175 analyzes the seedlings presented excessive accumulation of  $\text{Na}^+$  and  $\text{Cl}^-$  ions.

176 The excess  $\text{Na}^+$  and  $\text{Cl}^-$  ions seem to play a role in reducing the physiological quality of  
 177 seeds, since they tend to cause a decrease in protoplasmic intumescence (the ions in  
 178 solution initially cause a decrease in intumescence, and only after absorption and  
 179 accumulation in the vacuoles and apoplast is that the rate of water absorption is normalized),  
 180 affecting the enzymatic activity, resulting, mainly, in the inadequate production of energy by  
 181 disturbances in the respiratory chain [28]. In addition, it is necessary to emphasize its toxic  
 182 effect, resulting from the concentration of ions in the protoplasm. As the saline concentration  
 183 increases in the soil solution, there is an increase in the osmotic pressure and, therefore, the  
 184 plant does not absorb the soil water, causing physiological and morphological disturbances  
 185 that hinder the survival of the plant in this stress condition [29].

187 When seeds conditioned or not subjected to water stress, no change in seed germination  
 188 was observed along the reduction of water availability and no difference between the two  
 189 treatments in any of the potentials tested.

190 However, as the water potential became more negative, there was a decrease in the rate of  
 191 germination of *L. leucocephala* seeds conditioned and not conditioned with potassium nitrate  
 192 ( $\text{KNO}_3$ ), and conditioned seeds presented higher results in relation to unconditioned seeds  
 193 (Figure 2 A). Similar behavior was observed by Lopes et al. [30] when studying the  
 194 physiological quality of seeds of *Galliesia integrifolia* (Spreng.) Harms, verified that seeds  
 195 treated with  $\text{KNO}_3$ , at concentrations of  $1 \text{ mg L}^{-1}$  and  $10 \text{ mg L}^{-1}$ , presented higher averages  
 196 for the germination speed index, when compared the not conditioned seeds.  
 197





199  
200  
201  
202  
203  
204

**Figure 2. Germination velocity index (A), seedling length (B) and dry mass (C) of *L. leucocephala* from seeds not conditioned and conditioned with potassium nitrate (and submitted to water stress).**

205  
206  
207  
208  
209  
210

When evaluating the viability of *Apuleia leiocarpa* (Vogel) J.F.Macbr. (Fabaceae) in different water potentials, Spadeto et al. [31] also observed that the more negative the potentials, the lower germination speed averages were -0.4 and -0.8 MPa obtained 3.80 and 3.74, respectively. When Felix et al. [34] evaluated the effects of water stress, verified that the germination speed in *L. leucocephala* seeds were reduced when submitted to the most negative water potentials, -0.3 and -0.6 MPa, obtaining a mean of 8.0 and 4.0, respectively.

211  
212  
213  
214  
215  
216

Therefore, the results of these researchers corroborate with those obtained in this research, indicating the negative effect of reducing the availability of water in substrate on the germination of seeds this species. This fact is probably due to decrease in water availability, which limits imbibition and reduction of seed water intake, reducing the metabolic activities expected during germination phases, slowing down the process of mobilization and utilization reserves, leading to reduction of the rate germination [5].

217  
218  
219  
220  
221

It is observed that *L. leucocephala* seeds presented high rate of germination in the conditions in which the lowest water restrictions were observed. According to Bewley et al. [5] Seeds such as *L. leucocephala* are tolerant to water stress due to their natural occurrence in semiarid regions, as they acquire biochemical and physiological strategies that can be attributed to membrane repair even in situations of low water potential.

222  
223  
224  
225  
226  
227

Conditioning with potassium nitrate ( $\text{KNO}_3$ ) has influenced the reversal of harmful changes in cell membranes, caused by inactivation of enzymes and inhibition of protein synthesis [29]. This shows that even under conditions of stress of abiotic origin, *L. leucocephala* seeds had positive effects on the physiological responses involved in the whole germination process, regulating percentage of germination and rate germination, as well as the growth, development and establishment of seedlings.

228  
229  
230  
231  
232  
233

There was a tendency to increase the length and dry mass of *L. leucocephala* seedlings as the water restriction was increased, regardless of the potassium nitrate ( $\text{KNO}_3$ ) or not. However, the seeds presented different results for length and dry mass of seedlings when compared to conditioned and unconditioned seeds. With the reduction of the water potential, the length of the seedlings resulting from conditioned seeds presented results with a tendency to increase the difference in relation to the unconditioned seeds, indicating an

234 increase in the size conditioned seedlings in relation the untreated seedlings under  
235 conditions of lack of water (Figure 2 B). On the other hand, the dry matter of seedlings  
236 showed a decrease in the water potential (Figure 2 C), in relation to the uncon solved seeds.

237

238 The ability to develop roots and aerial parts, even under conditions of stress and water  
239 scarcity, varies widely between species and may be related to the hypothesis of adaptation,  
240 due to the proliferation and the increase of the roots increase the capacity of water  
241 absorption in a limit situation, or may be related to the growth rate of a given plant [32].

242 Many proteins that are induced in the early stages of water stress are also involved in root  
243 morphogenesis and carbon/nitrogen metabolism, which may contribute to the nullification of  
244 stress by increasing root growth [33]. In the case of *L. leucocephala*, the literature reports  
245 that it is a plant adapted to inhospitable conditions and such rusticity can favor the tolerance  
246 of the seeds under different conditions.

247

#### 248 **4. CONCLUSION**

249

250 The use of potassium nitrate (KNO<sub>3</sub>) contributed to the maintenance of the physiological  
251 quality of *L. leucocephala* seeds under conditions of saline stress and water restriction.

252

#### 253 **COMPETING INTERESTS**

254

255 Authors have declared that no competing interests exist.

256

#### 257 **AUTHORS' CONTRIBUTIONS**

258

259 This work was carried out in collaboration with all authors.

260

#### 261 **REFERENCES**

262

- 263 1. Nogueira NO, Oliveira OM de, Bernardes CA da SM, Bernardes C de O. Utilização  
264 de leguminosas para recuperação de áreas degradadas. Enciclopédia Biosfera.  
265 2012;8(14):2121.
- 266 2. Cardoso EA, Alves AU, Cavalcante ÍHL, Farias SGG, Santiago FEM. Métodos para  
267 superação de dormência em sementes de leucena. Revista Ciências Agrárias.  
268 2012;55(3):220–4. Portuguese Available: <http://dx.doi.org/10.4322/rca.2012.062>
- 269 3. Mariano LG, Somavilla A, Silveira AG da, Salamoni AT. Análise de superação de  
270 dormência de sementes de *Leucaena leucocephala* e desenvolvimento inicial de  
271 plântulas. Electronic Journal of Management, Education and Environmental  
272 Technology. 2016;20(1):398–404. Portuguese. Available:  
273 <http://dx.doi.org/10.5902/2236117019719>
- 274 4. Begum MAJ, Selvaraju P, Venudevan B. Saline stress on seed germination.  
275 Scientific Research and Essays. 2013;8(30):1420–3. Available:  
276 <http://dx.doi.org/10.5897/SRE2013.5600>
- 277 5. Bewley JD, Bradford K, Hilhorst H, Nonogaki H. Seeds: physiology of development,  
278 germination and dormancy. 2013.
- 279 6. Cavallaro V, Barbera AC, Maucieri C, Gimma G, Scalisi C, Patanè C. Evaluation of  
280 variability to drought and saline stress through the germination of different ecotypes  
281 of carob (*Ceratonia siliqua* L.) using a hydrotime model. Ecological Engineering.  
282 2016;95:557–66. Available: <http://dx.doi.org/10.1016/j.ecoleng.2016.06.040>
- 283 7. Elias SG, Copeland LO, McDonald MB, Baalbaki RZ. Seed testing: principles and  
284 practices. East Lansing: Michigan State University Press.; 2012. 368 p. Available:



- 285 <http://dx.doi.org/1611860393,9781611860399>
- 286 8. Guedes RS, Alves EU, Viana JS, Gonçalves EP, Lima CR de, Santos SDRN dos.  
287 Germinação e vigor de sementes de *Apeiba tibourbou* submetidas ao estresse  
288 hídrico e diferentes temperaturas. *Ciência Florestal*. 2013;23(1). Portuguese.  
289 Available: <http://dx.doi.org/10.5902/198050988438>
- 290 9. Martins CC, Pereira MRR, Lopes MTG. Germinação de sementes de eucalipto sob  
291 estresse hídrico e salino. *Bioscience Journal*. 2014;30(3):318–29.
- 292 10. Matakiaadis T, Alboresi A, Jikumaru Y, Tatematsu K, Pichon O, Renou J-P, et al. The  
293 Arabidopsis Abscisic Acid Catabolic Gene CYP707A2 Plays a Key Role in Nitrate  
294 Control of Seed Dormancy. *Plant Physiology*. 2008;149(2):949–60. Available:  
295 <http://dx.doi.org/10.1104/pp.108.126938>
- 296 11. Zheng C, Jiang D, Liu F, Dai T, Liu W, Jing Q, et al. Exogenous nitric oxide improves  
297 seed germination in wheat against mitochondrial oxidative damage induced by high  
298 salinity. *Environmental and Experimental Botany*. 2009;67(1):222–7. Available:  
299 <http://dx.doi.org/10.1016/j.envexpbot.2009.05.002>
- 300 13. Pereira BLC, De Lima E Borges EE, Oliveira AC, Leite HG, De Carvalho Gonçalves  
301 JF. Influência do óxido nítrico na germinação de sementes de *Plathymenia reticulata*  
302 Benth com baixo vigor. *Scientia Forestalis*. 2010;(88):629–36. Portuguese.
- 303 14. Köppen W, Geiger R. *Klimate der Erde*. Gotha: Verlag Justus Perthes. 1928.
- 304 15. Brasil. Ministério da Agricultura, Pecuária e Abastecimento. Regras para análise de  
305 sementes. Brasília, DF: MAPA/ACS; 2009. 399 p. Portuguese.
- 306 16. Betoni R, Scalon S de PQ, Mussury RM. Salinidade e temperatura na germinação e  
307 vigor de sementes de mutambo (*Guazuma Ulmifolia* Lam.) (Sterculiaceae). *Revista*  
308 *Árvore*. 2011;35:605–16. Portuguese. Available: [http://dx.doi.org/10.1590/S0100-](http://dx.doi.org/10.1590/S0100-67622011000400004)  
309 [67622011000400004](http://dx.doi.org/10.1590/S0100-67622011000400004)
- 310 17. Michel BE, Kaufmann MR. The osmotic potential of polyethylene glycol 6000. *Plant*  
311 *Physiology*. 1973;51(5):914–6. Available: <http://dx.doi.org/10.1104/PP.51.5.914>
- 312 18. Maguire JD. Speed of germination-aid selection and evaluation for seedling  
313 emergence and vigor. *Crop Science*. 1962;2:176–7. Available:  
314 <http://dx.doi.org/http://dx.doi.org/10.2135/cropsci1962.0011183X000200020033x>
- 315 19. SAS Institute (2011) SAS/STAT Versão 9.3: User's Guide. SAS Institute Inc., 8621.
- 316 20. Amaro MS. Germinação de sementes e mobilização de reservas em plantas de  
317 copaibasobr estresse hídrico e salino. Tese (Doutorado em Fitotecnia). Universidade  
318 Federal do Ceará; 2012. Portuguese.
- 319 21. Cavalcante A de MB, Perez SCJG de A. Efeitos dos estresses hídrico e salino sobre  
320 a germinação de sementes de *Leucaena leucocephala* (Lam.) de Wit. *Pesquisa*  
321 *Agropecuária Brasileira*. 1995;30(2):281–9. Portuguese.
- 322 22. Hurtado Salazar A. Tolerância ao estresse abiótico (salinidade e seca) e influência  
323 de porta-enxertos na qualidade de frutos de *Passiflora* spp. Tese (Doutorado em  
324 Fitotecnia). Universidade Federal de Viçosa; 2016. Portuguese.
- 325 23. Brito SF, Bezerra AME, Pereira DS. Efeito da Temperatura e do KNO<sub>3</sub> na  
326 germinação de *Acnistus arborescens* (Solanaceae). *Floresta e Ambiente*.  
327 2016;22(3):406–12. Portuguese.
- 328 24. Nunes A da S, Souza LCF de, Scalon S de PQ, Pagnoncelli J. Nitrato de potássio e  
329 retirada do pericarpo na germinação e na avaliação do vigor de sementes de  
330 crambe. *Semina: Ciências Agrárias*. 2015;36(3):1775–82. Portuguese. Available:  
331 <http://dx.doi.org/10.5433/1679-0359.2015v36n3Supl1p1775>
- 332 25. Sousa Filho PH de, Pereira DP, Marques JB, Moreira ÉFA, Carvalho M, NAVES MF.  
333 Efeito do estresse salino na germinação de sementes de *Erythrina mulungu* Mart. ex  
334 Benth. *Anais do I Seminário de Pesquisa e Inovação Tecnológica*. 2017;1(1).  
335 Portuguese.
- 336 26. Maia JM, Ferreira-Silva SL, Voigt EL, Macêdo CEC de, Ponte LFA, Silveira JAG.  
337 Atividade de enzimas antioxidantes e inibição do crescimento radicular de feijão

- 338 caupi sob diferentes níveis de salinidade. Acta Botânica Brasilica. Portuguese.  
339 2012;26(2):342–9. Available: <http://dx.doi.org/10.1590/S0102-33062012000200010>
- 340 27. Marques EC, Freitas VS, Bezerra MA, Prisco JT, Gomes-Filho E. Efeitos do estresse  
341 salino na germinação, emergência e estabelecimento da plântula de cajueiro anão  
342 precoce. Revista Ciência Agronômica. 2011;42(4):993–9. Portuguese. Available:  
343 <http://dx.doi.org/10.1590/S1806-66902011000400023>
- 344 28. Larcher W. Ecofisiologia vegetal. Editora Rima. 2000. Portuguese.
- 345 29. Taiz L, Zeiger E. Fisiologia Vegetal. 3rd ed. Editora Artmed. 2013. 918 p.
- 346 30. Lopes JC, Freitas AR de, Beltrame RA, Venancio LP, Manhone PR, Silva FRN e.  
347 Germinação e vigor de sementes de pau d’alho sob estresse salino. Pesquisa  
348 Florestal Brasileira. 2015;35(82):169. Portuguese. Available:  
349 <http://dx.doi.org/10.4336/2015.pfb.35.82.631>
- 350 31. Spadeto C, Mengarda LHG, Paulucio MC, Lopes JC, Matheus MT. Embebição,  
351 osmocondicionamento e viabilidade de sementes de *Apuleia leiocarpa* (VOGEL.) J.  
352 F. MACBR. Ciência Florestal. 2018;28(1):80. Portuguese. Available:  
353 <http://dx.doi.org/10.5902/1980509831582>
- 354 32. Forti, VA, Cicero, SM, Pinto, TLF Efeitos de potenciais hídricos do substrato e teores  
355 de água das sementes na germinação de feijão. Revista Brasileira de Sementes.  
356 2009;31(2):63-70. Portuguese. Available: <http://dx.doi.org/10.1590/S0101-31222009000200007>
- 357
- 358 33. Masetto TE, Quadros J de B, Ribeiro DM, Rezende RKS, Scalon S de PQ. Potencial  
359 hídrico do substrato e teor de água das sementes na germinação do crambe. Revista  
360 Brasileira de Sementes. Portuguese. 2011;33(3):511–9. Available:  
361 <http://dx.doi.org/10.1590/S0101-31222011000300014>
- 362 34. Felix FC, Araújo FS, Silva MD, Ferrari CS, Pacheco MV. Estresse hídrico e térmico  
363 na germinação de sementes de *Leucaena leucocephala* (Lam.) de Wit. Revista  
364 Brasileira de Ciências Agrárias. 2018;2(13):e5515. Portuguese. Available:  
365 <http://dx.doi.org/10.5039/agraria.v13i2a5515>