

# Geology, Palynomorphs distribution, Stratigraphy and Depositional Environments of Lewumeji and Idogun wells, Eastern Dahomey Basin Southwestern Nigeria

## Abstract

Selected composited samples from Lewumeji (0-111m) and Idogun (0- 54m) Abeokuta Group, Eastern Dahomey Basin, were subjected to detailed lithologic and palynological studies. The studies aimed at determining the lithological sequence, relative age, palynological zone and paleoenvironments of deposition. The lithological description was done using hand lens, visual examination and dilute HCl. The palynologic sample preparation went through sample digestion, flotation and mounting on glass slides in order to determine palynomorphs contents such as pollen, spore and dinoflagellates present.

The lithologies from both wells revealed brownish clay, reddish to brown colour sandstone and dark grey shale denoting marine, fluvial, brackish and lagoonal environment. A total of 31 well preserved low to moderate diverse palynomorphs were recovered from the studied area. The palynomorphs frequency percentage distribution shows that both wells has a higher frequency of land derived pollen and spores to the marine dinoflagellates; (75%, 25% and 61%, 39%) for Lewumeji and Idogun wells respectively. The microfloral assemblages include abundant *Cyathidites* sp, *Cyathidites* minor, *Tubistephanocolpites* cylindricus, *Proteacidites* sp, *Trilete* spore, *Foveotriletes* margaritae, *monocolpites* marginatus, *monoporites* annulatus, *pteris* sp, *Distaverrusporites* simplex and *Laevigatosporites* sp. The dinoflagellates recovered were characterized by the likes of *Leiosphaeridia* sp, *Senegalinium* sp, *Oligosphaeridium* sp, *paleocytodinium* sp, *Cerodinium* sp and *Subtilisphaera* sp. The wells fall within *Cyathidites* Minor zone, characterized by the diagnostic occurrence of *Cyathidites* minor, *Cyathidites* sp and *monocolpites* marginatus dated Upper Maastrichtian to Early Paleocene. Paleoenvironmental deductions based on abundance of freshwater swamps pollen and Spores, diagnostic dinoflagellates cyst and Palynomorphs marine Index (PMI) indicated a continental to brackish to shallow marine environment with minor influx of freshwater.

Keywords: Dahomey Basin, lithostratigraphy, palynology, Paleoenvironment,

Word counts: 269

## 1 Introduction

The applications of biostratigraphy in the palynological studies have become more valuable tools and universally accepted methods of evaluating the stratigraphy and source rock potential of sedimentary basins. These include the modern and fossil pollen, spores and dinoflagellates cysts. This marker species gives reliable and accurate information about past environments. When these markers are efficiently utilized, many of the hindrances encountered in paleoenvironmental synthesis can be avoided (Adegoke, 2012). The study area, Lewumeji and Idogun wells, falls within the Abeokuta group of the Eastern Dahomey basin (Fig.1, Fig. 2). The Abeokuta group is the oldest formation in the Dahomey Basin, Southwestern Nigeria, lying non-conformably on the basement (Jones and Hockey, 1964) and it is the thickest group within the basin, with an average thickness of 200m (Fayose, 1970). The basin is a pre-cratonic basin that was developed during the initiation of rifting associated with the opening of Gulf of Guinea in Early Cretaceous to

Comment [S1]: sp.

Comment [S2]: all specific names must be written in italic!!!

42 Late Jurassic (Whiteman, 1982; Kingston *et al.*, 1983). Agagu, (1985) illustrated and described  
43 the lithostratigraphy of the basin to be dominated by Monotony of Sand and shale alterations  
44 with minor proportion of Limestone and clay.


45 Several workers have carried out considerable and intensive researches to characterize and  
46 deduce the age of the sediments in the basin. (Omatsola and Adegoke, 1981; Salami, 1987;  
47 Obaje and Okosun, 2013, Adeigbe and Amodu, 2015). The stratigraphy of the Dahomey Basin  
48 has been well established by various authors, Jones and Hockey, (1964), Omatsola and Adegoke,  
49 (1981), Coker *et al.*, (1983), Biliman (1992), Enu, (1990).

50 The studied wells are situated between latitudes  $06^{\circ}30'0''\text{N}$  -  $06^{\circ}37'0''\text{N}$  and longitude  $04^{\circ}45'0''\text{E}$   
51 -  $05^{\circ}00'0''\text{E}$ . and falls within the Abeokuta group of the Eastern Dahomey Basin (Fig.1 and  
52 Fig.2)

53 The present study focuses on using palynological and lithostratigraphic data to enhance the  
54 detailed general lithological description, deductions of age, varying depositional environment as  
55 well as to interpret the Biostratigraphy (biozones) of Lewumeji (0 – 111m) and Idogun wells (0  
56 – 54m).


## 57 2 Sampling and Methods

58 The core samples used for this study were collected from the Bitumen project base Ore in Ondo  
59 state, Nigeria. The cores were sampled at every 3.0 meters interval from top to bottom of the  
60 Boreholes. A total of Four (4) composited samples from the Lewumeji well and five (5)  
61 composited sample from Idogun well were used for this study.



62

63 Figure 1: [Location Map](#) of Dahomey Basin [showing and the sStudy wells and geological scheme](#)  
 64 [of the area](#) (modified after Biliman, 1992)



65  
 66 Figure 2: Location map of the exploration wells for this Study, Eastern Dahomey basin,  
 67 Southwestern Nigeria. (Inset map illustrates the position of Dahomey basin in Nigeria)

68 **2.1 Lithologic description**

69 Detailed lithologic description was carried out on the core samples provided, by following the  
 70 standard method of describing samples by using microscope hand lens, dilute HCl and physical  
 71 examination. The description was based on their texture, Fissility, colour, and fossil content in  
 72 term of plant remains and fossil fragment.

73 **2.2 Palynological Analysis**

74 The purpose of palynological preparation is to disaggregate the fossil palynomorphs from the  
 75 rock or sediment matrix. Factors that can make samples unreliable, such as: Laboratory  
 76 contamination, assemblages mixing in nature and misplacing of samples through human error  
 77 were put into consideration during analysis for optimum retrieval of palynomorph. A standard  
 78 extraction method was used. 10g of each sample was weighed, gently crushed to avoid  
 79 deforming the palynomorphs, and poured into a well labelled plastic beaker and placed in a fume  
 80 cupboard. Each samples was digested with 10% hydrochloric acid (HCl) for about 15minutes for  
 81 carbonate removal and soaked overnight with 40% hydrofluoric acid (HF) for the removal of  
 82 silicate. From the preceding preparatory stage a drop of potassium chlorate (KClO<sub>3</sub>) was added,  
 83 which was stirred and left for about 5minutes so as to react, it is then rinse twice to remove the  
 84 KClO<sub>3</sub>. A 5 micron sieve was then used under a branson sonifier to wash out the inorganic

85 matter (mud and clay). A drop of Norland Optical Adhesive was then deposited on the slides to  
86 be used. The slides were studied under a light transmitted microscope to obtain the  
87 palynomorphs.

### 89 3.0 Results

#### 90 Interpretation and Discussion

#### 91 3.1 Lithostratigraphy

92 The nine (9) composited samples of the studied sections of Lewumeji and Idogun wells, were  
93 carefully studied based on their Lithology. Three (3) litho units were identified in the study  
94 wells. All the Three (3) units occur in Idogun well with the alternation of shale and sandstone  
95 while two (2) units occurred in Lewumji well with a little clay intercalation. The three lithounits  
96 ~~are consisting respectively of~~ sandstone, clay and shale. The description of the facie units are  
97 presented below while the litholog is shown in Fig. 3 and Fig.4.

#### 98 Lewumeji well

##### 99 Litho unit 1 (0-15m)

101 This unit is on the topmost layer. The sandstone is reddish brown at the upper part of the unit  
102 then a light brown at the base of the layer. It has a fine to medium size grains. The unit is 15m  
103 thick and was deposited in a fluvial environment. This is further confirmed by the  
104 palynological study carried out which revealed the presence of an Angiosperm pollen

105 Tubistephanocolpites *Cylindricus cylindricus*

##### 106 Litho unit 2 (15 – 111m)

107 This unit is 96m thick. It is composed of shaly, dark to grayish, fissile to non- fissile,  
108 carbonaceous shale. Also the occurrence of *Microforaminiferal* wall lining and  
109 *Laevigatosporites* sp within the interval suggested that it could have been deposited in a marine  
110 environment.

#### 111 Idogun well

##### 112 Litho unit 1 (0 -9m)

114 This units is 9m thick, it is reddish brown, non- carbonaceous clay. This litho unit portrays a  
115 mixed depositional environment in which there is strong influence of fluvial on lagoonal  
116 environment.

##### 117 Litho unit 2 (9 – 15m)

118 This interval is composed of fine to medium grain sandstone with an evidence of shelly whitish  
119 material in some horizons. It is 6m thick and reddish brown to brown in colour. The sediment  
120 was deposited in a fluvial environment.

Comment [S3]: ????

Comment [S4]: put on later

Comment [S5]: Italic

Comment [S6]: what are they?

Comment [S7]: ital

Comment [S8]: put on later

Formatted: Font: Bold

Comment [S9]: put on later

Comment [S10]: ????

121 **Litho unit 3 (15 – 24m)**

Formatted: Font: Bold

122 This unit which is about 9m thick is shally, dark grey in colour, non- fissile and could have been  
 123 deposited in a marine environment.

124 Litho unit 4 (24 – 42m)

125 This units is made up of grey coloured sandstone. Fine to medium grained. The occurrence of  
 126 monocolpites marginatus, *Tubistephanocolpites ceylindericus* also suggests deposition in a  
 127 fluvial environment.

Comment [S11]: Italic

Comment [S12]: put on later

128 **Lithounit 5 (42 -54m)**



Formatted: Font: Bold

129 This interval consists of a dark to greyish non- fissile shale. It is carbonaceous. The units is  
 130 about 9m thick and the high occurrence of dinoflagellates cysts like *Senegalinium* sp,  
 131 *Paleocytonium* sp, *Ssubtilisphaera* sp suggests position in a marine setting.

Comment [S13]: italic

Comment [S14]: it

Comment [S15]: italic



133 Figure 3: Litholog of the studied interval of  
 134 Lewumeji well, Dahomey Basin, Nigeria.

133 Figure 4: Litholog of the studied interval of  
 134 Idogun well, Dahomey Basin, Nigeria

135 **3.2 PALYNOLOGICAL STUDIES**

136 Analytical breakdown of the palynomorphs showed that the samples are well preserved with a  
 137 low to moderate occurrence and moderately diverse pollen, spores and the dinoflagellates. Some  
 138 of the palynomorphs recovered in Lewumeji well are *Tubistephanocolpites cylindricus*,  
 139 *Proteacidites* spp, *Monocolpites marginatus*, *Cyathidites* spp., *Laevigatosporites* spp.,  
 140 *Cyathidies minor*, *Leiosphaeridia* spp and marine diagnostic specie microforaminiferal wall  
 141 lining were recorded. palynomorphs recovered in Idogun well are *Monoporites annulatus*,  
 142 *Monocolpites marginatus*, *Proteacidites* spp., *Foveotritetes Margaritae*, *Mauritiidites lehmani*,  
 143 *Tubistephanocopites ceylindericus*, *Cyathidites* spp., *Laevigatosporites* spp., *Trilete* spore,

Comment [S16]: all the specific name in italic!!!!

144 | *Distaverrusporites simplex*, *Pteris* spp., *Leiosphaerida* spp., *Cerodinium* spp., *Oligosphaeridium*  
145 | spp., *Paleocystodinium* spp., *Senegalinium* spp., *Subtilisphaera* spp. and marine diagnostic specie  
146 | microforaminiferal wall lining. The marine dinoflagellates cyst makes up to 39%, while the  
147 | pollen and spores makes up to 27.77% and 33.3% respectively of the total palynomorphs in  
148 | Idogun well while the marine dinoflagellates cysts makes up about 25%, while the pollen and  
149 | spores makes up about 33.33% and 41.66% respectively of the palynomorphs in Lewumeji well.

Comment [S17]: italic

Comment [S18]: all the specific name in italic!!!!

Comment [S19]: all the specific name in italic!!!!

### 150 3.2.1 Palynological zones and Correlation

151 | The erection of biozones is dependent of the evolution, extinction and quantitative occurrence of  
152 | marker forms present in the sediments (Ola- *Bburaimo*, 2012). The palynological interpretation  
153 | of the analyzed interval was based on diagnostic marker species. For the entire section of the  
154 | Lewumeji (0 – 111m) and Idogun (0 -54m) wells, the recovered palynomorphs enabled the  
155 | delineation of one major zone which is the *Cyathidites Minor Assemblage* zone, based on the  
156 | occurrence of *Cyathidites minor*, *Cyathidites* sp and *Monocolpites marginatus* this erected zone  
157 | can also be correlated with *spinizonocolpites Bacculatus* zone of *Lawal and Moullade*, (1987).  
158 | The details of the palynological zones recognized for lewumeji and Idogun well are discussed  
159 | below and shown graphically in the palynology distribution chart (Fig. 5 to Fig. 8). The chart  
160 | shows the ages of the recovered palynomorphs and the Index palynomorphs which marked the  
161 | zones as recorded in the bioevent section of the chart. The basis of characterization of Lewumeji  
162 | and Idogun well is given below:

Comment [S20]: all specific name in italic

163 | **Zone:** *Cyathidites mMinor* Assemblage zone

164 | **Interval:** 0.00m – 111.0m; 0.00m – 54.0m

165 | **Age:?** Upper Maastrichtian – Early Paleocene

#### 166 Characteristics

167 | For the Lewumeji well, the zone is marked at the base (75.00-111.00) by the occurrence  
168 | *Cyathidites* sp, *Cyathidites minor*, *Tubistephanocolpites cylindricus*, and the acritarch  
169 | *Leiosphaeridia* Sp. The part near the base (45.00- 75.00) is characterized by the new appearance  
170 | of *Monocolpites Marginatus*, *Laevigatosporites* spp, Microforaminiferal wall lining and  
171 | continuous occurrence of *Leiosphaeridia* Spp. Close to the top of the well (15.00- 45.00) is the  
172 | new appearance of *Proteacidites* sp, continuous occurrence of *Laevigatosporites* sp and  
173 | *Cyathidites* sp. while the topmost part (0.00-15.00) is very sparse in spores and dinoflagellates  
174 | cyst but marked by the single occurrence of an angiosperm pollen which is  
175 | *Tubistephanocolpites cylindricus* (table 1). A considerable amount of palynomorphs assemblage

Comment [S21]: italic

Formatted: French (France)

Formatted: French (France)

Formatted: French (France)

Comment [S22]: all specific name in italic

Comment [S23]: italic


176 found in this well have been reported for late Maastrichtian to Paleocene sediment in the basal  
 177 part of Araromi (Salami 1984, Adeigbe and Amodu, 2015), for the Paleocene sediment of Pan  
 178 tropical area (Germeraad *et al.*, 1968), for the cretaceous sediment of upper Benue trough (Lawal  
 179 and molluade, 1986; Awad, 1994), Major forms present in the upper Maastrichtian facies are  
 180 often present in Paleocene sediments. (Ola-Buraimo, 2012; Ayinla *et al.*, 2013).

181 For Idogun well, the interval study also belong to the Cyathidites minor assemblage zone.  
 182 Dinoflagellates cyst dominate the basal part (42.00-54.00m) of the well which is an indication of  
 183 more marine influence, these include the assemblages of Senegalinium sp, Oligosphaeridium  
 184 sp, *Subtilisphaera* sp, Cerodinium sp and relative high frequency of Paleocytodinium sp. The  
 185 diagnostic marker forms present are *Cyathidites* sp, *Monoporites anulatus* and *Monocolpites*  
 186 *Marginatus*. At depths 24.00 to 42.00m there is re-occurrence of *Monocolpites Marginatus* and  
 187 new forms that are diagnostics of late maastrichtian age, emerged, they include *Mauritiidites*  
 188 *lehmani*, *Tubistephanocolpites Cylindricus*, and *Pteris* Sp. the overlying interval (15.00 – 24.00)  
 189 is characterized by occurrence of new forms *Distaverrusporites simplex* which supports the late  
 190 Cretaceous age (Durugbo and Aroyewun, 2012). The overlying interval 9.00m – 15.00m is  
 191 relatively rich in palynomorphs, it is composed of continuous occurrence of *Cyathidites* sp.  
 192 Miospores and dinocysts that appear for the first time are *Leiosphaeridia* sp, *Trilete* Spore,  
 193 *Microforaminiferal wall lining*, *Foveotriletes Margaritae*, and *Laevigatosporites* sp. The topmost  
 194 interval 0.00m -9.00m is characterized by the re occurrence of *Cyathidites* sp and new  
 195 appearance of *Proteacidites* sp as shown in table 2

Comment [S24]: italic

Comment [S25]: all specific name in italic

Comment [S26]: all specific name in italic



196  
 197 Fig 5: Palynomorphs zones recognised in  
 198 Lewumeji well.






Fig 6: Palynomorphs zones recognised in  
 Idogun well.



199

200 Figure 7: The palynomorph distribution Chart of in Lewumeji well (0.00 – 111m).



201

202 Figure 8: The palynomorph distribution Chart of in Idogun well (0.00 – 54m).

203 Table 1: The distribution of palynomorphs species recovered in Lewumeji well and the number  
204 of counts for taxaspécie type

SAMPLE NO	Depth (m)	Lithology	Palynomorphs recovered	Counts/Species Type
A1.1	0 - 15.00	Sandstone	Tubistephanocolpites Cylindricus	1 (P)
A1.2	15 - 45	Shale	Cyathidites sp Proteacidites sp Laevigatosporites sp	2(S) 1(P) 2(S)
A 1.3	45 - 75	Shale	Monocolpites Marginatus Leiosphaeridia sp Microforaminiferal wall lining Laevigatosporites sp	1 (S) 1 (DC) 1 (DC) 1(S)
A 1.4	75 - 111	Shale	Leiosphaeridia sp Cyathidites sp Cyathidites Minor Tubistephanocolpites Cylindricus	1 (DC) 1 (S) 1(S) 1(P)

**Legend**  
P - Pollen  
S - Spores  
DC - Dinoflagellates  
Sandstone  
Shale

205




206 Table 2: The distribution of palynomorphs species recovered in Idogun well and the number  
 207 counts for specie type of taxa

Sample no	Depth (m)	Lithology	Palynomorphs Recovered	Counts/ Species Type
B1.1	0 - 9.00	Clay	Cyathidites sp	1(S)
			Proteacidites sp	1(P)
B1.2	9.00 -15.00	Sandstone	Leiosphaeridia sp, Trilete spores	1 (DC), 2(S)
			Microforaminiferal wall lining, Cyathidites sp	1 (DC), 1 (S)
B1.3	15.00 -24.00	Shale	Foveotriletes Margantae, Laevigatosporite sp	1(P), 1 (S)
B1.4	24.00- 42.00	Sandstone	Distavertuspontes simplex	1 (S)
			Monocolpites Marginatus	1(P)
B1.5	42.00 - 54.00	Shale	Maunitidites Lehmani	1(P)
			Tubistephanocolpites Cylindricus	1(P)
B1.5	42.00 - 54.00	Shale	Pteris sp	1(S)
			Senegalium sp	1 (DC)
			Paleocyrodinium sp	4 (DC)
			Subtilisphaera sp	2(DC)
			Cyathidites sp	1 (S)
			Monocolpites Marginatus	1(P)
			Cerodinium sp	1 (DC)
			Oligosphaeridium sp	1(DC)
Monopontes Annulatus	2(P)			

LEGEND  
 S - Spore  
 P - Pollen  
 DC - Dinoflagellates cysts

Clay  
 Sandstone  
 Shale

208  
 209 Correlation of intervals (fig.9) within both wells using terrestrially sourced spores and pollen  
 210 shows alots of similarities, this suggests that the sediments were deposited under the same  
 211 conditions and the miospores might have come from the same origin during the same period and  
 212 sediments were partly deposited under the same condition.



213  
 214 Figure 9: Correlation chart of the study sections using the recovered palynomorphs from both  
 215 wells (a) Idogun well (b) Lewumeji well

### 216 3.2.2 Environment of deposition

217 Deduction of Paleoenvironment of deposition was carried out using different means based on  
 218 preferable environment of deposition of environmentally indicative forms, palynomorphs  
 219 frequency distribution, and comparison of land derived forms to marine source. The  
 220 palynomorphs frequency percentage distribution shows that both well has a higher frequency of  
 221 land derived miospore to the marine dinoflagellates; (75%, 25% and 61%, 39%) for Lewumeji

222 and Idogun well respectively. This suggests that the source of organomacerals are plants and  
223 environment of deposition is likely to be from a continental to brackish environment of  
224 deposition (Adeigbe et al., 2013).

225 The occurrence of environmentally indicative forms in Lewumeji and Idogun well such as  
226 Leiosphaeridae Sp indicative of neritic environment (Ayinla et al., 2013), *Monocolpites*  
227 *marginatus* suggestive of coastal plain habitat (Adeigbe and Amodu 2015), foraminifera wall  
228 linings suggestive of nearshore environments .However, the moderate records of fern spores  
229 such as *Cyathidites* sp., *Cyathidites minor* are indicative of open fresh water swamps (Lawal and  
230 moullade 1987). And the presence marine loving forms such as *Cerodinium* sp.,  
231 *Paleocystodinium* sp., and *Senegalinium* sp. and *Subtilisphaera* sp in Idogun well are indicative  
232 of shallow marine environments. This suggest a depositional environment that vary from  
233 continental to brackish to shallow marine environment with minor influx of freshwater.

234 The Palynomorphs marine Index which is a semi quantitative interpretation technique was  
235 employed to further determine the Interval of Idogun (0.00-54.00m) and Lewumeji well (0.00-  
236 111.00m). This method depends on the amount of terrestrial and marine palynomorphs  
237 separately, to deduce the paleoenvironments of fossil forms in respect of fluvial and marine  
238 environment. Helenes *et al.*, 1998 define PMI (Palynological Marine Index) as:

239  $PMI = Rm/Rt + 1 * 100$

240 Range of classification follows

241  $>100 =$  Fluvial environment

242  $100-200 =$ Fluvial/ marine environment

243  $>200 =$  Marine environment.

244 Where  $R_t$  = Richness/number of terrestrial palynomorphs (pollen + spores + Fungal remains)

245  $R_m$  = Richness/number of aquatic palynomorphs (Dinoflagellates+ Acritarch + foraminifera wall  
246 linings + Prasinophytes). High, Low and nil values of palynomorph marine index (PMI) indicate  
247 a marine, brackish and fresh water environment respectively (Chukwuma-Orji *et al.*, 2017).

248 Quantitative interpretation technique applied using Palynomorph Marine Index (PMI) values  
249 show that in Lewumeji well (table 3) PMI value of about 100, indicative of interval 0.00-  
250 111.00m which are equivalent to fluvial deposit due to dominance of land derived  
251 palynomorphs. The PMI values (table 4) show that in Idogun well, intervals with PMI values of  
252 about 100, indicative of interval 0.00-9.00m, 9.00-15.00m, 15.00-24.00m, and 24.00 -42.00m are  
253 equivalent to fluvial deposits, while the lowermost part with the depth range of 42.00 –

Comment [S27]: italic

Comment [S28]: indicate

Comment [S29]: italic

Comment [S30]: italic

Comment [S31]: italic

Comment [S32]: italic

Comment [S33]: italic

Comment [S34]: sp.


Formatted: French (France)

254 54.00m has a PMI value between 100-200 which is indicating an alternation of continental and  
 255 marine deposits. Therefore, from the general view of the PMI values against analyzed  
 256 stratigraphic interval (fig 10, fig.11), a brackish to Shallow marine environments with minor  
 257 freshwater incursions is suggested for the study area.

258 Table 3: Paleoenvironment Interpretation of Lewumeji well from P.M.I. Value of the Palynomorphs Distribution.

Sample No	Depth (m)	Pollen	Spores	Dinoflagellate Cyst	Total	PMI	Paleoenvironment
A1.1	0.00 - 15.00	1	0	0	1	0	Fluvial Deposit / Freshwater environment
A1.2	15.00 - 45.00	4	1	0	5	0	Fluvial deposit / Freshwater environment
A1.3	45.00 - 75.00	1	1	2	4	67	Fluvial deposit / Brackish environment
A1.4	75.00 - 111.00	1	2	1	4	25	Fluvial deposit / Brackish environment

259




260

261 Figure 10: Palynomorphs Marine Index (PMI) chart of Lewumeji well

262 Table 4: Paleoenvironment Interpretation of Lewumeji well from P.M.I. Value of the  
 263 Palynomorphs Distribution.

Sample No	Depth (m)	Pollen	Spores	Dinoflagellate Cyst	Total	PMI	Paleoenvironment
B1.1	0.00 - 9.00	1	1	0	2	0	Fluvial Deposit / Freshwater environment
B1.2	9.00 - 15.00	1	4	2	7	34	Fluvial deposit / brackish environment
B1.3	15.00 - 24.00	0	1	0	1	0	Fluvial deposit / Freshwater environment
B1.4	24.00 - 54.00	1	2	1	4	0	Fluvial deposit / freshwater environment
B1.5	42.00 - 54.00	3	1	9	13	180	C/Marine deposit / marine environment

264




265

266 Figure 11: Palynomorphs Marine Index (PMI) chart of Idogun well

267

268 **Plate 1: Some selected Palynomorphs photomicrographs recovered from Idogun well**




270

271 1. Proteacidites sp 2. Foveotriletes margaritae 3. Pteris sp 4 &5. Cyatidites minor 6.  
 272 Laevigatosporites sp 7. Monocolpites marginatus 8. Microforaminiferal wall linings  
 273 9. Distaverrusporites simplex 10. Oligosphaeridium sp.

274

275 **Plate 2: Some selected Palynomorphs photomicrographs recovered from Lewumeji-1 well**



276

277



278 1. *Cyathidites minor* 2. *Monocolpites marginatus* 3. *Leiosphaeridia* sp 4. *Foveotriletes*  
279 *margaritae* 5. *Tubistephanocolpites cylindricus* 6. *Laevigatosporites* sp 7. Microforaminiferal  
280 wall linings.

### 281 Conclusions

282 The palynological and Lithostratigraphy studies has been appropriately employed to study the  
283 sediments of Abeokuta group a part of Eastern Dahomey basin through the use of nine (9)  
284 composited core samples from Lewumeji and Idogun well with depth ranging from 0 -111m and  
285 0-54m respectively.

286 The wells were examined lithologically and five units were delineated which can be further  
287 grouped into three for Idogun well two units of shale, two units of sandstone and a clay unit  
288 while the lewumeji well has a lithology of sandstone and shale. Both well are dominated by  
289 fissile to blocky, light to dark grey colour shale and the sand grain varies from medium to fine  
290 grained texture and the clay unit covers a small interval having a reddish brown colouration. This  
291 lithology denote Marine, fluvial and Lagoonal or brackish environment respectively. The thirty-  
292 one (31) palynomorphs recovered within the two well are well preserved with low to moderate  
293 diverse pollen, spores and the dinoflagellates cysts. The microfloral assemblages include  
294 abundant *Cyathidites* sp, *Cyathidites minor*, *Tubistephanocolpites cylindricus*, *Proteacidites* sp,  
295 *Trilete* spore, *Foveotriletes margaritae*, *monocolpites marginatus*, *monoporites annulatus*, *pteris*  
296 sp, *Distaverrusporites simplex* and *Laevigatosporites* sp. The dinoflagellates recovered were  
297 characterized by the likes of *Leiosphaeridia* sp, *Senegalinium* sp, *Oligosphaeridium* sp,  
298 *paleocytodinium* sp, *Cerodinium* sp and *Subtilisphaera* sp. The palynological assemblage zone  
299 identified within the two wells is the *Cyathidites minor* zone, these zone is correlatable with the  
300 *Spinizonocolpites Bacculatus* zone of Lawal and moullade, (1987). The zone is characterized by  
301 the presence of *Monocolpites marginatus*, *Cyathidites minor* and *Cyathidites* Sp. The studied  
302 sediments from the wells were deposited in a continental to brackish to shallow marine  
303 environment with minor freshwater incursions during the Upper Maastrichtian – Early Paleocene  
304 period based on environmental diagnostic species, palynomorphs marine index and frequency  
305 distribution of palynomorphs.

306

307

308 (1) Adegoke, A.K., (2012). Biostratigraphy and depositional environment of the sediments in  
309 Bornu Bornu Basin, North Eastern Nigeria. *Indian J. Sci. Technol.* 5(6), 2800-2809.

### References

- 310 (2) Adeigbe, O.C., Ola-Buraimo, A.O., Moronhunkola, A.O., (2013).palynology and  
 311 biostratigraphy of Emi – 1 well, Moroh field deep offshore, Eastern Dahomey Basin,  
 312 Southwestern Nigeria. *International Journal of Scientific and technology research*, 2(1),  
 313 58 -70.
- 314 (3) Adeigbe, O.C., and Amodu O.I., (2015). Palyno- environmental study of the Araromi -1  
 315 well, Eastern Dahomey Basin, South- west Nigeria. *Journal of science research*, Vol. 14,  
 316 ISSN: 1119 7333
- 317 (4) Agagu, O.K., (1985). Geological guide to bituminous sediment in South Western Nigeria.  
 318 (Unpublished monograph), Department of Geology, University of Ibadan, Ibadan. Vol. 3,  
 319 Pp17.
- 320 (5) Awad, M.Z., (1994): Stratigraphic, palynological and palaeoecological studies in the  
 321 East-Central Sudan (Khartoum and Kosti Basins), Late Jurassic to Mid- Tertiary.  
 322 Berliner geowiss Abh
- 323 (6) Ayinla, H.A., Adeigbe, O.C., Ola-Buraimo, O.A., Bankole, S.A., Adebowale, M., (2014).  
 324 Palynostratigraphy and high resolution paleoenvironmental reconstruction of part of  
 325 Kemar-1 well, Bornu Basin, northeastern Nigeria. *The pacific journal of science and*  
 326 *technology*, 15 (1), Pp 392 – 403.
- 327 (7) Biliman, H.G., (1992). Offshore Stratigraphy and Paleontology of the Dahomey (Benin)  
 328 Embayment, West Africa. *NAPE BULL.*, Vol. 7, No. 2 Pp. 121 -130.
- 329 (8) Chukwuma- Orji J.N., Okosun, E.A., Ekom, J.C., Okosun, E.A., Ekom, J.C., Abolarin, J.F  
 330 (2017). A palynology and paleoenvironment of a section from the Amansiodo -1 well,  
 331 Anambra basin South eastern Nigeria. *Journals.agh.edu.pl/geol*, Vol. 43(4) Pp. 293-301.
- 332 (9) Coker, J.L., and Ejedawe, J.E. (1983). Dynamic interpretation of organic matter  
 333 maturation and evolution of the oil generative window. *AAPG Bulletin*, vol. 68(8),  
 334 Pp.1024-1028.
- 335 (10) Durugbo, E. U., & Aroyewun, R. F. (2012). Palynology and Paleoenvironments of the  
 336 Upper Araromi Formation, Dahomey Basin, Nigeria. *Asian Journal of Earth Sciences*,  
 337 5(2), 50-62.
- 338 (11) Enu, E.I., (1990). Nature and occurrence of Tar sands in Nigeria in: B.D. Ako and E.I.  
 339 Enu (eds), occurrence, utilization and economic of Nigerian tar sands: 1<sup>st</sup> workshop on  
 340 Nigerian tar sands, NMGS published Ibadan Chapter, Ibadan, Nigeria Pp. 11-16.
- 341 (12) Fayose, E.A., (1970): Stratigraphic paleontology of Afowo well 1 South Western  
 342 Nigeria. *Geol.Surv.Nig*.Vol. 5 Pp. 65-68.
- 343 (13) Helenes, J. Guerra, C., and Vasquez, J. (1998): Palynology and Chronostratigraphy of  
 344 the Upper Cretaceous in the subsurface of Barinas area, Western Venezuela. *AAPG*  
 345 *Bull.*, 82(7), 1308-1328.
- 346 (14) Kingston, D.R., Dishroon, C.P. and Williams, P.A., (1983): Global Basin classification  
 347 system. *American. Assoc. of petroleum Geologists .Bulletin*. Vol.67, Pp 2175 -219331,
- 348 (15) Lawal, O., and Moullade, M., (1987). Palynological Biostratigraphy of the Cretaceous  
 349 sediments in the upper Benue Basin, N.E. Nigeria. *Revue micropaleontologie*, vol. 29,  
 350 n.1, p. 61-83.
- 351 (16) Obaje, S.O. and Okosun, E.A. (2013). Paleoenvironmental interpretation of Tomboy  
 352 field, Offshore western Niger delta, Nigeria. *International journal of science and*  
 353 *technology*, Vol 2, No, 9, Pp. 628 -638.
- 354 (17) Ola-Buraimo, A. O. (2012): Lithostratigraphy and Palynostratigraphy of Tuma-1 well,  
 355 Bornu Basin, Northeastern Nigeria. *J. Biol. Chem. Res.* 29(2), 206- 223.
- 356 (18) Omatsola, M. E., and Adegoke, O.S., 1981: Tectonic Evolution and Cretaceous  
 357 Stratigraphy of the Dahomey Basin. Nigeria. *Journal of Mining and Geology* 18 (01),  
 358 130– 137.

- 359 (19) Salami M.B., (1987): Petrography and palynology of Upper Maastrichtian Abeokuta  
360 Formation of South- western Nigeria, *Nigerian journal of Science*. Vol. 21, Pp. 140-  
361 146.
- 362 (20) Whiteman, A.J., (1982). Nigeria, its petroleum Geology of Nigeria, resources and  
363 potential, Vol 2 Pp 223-230 Graham and Trotman, London.
- 364

365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385

UNDER PEER REVIEW