

Original Research Article

Antibacterial Activity of Saponin Extracted from *Phyllanthus niruri* on Methicillin-Resistant *Staphylococcus aureus* (MRSA)

ABSTRACT

The antimicrobial activity of saponin extracted from *Phyllanthus niruri* was investigated on methicillin-resistant *Staphylococcus aureus* (MRSA). The nuclear magnetic resonance (NMR) was used to determine the structure spectra of the extracted purified saponin. The ¹³carbon NMR predicted on the basis of chemical shift that appeared in the resonances of 20 – 60 ppm gave a structure named Phylagenin-13-O-α-D-glucopyranoside and Phylagenin-25-O-β-D-glucopyranoside. The susceptibility profile of MRSA determined by the agar-diffusion method showed that 97.0% and 90.0% of the test bacterium were resistant to Tetracycline and Cotrimoxazole respectively and 60% of the bacterium was susceptible to saponin extract. The ability of saponin extracted from *P. niruri* to treat clinical manifestation like chest congestion and skin desquamation from which *S. aureus* resistant to conventional antibiotics have been isolated has been confirmed in this study. The fact that this extract exerted an inhibitory effect on MRSA indicates that they can potentially be further developed into antimicrobial clinically used agents.

Keywords: Antibacterial, Methicillin-Resistant, *Phyllanthus niruri*, Saponin, *Staphylococcus aureus*

1. INTRODUCTION

Historically, plants have provided a source of inspiration for novel drug compounds, as plant-derived medicines have made large contributions to human health and well-being [1]. Their roles are **two-fold** in the development of new drugs. They may become the base for the development of a medicine, a natural blueprint for the development of new drugs or, a phytomedicine to be used for the treatment of disease [2].

Higher plants especially tropical species, produce secondary metabolites with antibacterial activity [3]. Among higher plants, *Neurolaena lobata* and *Aristolochia* species which are commonly used to treat infections in Belizean folk medicine, have been shown to have activity against *Staphylococcus aureus*, *Escherichia coli*, *Proteus vulgaris*, and *Pseudomonas aeruginosa* [3].

Phyllanthus niruri L., (Syn. *P. fraternus* Webster) (Plate 1) is a common kharif (rainy season) weed found in both cultivated fields and wastelands [4]. It carries different nomenclature in different parts of the world. However, in Nigeria it is called *Asasa* or *Arunjeran* in Yoruba, *Majiryar Kurumi* in Hausa, *Asivi* or *Igbehen* in Edo, *Egu eza* in Ibo and *Oyomo-ke-iso-aman-ke-edem* in Efiki [5]. Although considered a problematic weed for farmers, it is a valuable medicine for herbalist [6] and holds a reputable position in both Ayurvedic and Unani

systems of medicine. Recently it has attracted the attention of researchers, because of its hepatoprotective properties [6]. Although no effective specific therapy is available for viral hepatitis, *P. niruri* has shown clinical efficacy in the treatment of viral Hepatitis B [7].



Plate 1: *Phyllanthus niruri* L. (Syn *P. fraternus* Webster)

Saponins are naturally occurring surface-active glycosides that are produced by plants. They derive their name from their ability to form stable, soap-like foams in aqueous solutions. This easily observable character has attracted human interest from ancient times [8]. Saponins are known to be antimicrobial, to inhibit mold, and to protect plants from insect attack [9, 10]. Saponins may be considered a part of plants' defense system, and as such have been included in a large group of protective molecules found in plants named phytoanticipins' or 'phytochemicals' [11]

A large number of the biological effects of saponins have been ascribed to their action on membranes. In fact, their specific ability to form pores in membranes has contributed to their common use in physiological research [12, 13]. Saponins have long been known to have a lytic action on erythrocyte membranes and this property has been used for their detection. The hemolytic action of saponins is believed to be the result of the affinity of the aglycone moiety for membrane sterols, particularly cholesterol [10, 14] with which they form insoluble complexes.

Methicillin-resistant *Staphylococcus aureus* (MRSA) is a specific strain of *S. aureus* bacterium which is intrinsically insensitive to methicillin and all **beta-lactamase** (β -lactamase) antibiotics like dicloxacillin, cloxacillin, nafcillin, penicillin, and oxacillin [15]. Since 1960s, successive waves of epidemic **methicillin-resistant** *S. aureus* have spread out throughout hospitals and other chronic health care facilities worldwide, to the extent that it is now the most commonly isolated antimicrobial resistant pathogen in many countries [15, 16].

The mechanism of methicillin resistance is an altered **penicillin-binding** protein (PBP2a) in MRSA that markedly reduces affinity for all available **beta-lactamase** antibiotics, while maintaining effective cell wall binding activity. The **PBP2a** is encoded by the *mecA* gene that is carried on a mobile DNA element, the staphylococcal cassette chromosome [17].

Comment [DM1]: Please provide a high resolution picture of the plant without any intervention

Healthy individuals may carry MRSA asymptomatically for periods ranging from a few weeks to many years. Patients with compromised immune systems are at a significantly greater risk of symptomatic secondary infection [18]. MRSA may progress substantially within 24 hours to 48 hours of initial topical symptoms, after 72 hours. MRSA can remain in human tissues and eventually become resistant to treatment.

Despite successes achieved in controlling many infectious diseases, efforts to defend against the wide range of microbes that threaten human health continued to be challenged by emerging and re-emerging of infectious pathogens and possible use of a variety of virulent agents as biological weapons [19]. A defensive strategy based solely on developing new vaccines and antimicrobial and antiviral drugs, each specific for only one or a few agents, is unlikely to be successful in dealing with potential microbial threats and these are exceedingly expensive [20]. Therefore, the objective of this study is to extract crude saponin from the whole plant of *P. niruri* and determine the *in-vivo* antimicrobial potency and efficacy on strains of MRSA.

Comment [DM2]: Several plants have numerous classes and categories of saponin.....is there any specific type of saponin mentioned in literature or authors tried to study the physicochemical aspects of the saponin

2. MATERIAL AND METHODS

2.1 Collection of Plant Material

The matured and fresh leaves of the plant, *Phyllanthus niruri* were collected from farmlands in Ado-Ekiti during the raining season between the months of May and October (2004 – 2006). The plant material was air-dried at room temperature ($27^{\circ}\text{C} \pm 1^{\circ}\text{C}$), was ground into the powdered form using milling machine (Retsch GmbH 5657 HAAH) and stored in air-tight plastic container. Identification and authentication of plant were performed in the Department of Plant Science, University of Ado-Ekiti, Nigeria and a voucher specimen was deposited.

Comment [DM3]: Provide specimen number

2.2 Extraction of Saponins

The method described by Martson *et al.* [21] was employed. The dried and powdered plant material (500 g) was defatted in a Soxhlet with petroleum ether at between 40°C and 60°C for 16 h. The residue was added to 100 mL of absolute methanol and left overnight under reflux at 70°C . It was filtered with Whatman No. 2 mm filter paper and the filtrate evaporated to dryness with a rotatory evaporator. The yield was dissolved in 100 mL of distilled water, extracted in a separating funnel with 1-butanol three times and dried by evaporating. Finally, the extract was dissolved in 25 mL of absolute methanol and the saponins compound was precipitated by adding 75ml of diethyl ether.

Comment [DM4]: There are some confirmatory test for the saponins, such as blood lysis test.....have you done those tests for the confirmation? Provide a section: physicochemical characterization

2.3 ^{13}C Carbon Nuclear Magnetic Resonance (NMR) spectral analysis of saponin

The ^{13}C -NMR was used to determine the structure spectra of the purified saponin extract. A dilution of 10 mg of the saponin was made in 10ml of distilled water. This was injected into the vial of the equipment. The analysis was done at standard conditions of optical rotations of 20° taken on a Perkin Elmer 241 polarimeter, Pulse of 92.9° , total time of 9hr, 20 min, 51 sec; WALTZ-16 modulated and 19237 repetitions.

Comment [DM5]: Write about the NMR system used...model or make?

2.4 Isolation and identification of Methicillin-Resistant *Staphylococcus aureus* (MRSA)

From the year 2005 through 2007 samples were collected from patients visiting State Specialist hospitals in Ado-Ekiti, Ikere-Ekiti, Ifaki-Ekiti, and Medical Centre, Ido-Ekiti; in Ekiti

State, Nigeria. The samples collected included wound/pus, sputum, urine, respiratory swabs, gynecologic specimens, and stool, synovial fluid. Samples for MRSA detection were put on plates. *S. aureus* was confirmed by bound coagulase test. Isolates found to be oxacillin resistant ($\leq 10\text{mm}$) or have intermediate resistance to oxacillin ($11 - 12\text{ mm}$) on a Kirby-Bauer disk diffusion assay were further tested by E test on Mueller-Hinton agar with 2% Sodium chloride incubated for 24 h, at 37°C . Those with a $\text{MIC} \geq 4\text{ }\mu\text{mol/mL}$ were considered to be MRSA [22].

2.5 Determination of antibacterial potency of crude saponin

The disk diffusion method described by Brady and Katz [23] was employed. Various concentrations of the extract were prepared on test tubes ($0.625 - 10\text{ }\mu\text{g/mL}$). Disks obtained from Whatman No 5 MM filter paper were sterilized in an oven at 60°C for 30 minutes and soaked in the extract for 2 hrs. A loopful of the final dilution (10^3) of the test bacterial suspension was spread on dried nutrient agar (Oxoid). The disks of different concentrations of the extracts were placed equidistance on the agar and incubated at 37°C for 24 hrs. Zones of inhibition were measured in milliliters with a meter rule. Each procedure was repeated three times.

2.6 Determination of antibacterial efficacy of crude saponin

This was carried out using the agar dilution method as described by Smith *et al.* [3]. A colony from stock was sub-cultured into 5 mL of nutrient agar (LAB) and incubated at 37°C for 18 h. Overnight broth (0.1 mL) of each organism was pipette into 9.9 mL of the broth to yield 10^1 dilutions. Serial dilution was carried out to obtain a dilution of 10^3 . A 2cm streak of bacteria strains was made on dried nutrient agar plates containing increasing concentrations ($0.625\mu\text{g/mL}$ - $10\mu\text{g/mL}$) of the extract. The lowest concentration that gave no visible growth after overnight incubation at 37°C was taken as the minimum inhibitory concentration (MIC) of the crude extract.

3. RESULTS AND DISCUSSION

3.1 Results

The $^{13}\text{C-NMR}$ spectra result of the purified saponin extract from *P. niruri* is shown in Figure 1. The site of attachment of one sugar to another sugar can be predicted on the basis of chemical shifts in resonances of $20 - 60\text{ ppm}$. A close resemblance of the chemical shifts due to a terminal sugar with respect to a methyl-O-glycoside lead to its immediate characterization whereas the chemical shift of other (inner) sugars differ significantly in comparison to methyl-O-glycosides. In oligoglycosides, the glycosylation causes a downfield shift of $20 - 60\text{ ppm}$ of the α -carbon, the hydroxyl of which has been directly involves in the glycosylation while neighboring β -carbon atoms show an upfield shift of $80\text{-}100\text{ppm}$. These α - and β -shifts are independent of the nature of the monosaccharide and provide a conclusive method for the establishment of interglycosidic linkages.

The susceptibility profile of MRSA is shown in Table 1. Twenty-nine 29(97%) of 30 isolates showed resistance to tetracycline while only 2 (7%) showed resistance to oflotarivid. The highest susceptibility of 25 (83%) was observed in oflotarivid. The isolates were also susceptible to gentamicin 13(43%) and nitrofunrantoin 12 (40%).

Comment [DM6]: How these chemical components play role in expressing biological activities? An Expert Opinion by the authors must be preferred

The susceptibility of MRSA to the crude saponin is shown in Figure 2. Of the 30 isolates, 18 (60%) isolates showed a zone of inhibition of diameter ranging between of ≥ 9.0 mm and 5.0 mm while 12 (40%) of the isolates showed a zone of inhibition of diameter ≤ 4.0 mm.

The antibacterial activity of different concentration of crude saponin extract to MRSA is shown in Figure 3. At a higher concentration (1.5 mg/mL), the optical density the broth culture decreases from an initial optical density of 0.02 nm to 0.30 nm. The control culture shows a progressive increase in the optical density to 1.20 nm at an interval of 5 hours incubation. There is evidence of increased antibacterial activity as the concentration of the extract increases.




Fig. 1. The ¹³Carbon NMR Spectra of the crude saponin extract

Table 1. Susceptibility profile of MRSA (n = 30)

Antimicrobial agent (μg)	Resistance Pattern		
	S (%)	I (%)	R (%)
Amoxicillin	2 (6.8)	9 (30)	19 (63.3)
Cotrimoxazole	1 (3.3)	2 (6.8)	27 (90)
Nitrofurantoin	12 (40)	4 (13)	14 (47)
Gentamicin	13 (43)	12 (40)	5 (17)
Nalidixic acid	11 (37)	11 (37)	8 (27)
Oflotarivid	25 (83)	3 (10)	2 (7)
Tobramycin	6 (20)	8 (27)	16 (53)
Tetracycline	0 (0)	1 (3.3)	29 (97)

S – Sensitive, I – intermediate, R – resistant




Fig. 2. Susceptibility of strains of **MRSA** to saponin extract of *Phyllanthus niruri* at a concentration of 1.5 mg/mL




Fig. 3. The **optical** activity of different concentration of crude saponin extracts on **MRSA**

3.2 Discussion

The resistance pattern of *S. aureus* showed that a high percentage of this isolate is resistant to most of the commonly used antibiotics in the classes of **β -lactams** (amoxicillin),

Comment [DM7]: Please provide the scale in the figure

tetracycline, folate inhibitors (co-trimoxazole), aminoglycosides (tobramycin and nitrofurantoin). It has been reported that similar to pathogenic bacteria, commensals are exposed to the selective pressure of antimicrobial agents [24]. The findings of this study were in substantial agreement with previous reports of high resistance rates of bacteria from low resource settings [22, 25].

In recent years, outbreaks of MRSA infection have been reported in different settings, outpatients and settings visiting hospitals, including hospitalized or surgery patients [26], as well as nursing homes and convectional facilities [22]. The problems of MRSA are increasing worldwide. MRSA is no longer restricted to hospital settings but is found in homes, places of work and kindergartens [27]. A number of risk factors for MRSA infection have been identified in those studies to include antimicrobial drug use, close contact with persons colonized with MRSA and barriers to medical care. Antibiotic drug self-medication is a cause for concern because it has contributed to the spread of antimicrobial drug resistance. Self-treatment with a drug that is ineffective against a disease causative organism or with an inappropriate dosage may increase the risk of selection of resistant organism that may be difficult to eradicate. These resistant organisms may then be transferred into the community. Unrecognized community associated-methicillin resistant *Staph aureus* (CA-MRSA) colonization during hospitalization could become an additional method of its dissemination in the community. Increased prevalence of CA-MRSA has been reported in Chicago, Los Angeles, Texas, and Minnesota [28].

Over the last 60 years or more, bacteria and, in particular, those pathogenic for humans have evolved toward antimicrobial drug resistance. Humans cannot control emergence because it occurs by chance and represents a particular aspect of bacterial evolution. Emergence can result from mutations in house-keeping, structural or regulatory genes or from acquiring foreign genetic information [20]. However, much can be done to delay the subsequent spread of resistance. Dissemination can occur at the level of the bacteria (clonal spread), replecons (plasmid epidermis), or of genes (transposons). These three levels of dissemination, which occur in nature, are not only infectious but also exponential, since all are associated with DNA duplication.

The activity of crude saponin extract showed high efficacy on multidrug-resistant *S. aureus* 60% of the isolates showing susceptibility. The highest percentage of resistance (40%) was observed. These findings showed that saponin extract from *P. niruri* is potent against multidrug-resistant *S. aureus* and it further substantiates the findings of Sen *et al.* [29] where it was observed that the growth of *S. aureus* was lowered at levels of 0.25% (w/v) by *Quillaja saponaria* saponin. Saponin was reported to have the potential to modulate microbial growth [30]. Membrane fluidity controls the enzyme activity of biological membranes and plays an important role in ion transport [31] and also controls the ability of saponins to affect the cellular function of bacteria.

4. CONCLUSION

The ability of saponin extracted from *P. niruri* to treat skin infection and pneumonia has been observed in this study. These favorable effects point to the potential of the saponin as a remedy against these two major health hazards in many countries including Nigeria. The fact that this extract exerted an inhibitory effect on MRSA indicates that they can potentially be further developed into antimicrobial agents.

Comment [DM8]: Saponins are considered toxic and even trials have been denied by several regulatory bodies with saponin based formulations.....write your own idea or hypothesis or an opinion of how these toxic components that even lyse human RBC on administration, can be used safely for clinical applications in the future?

Comment [DM9]: Please provide a more realistic conclusion with a direction towards the future perspective

ETHICAL APPROVAL (WHERE EVER APPLICABLE)

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

REFERENCES

1. Ang-lee M, Moss J. and Yuan C. Herbal medicines and perioperative care. *JAMA*. 2001; 286:208-216
2. Jiang W. Therapeutic wisdom in traditional Chinese medicine: a perspective from modern science. *Trends in Pharmacological Science*. 2005; 26:558-563.
3. Smith RA, Calviello CM, DerMarderosian A, Palmer ME. Evaluation of antibacterial activity of Belizean Plants: an improved method. *Pharmaceutical Biology*. 2000; 38:25-29.
4. Ajibade VA, Ajenifuja OA, FT Akinruli, Ajayi FA and Famurewa O. Antifungal efficacy of saponin extracted from *Phyllanthus niruri*. *International Journal of Pathogen Research*. 2018; 1(3): 1-8
5. Irvine FR. West African Botany (3rd edition) Oxford University Press, Amen House, London E.C.4; 1986.
6. Qudhia P, Tripathi RS. Prospects of cultivation of medicinal plants in Chattisgarh, India. In: Recent progress in medicinal plants, crop improvement, production technology, trade and commerce. Sci Tech publ, USA. 2002;5:211-236.
7. Paranjpe P. Indian medicinal plants: Forgotten healers. Chaukhamba Sanskrit Pratisthan, Delhi. 2001;48-49.
8. Yoshikawa M, Murakami T, Kishi A, Kageura T, and Matsuda H. Medicinal flowers 111, marigold (1): hypoglycemic gastric emptying inhibitory, and gastro protective principles and new oleanane-type triterpene oligoglycosides, calendasaponins A,B,C and D, from Egyptian *Calendula officinalis*, *Chemical and Pharmaceutical Bulletin*. 2001; 49:863-870.
9. Flaoven A, Wilkins AL, Deng D, and Brekkt T. Ovine metabolisms of saponin: evaluation of a method for estimating the ovine uptake of steroidal saponins from *Nartheicum ossifragnum*. *Veterinary Research Communications*. 2001; 25: 225-238
10. Rai S, Wahile A, Mukherjee PK, Saha BP and Mukheek A. Antioxidant activity of *Nelumbo mucifera* (sacred lotus) seeds. *Journal of Ethnopharmacol*. 2006; 104:322-327.
11. Morrissey JP and Osbourn AE. Fungal resistance to plant antibiotics as a mechanism of pathogenesis. *Microbiological and molecular Biological Reviews*. 1999; 63:708-724.
12. Choi S, Jung SY, Kim CH, Kim HS, Kim SC and Nah SY. Effect of ginsenosides on voltage dependent Ca^{2+} channel subtypes in bovine chromatin cells. *Journal of Ethno pharmacology*. 2001; 74:75-81
13. Plock A, Sokolowska-Kohler W, and Presber W. Application of flow cytometry and microscopical methods to characterize the effect of herbal drugs on *Leishmania* spp. *Experimental Prarsitology*. 2001; 97:141-153
14. Mukherjee PK. Plant products with hypocholesterolemic properties. In Taylor, Steve L. (Ed). Advance food and Nutrition Research, 47. Elsevier Science U.S.A. 2003; pp277-338.

Comment [DM10]: Please arrange them according to the instructions provided by the journal

15. Diekema DJ, BootsMiller DJ, Vaughn TE, Woolson RF, Yankey JW. Antimicrobial resistance trends and outbreak frequency in United States Hospitals. *Clin. Infect. Dis.* 2004; 38: 78-85.
16. Goossens H. European status of resistance in nosocomial infections. *Chemotherapy.* 2004; 51: 177-181.
17. Katayama Y, Ito T, Hiramatsu K. "A new class of genetic element, *Staphylococcus* cassette chromosome mec, encodes methicillin resistance in *staphylococcus aureus*". *Antimicrob. Agents Chemother.* 2000; 44: 1529-1555.
18. Ajibade VA, Fajilade TO and Famurewa O. Incidence and *in vitro* Susceptibility of Methicillin- Resistant *Staphylococcus aureus* Isolated from Ekiti State to Saponin Extract from *Phyllanthus niruri*. *J. Pharm. Biomed. Sci.*, 2010; 1(1): 1-6.
19. Douglas WM, Brian DG, William BB, Shukal B, Paul OG, Paul H, Segarra N. Population Mobility, Globalization, and Antimicrobial Drug Resistance. *Emerg. Infect. Dis.* 2009; 15(11):1727-1732
20. Smolinski MS, Hamburg MA and Leaderberg J. Microbial threats to health: emergence, detection, and response. Washington: Institute of Medicine. 2003; p. 32.
21. Martson A, Wolfender JL and Hostettmann K. Analysis and isolation of saponins from plant material. In saponins in food, feedstuffs and medicinal plants. *Annual proceedings of the Phytochemical Society.* 2000; 120-138.
22. Louiselle L, Jacques P, Krystel T, Marie-France Q, Marie-Andree C, Marie-Rier C, and Marie-Ere A. Flouroquinolones and risk for Methicillin-resistant *Staphylococcus aureus*, Canada. *Emerg. Infect. Dis.* 2006; 12 (9):1398-1405
23. Brady MS and Katz SE. Factors influencing optimization of diffusion assays for antibiotics. *J. Assoc official Anal. Chemist.* 1990; 73: 202-205
24. Okeke IN, Lamikanra A and Edelma R. Socio – economic and behavioural factor leading to acquired bacterial resistance to antimicrobial agents in developing countries *Emerg. Infect. Diseases.* 2000; 5:18-27
25. Philip VG, Loreen AH, Bret A, Mary D, Kristopher H, Gary D, Patricia W, Diana DV and Daniel D. Community-associated methicillin-resistant *Staphylococcus aureus* Iowa, USA. *Emerging infectious Diseases.* 2009; 15(10):1582-1589.
26. Moel GS, Jones RN, Biedenbach DJ, Shlwell MG and Fritsche TR. Contemporary causes of skin and soft tissues infections in North America, Latin America, and Europe: report from the SENTRY Antimicrobial Surveillance Program (1998-2004). *Diagn Microbiol Infect Dis.* 2007; 57: 7-13
27. Wulf MM, Sorum M, VanNes A, Skov R, Melchers WJ and Klaassen CH. Prevalence of methicillin- resistant *Staphylococcus aureus* among veterinarians: an international study. *Clin. Microbiol Infect.* 2008; 14 :29-34.
28. Fergie JE and Purcell K. Community-acquired methicillin- resistant *Staphylococcus aureus* infections in South Texas children. *Pediatr Infect. Dis. J.* 2001; 20:860-863.
29. Sen S, Makkar HPS, Muetzel S and Becker K. Effect of *Quillaja saponaria* saponins and *Yucca schidigera* plant extract on growth of *E. coli*. *Letters in Applied Microbiology.* 1998; 27(1): 35-38.
30. Apers S, Baronikova S, Sindambiwe JB, Witvruw ME, Vander BD, Vanmarck E, Bucetinck A and Peters I. Antiviral, haemolytic and muluscidal activities of triterpenoid saponins from *Maesa lanceolata*: establishment of structure activity relationships. *Planta medica.* 2006; 67:528-532.

31. Ma LY and Xiao PG. Effects of *Panax notoginseng* saponins on platelets aggregation in rats with middle cerebral artery occlusion or *in-vitro* and on lipid fluidity of platelet membrane. *Phytotherapy Research*. 1998; 12:138-140.

UNDER PEER REVIEW