

Prevalence of Soil Transmitted Helminthes (STHs) Among Pupils of Community Primary Schools in Nkpor and Mgbodohia Communities in Rivers State, Nigeria.

ABSTRACT

Soil transmitted helminthes (STHs) are common public health concern among children in Sub saharan Africa. A study to determine the prevalence and intensity of these parasites among pupils of two primary schools in Nkpor and Mgbodohia communities, Obio/Akpor Local Government Area, Rivers State, Nigeria was conducted. The formo-ether concentration technique was used to concentrate and separate the eggs and cysts from the faeces. Out of 107 pupils (56 males and 51 females) investigated, 81 (75.7 %) were positive for at least one gastrointestinal helminth. The parasites identified included *Ascaris lumbricoide* (60.5%), Hookworms (23.5%), *Trichirus trichiura* (13.6%) and *A. lumbricoide* + *T. trichiura* (2.5%). *Ascaris lumbricoide* was significantly ($P = .05$) higher in prevalence than other parasites. Although more males (53.1%) were infected than females (47.0%), there was no significance ($P = .05$) difference in the prevalence in relation to sex. Of the 43 males infected, 26 (46.4%), 6 (10.7%), 11 (19.6%) and 0 (0%) harboured *A. lumbricoide*, *T. trichiura*, Hookworms and mixed infection (*A. lumbricoide* + *T. trichiura*) respectively. Out of the 38 females infected, 23 (45.1%), 5 (9.8%), 8 (15.7%) and 2 (3.9%) harboured *A. lumbricoide*, *T. trichiura*, Hookworms and mixed infection (*A. lumbricoide* + *T. trichiura*) respectively. There was no significance difference in the prevalence of *A. lumbricoide* between males (46.4%) and females (45.1%). There was significant difference in prevalence of soil transmitted helminthes (STHs) among two major age groups (5-10 years -47.0% and 11-15 years -44.4%). Children within the age group of 16-20 years had the least infection (8.6%). The intensities of *A. lumbricoide*, *T. trichiura* and hookworm were 246.5, 107.5 and 187 Epg respectively. The intensity of *A. lumbricoide* was significantly higher than other parasites identified. Soil transmitted helminthes remain a public health concern among children in the study area. Provision of portable water, toilet facilities, good education on the epidemiology of STHs and regular de-worming exercise will enhance control measures.

Keywords: Prevalence, Intensity, Soil-transmitted helminth, Nkpor, Mgbodohia

1.0 Introduction

Soil-transmitted helminths are a group of gastrointestinal parasites whose developmental stages requires the soil environment, that are transmitted to human through contact with soil, food or water contaminated with infective stages of the parasites. Soil [1]. The eggs of these parasites are passed out via faeces of infected persons into the environment, hence contaminating the soil in areas with poor sanitation especially where there are no modern toilet facilities. The four major gastrointestinal parasites known to cause STH infections include *Ascaris lumbricoides* (roundworm), *Trichuris trichura* (whipworm), and the species of hookworm (*Anclostoma duodenale* and *Necator americanus*) [2]. Helminthiasis (Ascariasis, trichuriasis and hookworm disease) caused by these parasites are the most prevalent in developing countries of the world possibly because of poor sanitary condition coupled with poverty and other environmental factors that encourage the thriving of the parasites [1,2].

The World Health Organization recorded that an estimated 1.5 billion people or 24% of global population are infected with soil-transmitted helminths worldwide [3], out of which an estimated 300 million cases resulted in severe morbidity [4]. In 2010, about 438.9 million people were infected with hookworm, 819.0 million with *A. lumbricoides* and 464.6 million with *T. trichiura* [5,6,7]. Ascariasis caused an estimated 2,824 million deaths, a large percentage of which occurred in Asia [6]. In 2016, one-third of an estimated three billion people living on less than two US dollars per day in developing countries of the world including Sub-Saharan Africa were at risk of at least one soil transmitted helminthic infection [8,9,2].

In sub-Saharan Africa , about 866 million people were infected with STH in 2012. School children are the most vulnerable group at risk of STH due to the habit of walking and playing barefooted, coupled with poor access to nutritional foods, poor awareness about routes of STH transmission and lack of access to basic water, sanitation and hygiene resource [5,10-13]. STH infections impact heavily on the physical growth and cognitive development of children [14,15].

Globally, an estimated 267 million pre-school-age children and over 568 million school- age children live in endemic areas and are in need of treatment and preventive interventions [9]. These infections cause a wide range of abdominal complications, iron-deficiency anaemia and dysentery syndrome in children [16]. [17] reported that STH weaken the nutritional status of the infected individuals in many ways including feeding on host tissues and blood leading to loss of iron and protein (anaemia) as in the case of hookworm, increased malabsorption of nutrients and competition for vitamin A. *Trichiura trichiura* causes dysentery and diarrhea while others have been implicated in loss of appetite which invariable reduce intake of nutrients causing poor physical growth.

In 2001, delegates at the World Health Assembly unanimously endorsed resolution WHA 54.19 which urged member countries in endemic regions to give serious attention to the tackling of STHs with the view to control morbidity through periodic deworming of school children and other persons at risk of the disease. In complementing preventive chemotherapy, the resolution also

recommends provision of adequate sanitation and personal hygiene as intervention measures for the control of STHs, and recommended albendazole and mebendazole as drugs of choice for the treatment of infected persons [17].

In Nigeria, several studies have been conducted on the prevalence of STH in different regions of the country[11,17,18]. In the study area, However, there is currently published no evidence on the prevalence of STH parasites in Nkpor and Mgbodohia communities in Obio/Akpor local government area in Rivers State. This study therefore provided such evidence to guide STH control programme officers and policy markers in the area.

2.0 MATERIALS AND METHODS

2.1 Study Area

Nkpor and Mgbodohia are two communities in Obio/Akpor Local Government Area. The communities have common boundaries and share common facilities(reservoir for water supply, road, electricity and drainage system). They are clans within Rumuolumeni kingdom and are host to several multinational oil companies including Agip oil company and Eni group that have several subsidiaries. The Local Government Area is basically a low land area, about 30 meters above sea level. It covers about 100sq mi (260 km²); and due to high rain fall, the soil consists of sandy or sandy loam [23].The vegetation consist of light rain forest and thick mangrove forest. According to the 2006 Census, it has a population of 464,789[24]. It is located between latitudes 4°45'N and 4°60'N

and longitudes 6°50'E and 8°00'E. Covering around 100 sq mi, Obio-Akpor is a lowland area with an average elevation below 30 meters above sea level.

UNDER REEER REVIEW

2.2 Study Population

The study subjects were primary school pupils between the age of 5 to 20 years. The only two public primary schools in the area (Community Primary school, Mgbodohia and Community Primary School, Nkpor) were selected for the study. The study was conducted between November 2017 and January, 2018.

2.3 Sample size

The sample size was estimated from the population of each school, that was below the age of 20 years. Official school records in both schools indicated that Community Primary School (CPS), Nkpor and Community Primary School (CPS), Mgbodohia had a population size of 63 and 55 pupils respectively. Out of this population, 57 pupils in CPS, Nkpor were below the age of 20 years while 50 pupils in CPS, Mgbodohia were below the age of 20 years. These became the sample size for the respective schools.

2.4 Sample Collection

A total of 150 pupils were randomly selected for study and the method of [25] was adopted in the collection of faecal samples. Well labeled plastic specimen bottles containing 10% formaldehyde and wooden scapula were distributed to the pupils, for collection of their faecal sample. The pupils were properly instructed on how to carefully collect their faecal samples into the specimen bottles. The samples were transported to the Research Laboratory, Department of Biology, Ignatius Ajuru University of Education for parasitological examination. Furthermore, demographic information on the age and sex of participants were

collected from the school record, the consistency of the stool (formed, soft, semi-soft and watery) were recorded for each pupil in a prepared recording format.

2.5 Parasitological Examination

Parasitological examination of the faecal samples was done using formo-ether concentration technique [26] to concentrate the parasites. About 1g of each faecal sample was thoroughly mixed in a test tube containing 4ml of 10% formol water and sieved through cotton gauze into centrifuge tube containing another 4ml of formol water. 3ml of diethyl ether was then added and thoroughly mixed. The mixture was then centrifuged for one minute at 3000rpm. The sediments were transferred onto a microscopic slide after the supernatant was discarded. Parasites eggs and cyst were observed and identified using X10 and X40 magnification of the microscope.

2.6 Ethical Consideration

The head teachers and Parent Teachers Association (PTA) of the selected schools were formally contacted in writing with regard to the research work. This was followed by official visit during which detail explanation on the essence of the research work made to the management of the schools and PTA. Further, the researchers interacted with the pupils at the early morning assembly ground during which orientation was given to the pupils about the research work. Subsequent upon this, written informed consent was also obtained from the Head Teachers and Parent Teacher Associations (PTA) of the schools investigated. Another approval was also given to the work by the Ethics Committee, Ignatius Ajuru University of Education, Port Harcourt.

2.7 Data Analysis

The prevalence rate (Pr) was determined by dividing the number of infected pupils (Ni) by the total number of pupils examined (Ne) multiplied by 100.

$$\text{Pr} = \frac{\text{Ni}}{\text{Ne}} \times 100$$

Parasite intensity (Pi) among the pupils was calculated as the number of eggs (Ng) per gram of faeces (Fg) while the parasite load was expressed as the mean number of eggs of each parasite species per gram of faeces of each pupil.

$$\text{Pi} = \frac{\text{Ng}}{\text{Fg}}$$

$$\text{Parasite Load} = \frac{\text{Mean number of eggs of parasite species}}{\text{Gram of faeces}}$$

Statistical analysis of the data was done using SPSS (version 16.0). Chi-square test was also used to compare difference among variables at significance level of $p = .05$.

3.0 Results

Out of the 150 pupils that received specimen bottles, only 107 returned their faecal samples. Of the 107 samples (56 males and 51 females) received and examined, 81 (75.7%) were positive for at least one helminthic infection (Fig. 1). Of the 81 positive samples, 43 (53.1%) were females while 38 (47.0%) were males. Of the 43 males infected, 26 (46.4%), 6 (10.7%), 11 (19.6%) and 0 (0%) harboured *A. lumbricoides*, *T. trichiura*, Hookworms and mixed infection (*A. lumbricoides* + *T. trichiura*) respectively (Table 1). Similarly, of the 38 females positive for helminth parasites, 23 (45.1%), 5 (9.8%), 8 (15.7%) and 2 (3.9%) harboured *A. lumbricoides*, *T. trichiura*, Hookworms and mixed infection (*A. lumbricoides* + *T. trichiura*) respectively (Table 1). There was no significance in

the prevalence of *A. lumbricoide*s between males (46.4%) and females (45.1%). However, there was statistical significance in the prevalence of *T. trichiura* between males (10.7%) and females (9.8%). Similarly, there was a statistical significance in prevalence of mixed infection between males (0%) and females (3.9%).

The results indicated that there was no significance difference ($p=.05$) in the prevalence of the parasites between the two major age groups, 5-10years (47% and 11-16years (43.2%). However, children that were above 17years old had the least infection (9.9%). Similarly, *A. lumbricoide*s had no significance prevalence across age group (Table 1). No significance difference in prevalence of Hookworms between age group 5-10years (21.1%) and age group 11-16years (25.7%). However, a statistical difference was observed in the prevalence of mixed infection (*A. lumbricoide*s + *T. trichiura*) among children above 17years old and the rest of the age groups (Table 1). Out of the 81 children positive for STH, 49 had *Ascaris lumbricoide*s, 19 had Hookworm, 11 had *Trichirus trichiura* while 2 had mixed infection (*A.lumbricoide*s + *T. trichiura*) representing 60.5%, 23.5%, 13.6% and 2.5% of the total worm burden respectively (Table 1). The infection was statistically significance for *A. lumbricoide*s with Nkpor (56.3%) higher than Mgbuodohia (46.8%). There was a significance difference ($p=.05$) in prevalence of STHs between Nkpor (59.3%) and Mgboudohia (40.7%). The prevalence of hookworms (23.5%) in the study area was not significance. However, there was significance difference in the prevalence of hookworm between the two schools; Nkpor (60.9%) and Mgboudohia (15.5%). In all, statistical significance difference ($p=.05$) was observed in the prevalence of *T.*

trichiura, hookworms and mixed infection (*A. lumbricoides* and *T. trichiura*) between schools in the study area (Table 1).

Parasite Intensity: The intensity of parasites was expressed as the number of eggs of specific parasite per gram of faeces. Among pupils of community primary school Mgbodohia, the intensity was 216.8, 188.7, 172.2 and 8.6 for *A. lumbricoides*, *T. trichiura*, Hookworm and Mixed infection (*A. lumbricoides* and *T. trichiura*) respectively (Table 3). For community primary school, Nkpor, the intensity included 276.8, 26.8, 201.8 for *A. lumbricoides*, *T. trichiura* and Hookworm respectively (Table 2). There was no statistically significance difference in the intensity of *A. lumbricoides* and Hookworm in pupils of both schools. However, a significance difference ($p = .05$) existed in the intensity of *T. trichiura* and mixed infection (*A. lumbricoides*, *T. trichiura*) among pupils of both schools (Table 4). The intensity of *A. lumbricoides* was higher in children in the age group of 5-10 years and lower in children above 17 years. Similar trend was observed in the intensity of hookworm and *T. trichiura* (Table 2).




Fig. 1: Overall prevalence of Soil Transmitted Helminthes (STH) among primary school pupils in Akpor and Mgbodohia

Table 1 : Prevalence of helminth parasite in relation to sex, age and school (n= 107)

Sex	No. examined	Species of helminthes				Total infected (%)
		<i>A.lumbricoides</i> No. infected (%)	<i>T. trichiura</i> No. infected(%)	Hookworms No. infected(%)	Mixed Infection (<i>A. Lumbricoide</i> + <i>T. trichiura</i>)	
Male	56	26 (46.4)	6(10.7)	11(19.6)	0 (0)	43(53.1)
Female	51	23(45.1)	5(9.8)	8 (15.7)	2(3.9)	38(47.0)
Age (yr)						
5-10	39	25 (65.8)	5(13.2)	8(21.1)	0(0)	38(47.0)
11-16	35	22(62.9)	4(11.4)	9(25.7)	0(0)	35(43.2.)
>17	33	3(37.5)	2(25.0)	1(12.5.)	2(25.0)	8(9.9)
School						
CPS, Mgbuodohia	50	22(46.8))	4(39.1)	5(15.2)	2(6.1)	33(40.7)
CPS, Nkpor	57	27(56.3)	7(63.6)	14.60.9	0(0)	48 (55.3)
TOTAL	107	49(60.5)	11(13.6)	19(23.5)	2(2.5)	81(75.7)

P=.05

Table 2: Intensity(epg) of Soil Transmitted Helminthes among primary school pupils in Nkpor and Mgboudohia in relation to age

Species of helminthes				
Age (yr)	<i>A. lumbricoide</i>	<i>T. trichiura</i>	Hookworms	Mixed Infection (<i>A. Lumbricoide</i> + <i>T. trichiura</i>)
5-10	117.3	98.2	145.2	0
11-16	98.1	7.8	139.2	0
>17	31.1	1.8	89.6	8.6

Table 3 : Intensity(epg) of Soil Transmitted Helminthes among primary school pupils in Nkpor and Mgboudohia

School	Species of helminthes			
	<i>A. lumbricoide</i>	<i>T. trichiura</i>	Hookworms	Mixed Infection (<i>A. Lumbricoide</i> + <i>T. trichiura</i>)
CPS, Mgbuodohia	216.8	188.7	172.2	8.6
CPS, Nkpor	276.2	26.8	201.8	0
TOTAL	246.5	107.8	187	4.3

p=.05

4.0 Discussion

The occurrence of soil transmitted helminthes among primary school children in Mgbodohia and Nkpor communities was investigated. A statistically significant proportion (75.7%) of the population in the study area harboured at least a helminth parasite, an indication that the prevalence rate of these parasites is high in the study area and still remains a public health concern among children [13,27]. Similar trend has been reported among school age children in various parts of Nigeria [28-35]. However, the high prevalence rate recorded in this study is higher than the 59.2% recorded by [11] among school-age children in Ife, Osun state and the 30.3% reported by [36] among school children in Amaruru community, Imo State, Southeast Nigeria. It is also higher than the 24.6% observed by [34] among among rural Fulani children in Vom, Plateau State. The result is also higher than the 58.0% reported by [37] among primary school children in Bo district of Sierra Leone, the 51.5% reported by [38] among adults of Akonolinga health district of Cameroon and the 19.3% recorded by [39] in the middle-belt of Ghana. The result was however lower than the 90% prevalence rate recorded among high risk area in India by [40] and the 75.6% recorded among primary school children in another part of India by [41]. The high prevalence recorded in this study could be attributed to lack of toilet facilities and portable drinking water [42] in the area. The poor socio-economic status and occupation of the parents may also be responsible for the high prevalence of the parasites among these children [11,42]. Furthermore, the high prevalence rate could also be attributed to the fact that the studied primary schools are located in rural communities. The two communities

are located at the periphery of Port Harcourt metropolis and had no urban flavor. The communities lack social infrastructures such as safe drinking water, good road and electricity. Ojurongbe et al. reported that rural communities are characterized by high poverty rate, poor infrastructural development, poor sanitation and poor hygiene, factors that enhance the transmission of helminth parasites.

Most of the pupils (60.5%) harboured *Ascaris lumbricoide*. The prevalence rate of *A. lumbricoide* observed in this study is higher than the 21.8% and 13.1% reported by [13] in Ifewara and Ajobandele respectively, 4% in Edo [44], 18.8% in Adamawa [45], 1.5% in Ghana [39] and 18.8% in Cameroon [38]. The prevalence (60.5%) reported in this study is also comparatively higher than the 44.8%, 13.1% and 17.7% reported by [11], [13] and [37] respectively. The parasite also occurred in all age groups investigated confirming the findings of [29, 33]. These authors recorded the prevalence of *A. lumbricoide* in all age groups investigated.

The high prevalence rate (47%) among children of lower age group (5-10yrs) recorded in this study is contrary to the findings of [11,37]. The researchers recorded the lowest prevalence rate (31.1%) and 7.7% respectively among lower age groups below 10years. The difference in our results could be attributed to the differences in study area and the attitude of parents towards their children. Our study was conducted in rural communities where little children are allowed to crawl on the ground and play indiscriminately on the ground and sometime bare

footed. Moreso, they eat with unwashed or improperly washed hands. The lower prevalence among children above 16years old is in consonance with the report of other authors [11,33,38]. The lower prevalence among this age group could be as a result of the fact that as age increases, the children become more conscious of personal hygiene and avoid playing with soil and unhygienic places [11,46]. They have less contact with soil as they are old enough to control their playing habit and are more conscious of their personal hygiene [19]. Similar observation has been recorded by [28,32].

The prevalence of Hookworms (23.5%) in the present study is lower than the 94.2% recorded among Primary School Children in a Amaruru Community in Imo State by [19]. It is also lower than the 33.3% recorded by [25] among abattoir worker in Port Harcourt, Rivers State. The difference in prevalence could be attributed to the period when the studies were embarked upon. [19] conducted their research in the rainy season while this study was conducted at the onset of dry season (Late November-December). The transmission of hookworm is reportedly high in rainy season [27] as the eggs are commonly distributed by the rain increasing the chance of parasite-human contact [19]. Invariably, during dry season, the transmission rate of hookworm is retarded during dry season.

The lower prevalence of *T. trichiura* recorded in this study agreed with the report of [47], among children in rural communities of Edo state and malnourished school age children in peri-urban area of Ibadan respectively. Similar lower prevalence was recorded by [11] while [42] did not observe any *T. trichiuru* in

their respective studies. However, the 13.6% prevalence observed in this study is slightly lower down the 14.9% recorded by [11]. Although more males (53.1%) were infected than females (47.0%), there was no statistically significant ($P=0.05$) difference in STHs infection in relation to sex. This is contrary to the record of [19] who reported more infection in males than in females. No clear reason could be advanced for this trend. However, this result is an indication that both sexes are vulnerable to soil transmitted helminthic infections as they both have playing contact with soil and are exposed to the same environmental, socioeconomic and hygienic conditions.

Our study showed that the only and most common co-infection was *A. lumbricoide* + *T. trichiura*. This observation is in consonance with the report of [11, 49] and who recorded a combined infection of *A. lumbricoide* + *T. trichiura* among primary school children of Ondo State, Nigeria. This result is however contrary to the records of [50, 51]. The researchers recorded co-infection of *A. lumbricoide* + Hookworm as the most common combined infection in Nsukka, Enugu and among palm wine drinkers in Ibadan respectively.

In spite of the relatively high intensity of *A. lumbricoide* in children in the age group of 5-10years (145.2epg) and 276.2epg recorded among children in Community primary school, Nkpor and the overall intensity of 246.5epg recorded in both school, none of the records reaches the threshold for light infection (1-4,000epg) standard stipulated by [52]. Similarly, the overall intensity of

T.trichura observed in both schools (107.8epg) and that of 187epg recorded for hookworm in both schools were also below the threshold for light infection, 1,999epg and 1-1999epg for *T. trichura* and hookworm respectively[52].

5. Conclusions

The results of this investigation indicated significant prevalence of STHs in the study area, hence there is need for deliberate effort to formulate sustainable control program by the government, Non-Governmental Organizations and other agencies including the multinational companies in the area. Provision of toilet facilities, clean drinking water, orientation on personal hygiene and improved sanitary habit will enhance control measures.

References

REFERENCES

1. Chukwuma MC, Ekejindu IM, Agbakoba NR, Ezeagwuna DA, Anaghalu IC, Nwosu DC. The Prevalence and Risk Factors of Geohelminth Infections among Primary School Children in Ebenebe Town, Anambra State, Nigeria. *Middle-East J. Sci. Res* 2009., 4 No.3: 211- 215.
2. Hotez PJ, Molyneux DH, Fenwick A, Ottesen E, Sachs SE, Sachs JD Incorporating a rapid impact package for neglected tropical diseases with programs for HIV/AIDS, tuberculosis and malaria, *PLoS Medicine* 2007; 3: e102
3. World Health Organization Soil transmitted Helminth Infections. *Fact Sheet* 2007; No. 366. Geneva,
4. Hotez PJ, da Silva N, Brooker S, Bethony J. Soil Transmitted Helminth Infections: The Nature, Causes and Burden of the condition. Working Paper No. 3, Disease Control Priority Project. Bethesda, Maryland: Fogarty International Centre, National Institute; 2003.
5. Nigeria Centre for Disease Control. Soil-transmitted helminthes. <http://www.ncdc.gov.ng/diseases/info/S>, 2017. Last updated. 2017-01-14, 15:35:38
6. Rachel LP, Smith JL, Jasrasaria R, Broker SM. Global numbers of infection and disease burden of soil transmitted helminth infections in 2010. *Parasites & Vectors* 2007, 7:37
7. Murray CJL, Salomon JA, Mathers CD, Lopez AD. Summary measures of population health: concepts, ethics, measurement and applications. World Health Organization, Geneva, 2002.
8. Ngonjo T, Okoyo C, Andove J, Simiyu E, Lelo AE, Kabiru E, Kihara J, Mwandawiro C. Current Status of Soil-Transmitted Helminths among School Children in Kakamega County, Western Kenya. *J. Par. Res Vol.*2016 Article ID 7680124,2007 9 pages <http://dx.doi.org/10.1155/2016/7680124>
9. World Health Organization. Soil Transmitted Helminths. Geneva.2007. Last updated: 2017-01-14, 15:35:38

10. Crompton DWT, Savioli L. Intestinal parasitic infection and urbanization. *Bullard World Health Organization*. 1993; 711: 1-7.
11. Salawu SA, Ughele VA. Prevalence of soil-transmitted helminths among school- age children in Ife East Local Government Area, Osun State, Nigeria. *FUTA J. Res. Sc.* 2015;1: 139-151.
12. Crompton DWT. Prevalence of Ascariasis. In Crompton D.W.T., Neshein.M.C.and Pawlowski Z.S (eds). *Ascariasis and its prevention and control*. Taylor and Francis London.1989.
13. Asaolu SO, Ofoezie IE, Odemuyiwa PA, Sowemimo OA, Ogunniyi TAB. Effect of water supply and sanitation on the prevalence and intensity of *Ascaris lumbricoides* among pre-school age children in Ajebandele and Ifewara Osun State, Nigeria. *Trans. R. Soc. Trop. Med. and Hyg.* 2002; 96: 600-604
14. Stollzfus RJ, Albonico M, Chwaya HM, Savioli L, Tielsh J, Schulze K, Yip R. Haemoduant determination of hookworm- related blood loss and its role in Iron deficiency in African Children. *Amer. J. Trop. Med. & Hyg.* 1996; 55: 399-404.
15. Miguel E, Kremer M. Worms: identifying impacts on duration and health in the presence of treatment externalities. *Econometricca* 2004; 72(1): 159- 217.
16. World Health Organization. Report of the WHO informal consultation on schistosomiasis control. World Health Organization. Document 1998; No. WHO/CDS/CPC/SIP/99.2.
17. World Health Organization. Soil-Transmitted Helminthes Infection. Key facts, Geneva, 2018.
18. Ofoezie IE, Bolton P, Imevbore AMA, Christensen NO. Schistosomiasis and other helminth infections in irrigation schemes in Sokoto, Katsina and Kebbi State of Nigeria. *Nig. J. Par.* 1996; 17: 31-37.
19. Odinaka KK, Nwolisa EC, Mbanefo F, Iheakaram AC, Okolo S. Prevalence and Pattern of Soil-Transmitted Helminthic Infection among Primary School Children in a Rural Community in Imo State, Nigeria. *J. Trop. Med.* 2005; Article ID 349439, 4 pages
20. Ekpo UF, Odoemene SN, Mafiana CF, Sam-Wobo SO. Helminthiasis and hygiene conditions of schools in Ikenne, Ogun State, Nigeria. *PLoS Negl. Trop.Dis.* 2008; 2(1):146(1-6).
21. Odu NN, Okonko IO, Erhi O. Study of Neglected tropical diseases (NTDs): Gastro-Intestinal Helminthes among school children in Port

- Harcourt, Rivers State, Nigeria. *Rep. & Opin.* 2011;3(9).
<http://.sciencepub.net/report>
22. World Health Organization. Soil-Transmitted Helminthiasis: Eliminating Soil-Transmitted Helminthiasis as a Public Health Problem in Children: Progress Report 2001–2010 and Strategic Plan 2011–2020, World Health Organization, Geneva, Switzerland, 2012.
 23. Eludoyin OS, Wokocha CC, Ayolagha G. GIS Assessment of Land Use and Land Cover Changes in Obio/Akpor L.G.A., Rivers State, Nigeria. *Res. J. Env. & Ear. Sci.* 2011; 3(4): 307-313.
 24. National Population Commission. Nigeria Demographic and Health Survey. NPC, Abuja. 2006.
 25. Gboeloh, LB & Elele, K. Incidence of Gastrointestinal Parasites Among Workers in Major Abattoirs in Port Harcourt, Rivers State, Nigeria. *Int'l. J. Med., H. Biomed. Bioeng. & Pharm. Eng.* 2013; 7(11): 750-752.
 26. Cheesbrough, M. *District Laboratory Practice in Tropical Countries*, Second Edition, London: Cambridge University Press. 2005; Pp.198-199
 27. Nwosu ABC. The community ecology of soil-transmitted helminth infections of humans in a hyperendemic area of southern Nigeria. *An. Trop. Med. Paras.* 1981;75 (2): 197–203.
 28. Ukpai OM Ugwu CD. The prevalence of gastro-intestinal tract parasites in primary school children in kwuano Local Government Area of Abia State Nigeria,” *Nig.J. Paras.* 2003; 24: 129–136
 29. Egwunyenga OA, Ataikiru DP. Soil transmitted helminthiasis among school age children in Ethiope East L.G.A. Delta State, Nigeria. *Afr. J. Biotech.* 2005; 4 (9):938-941.
 30. Ugbomoiko US, Onajole AT, Edungbola LD. Prevalence and intensity of geo-helminths infection in Oba- Ile Community of Osun State, Nigeria. *Nig. J. Paras.* 2006; 27: 62-67.
 31. Adeoye GO, Osayemi CO, Oteniya O, Onyemekeihia SO. Epidemiological Studies of Intestinal Helminthes and Malaria among Children in Lagos, Nigeria. *Pak. J. Bio. Sci.* 2007; 10: 2208-2212.
 32. Obiukwu MO, Umeanaeto PU, Eneanya CI, Nwaorgu GO. Prevalence of gastro-intestinal helminths in school children in Mbaukwu, Anambra State, Nigeria, *Nig. J. Paras.*, 2008, 29(1):15–19.

33. Sowemimo OA, Asaolu SO. The current status of soil transmitted helminthiasis among pre-school and school children in Ile Ife Osun State, Nigeria. *J. Helm.* 2011; 85:234-238.
34. Okolo SN, John C. Nutritional status and intestinal parasitic infestation among rural Fulani children in Vom, Plateau State, *Nig. J. Paedr.* 2008; 33 (2): 47–55.
35. Aisien MSO, Adams MA, Wagbatsoma VA. Intestinal helminthiasis in an Ochocerciasis endemic community on ivermectin treatment. *Nig. J. Paras.* 2002, 23:153-158.
36. Ugbomoiko US, Ofoezie IE. Multiple infection diagnosis of intestinal helminthiasis in the assessment of health and environmental development projects in Nigeria. *Journal of Helminthology* 2007; 81: 227–23.
37. Bangural ET, Ngegba MP, Nyalley F. Prevalence and intensity of soil-transmitted helminths (SIHs) and schistosomes in primary schools in Bo District, Southern Sierra Leone. *Global Journal of Bio-science and Technology.* 2016; 59(1): 55-61.
38. Bopda J, Nana-Djeunga H, Tenaguem J, Kamtchum-Tatuene J, Gounoue-Kamkumo R, Assob-Nguedia C, Kamgno J. Prevalence and intensity of human soil transmitted helminth infections in the Akonolinga health district (Centre Region, Cameroon): Are adults hosts contributing in the persistence of transmission? *Parasite Epidemiology and Control*, 2016; 1(2):199-20.
39. Adu-Gyasi D, Asante KP, Frempong, MT, Gyasi DK, Iddrisu LF, Ankrah L, Dosoo D, Adeniji E, Agyei O, Gyaase S, Amenga-Etego S, Gyan B, Owusu-Agyei S. Epidemiology of soil transmitted Helminth infections in the middle-belt of Ghana. *Parasite Epidemiology and Control*, 2018;3(3): e00071.
40. Salma N, Azam S. Prevalence and distribution of soil transmitted helminth infections in India. *BMC Public Health*, 2007; 17:201. DOI 10.1186/s12889-017-4113-2.
41. Ganguly S, Barkataki S, Karmakar S, Sanga P, Boopathi K, Kanagasabai K, Kamaraj P, Chowdhury P, Sarkar R, Raj D, James L, Dutta S, Sehgal R, Jha P, Murhekar M. High prevalence of soil-transmitted helminth infections among primary school children, Uttar Pradesh, India, 2015. *Infectious Diseases of Poverty*, 2017; 6:139 DOI 10.1186/s40249-017- 0354-74.
42. Ugbomoiko US, Ofoezie IE. Multiple infection diagnosis of intestinal helminthiasis in the assessment of health and environmental development projects in Nigeria. *Journal of Helminthology*, 2007; 81: 227–231.

43. Ojurongbe O, Oyesiji K F, Ojo JA, Odewale G, Adefioye OA, Olowe AO, Opaleye OO, Bolaji OS, Ojurongbe TA. Soil transmitted helminth infections among primary school children in Ile-Ife Southwest, Nigeria: A cross sectional study. *International Research Journal of Medicine and Medical Sciences*, 2014; 2(1): 6-10.
44. Oguanya FC, Okugu GRA, Akhile AO, Eloka CCV, Okoro CJ, Okpe AC. Prevalence of soil transmitted helminth infections among public school pupils in Ekpoma, Edo State, Nigeria. *International Journal of Community Research*, 2012; 1 (1): 30-34.
45. Shitta K, Akogun OB. Intestinal helminthes among nomadic Fulanis in two localities of Adamawa State, North-East Nigeria. *Nig. J. Para*, 2017; 38(1): 67-72
46. Adanyi CS, Audu PA, Luka SA, Adanyi DN. The influence of types of toilet used and personal on the prevalence of helminthosis among primary school children in Zaria, Kaduna State. *Scholar Research Library Archives of Applied Science Research* 2011; 3(3):257-260.
47. Osazuwa F, Ayo OM, Imade P. A significant association between intestinal helminth infection and anaemia burden in children in rural communities of Edo state, Nigeria, *N. Amer. J. Med. Sci.* 2011; 3(1):30-34.
48. Adeyeba OA, Tijani BD. Intestinal helminthiasis among malnourished school age children in peri-urban area of Ibadan, Nigeria. *Afr. J. Clin. & Exp. Microb.* 2002; 3(1), 24-28.
49. Oyewole F, Ariyo F, Sanyaolu A, Oyinbo WA, Faweya T, Monye P, Ukpong M, Okoro C. Intestinal Helinthiasis and their control with Albendazole Among Primary School Children in Riverine Communities of Ondo State, Nigeria. *Southeast Asian Journal of Tropical Medicine and Public Health*, 2002;33: 214-218.
50. Onuaha EO, Ofozie IE. Influence of Education Background and Personal Hygiene on Soil Transmitted Helminthiasis in Nsukka Zone, Enugu State, Nigeria in Many Species One Planet Future: Proceeding of International Conference of Institute of Ecology and Environmental Studies, Obafemi Awolowo University, Ile-Ife, 2010; 3:68-74.
51. Alli JA, Okonko IO, Oyewo AJ, Kolade AF, Nwaze JC, Ogunjobi PN, Tonade AA, Dada V K (2011). Prevalence of intestinal parasites among palm wine drinkers in Ibadan Metropolis, Nigeria, *Researcher*, 2011; 3(11):11-17.

52. WHO. Helminth control in school-age children: a guide for managers of control programmes, 2002, Geneva.

UNDER REEER REVIEW