

2
3 **The Effect of Different Organic Nutrients on**
4 **Some Quality Properties**
5 **of Popcorn (*Zea mays* L. *everta*)**

6
7 Haktan CIHANGIR¹, Abdullah OKTEM^{2*}

8 Corresponding author : aoktem@harran.edu.tr,

9 ¹Type Agriculture and Rural Development Support Institution, Mardin, Turkey

10 ²University of Harran, Faculty of Agriculture, Department of Field Crops, Sanliurfa, Turkey

11
12
13
14
15
16 **ABSTRACT**

17
Aims: Study was aimed to find out the effect of conventional and fifteen different organic nutrients (peat, compost, cattle manure, chicken manure, horse manure, sheep manure, pigeon manure, vermicompost, seaweed + cattle manure, compost + humic acid, cattle manure + humic acid, chicken manure + humic acid, horse manure + humic acid, sheep manure + humic acid, peat + humic acid) on some quality properties of popcorn.

Study design: Trial was designed in complete randomized block design with three replications. Ant-Cin-98 popcorn cultivar was used in the study. Each parcel comprised 4 lines. The planting was made into a depth of 5-6 cm along the lines 5 meters long with a row spacing of 70 cm and intra row of 20 cm.

Place and Duration of Study: This study was conducted in Diyarbakır - Cermik conditions of Turkey between 2010 and 2011.

Methodology: The effect of conventional and fifteen different organic materials to some quality parameters of popcorn such as cob ratio, 1000-kernel weights, test weight, popping volume and number of unpopped kernel were evaluated in the study. Physical and chemical properties of the trial area were determined by taking soil sample from a depth of 0-30 cm on the trial area before planting.

Results: According to the investigated results, the highest and the lowest values were ranked between 19.98% (peat + humic acid) and 17.26% (vermicompost) for cob ratio, 138.65 g (seaweed + cattle manure) and 122.48 g (chicken manure) for 1000-kernel weight, 81.29 kg hl⁻¹ (horse manure + humic acid) and 75.62 kg hl⁻¹ (vermicompost) for test weight, 19.71 cm³ g⁻¹ (peat) and 17.17 cm³ g⁻¹ (sheep manure + humic acid), for popping volume 5.92% (peat) and 3.65% (horse manure + humic acid) for number of unpopped kernel.

Conclusion: Higher values obtained from organic nutrient sources than conversational application in all tested quality parameters. The implementation of organic fertilizers together with humic acid in popcorn produced better results in comparison to alone implementation of organic fertilizers. Also it was determined that use of natural enemies of *Trichogramma* spp against corn borer can be possible without any chemicals.

18
19 **Keywords:** Organic nutrients, popcorn, popping volume, test weight

20 **1. INTRODUCTION**

21

22 The corn is used in human nutrition directly and indirectly in addition to the use as industrial
23 raw material and animal feed in the world. Although dent corn (*Zea mays* L. *indentata*)
24 varieties comprise the vast majority of corns grown both in world and in Turkey, no statistics
25 so far related to the cultivation area, manufacture and consumption amount of popcorn (*Zea*
26 *mays* L. *everta*) is available in Turkey. It is reported that planting is made around the
27 provinces of Adana, Canakkale, Adapazari, Antalya, Isparta and Burdur, in Aegean and
28 Mediterranean Regions of Turkey [1]. Consumption of popcorn is increasing every passing
29 day in Turkey.

30 Popcorn is also directly used in human nutrition. The sub type of corn having grains popped
31 when heated is popcorn. It generates pressure inside the grain through expansion when the
32 humidity in the endosperm is heated because its grain is hard, its hull is thick and
33 impermeable. At the same time the starch in the endosperm transforms with the effect of
34 heat. The hull can't resist this pressure and bursts by splitting suddenly. The volume of the
35 grains burst expands and they are eaten by salting or adding oil. Its consumption rises also
36 in Turkey because of low cost and easy to prepare with popping machines, in pans or pots.
37 Popcorn is commonly consumed while watching cinema and soccer matches and television
38 during winter months. Additionally, it is preferred much by children [2].

39 Nowadays environmental pollution has reached a significant level as a result of the use of
40 synthetic and chemical inputs in excessive amounts due to the production increase within
41 conventional agriculture system. Organic fertilizers/matters, which are essential inputs of
42 organic agriculture systems that have become widespread in parallel with the interest of
43 people in organic products, are made available for producers under a great variety of names
44 and contents in the market. It is necessary to utilize these matters in various ways to prevent
45 environmental pollution caused by wastes, and to enhance organic matter level of our soil.

46 Different results have been obtained in studies regarding the subject of the study. Anac and
47 Okur [3] reported that application of Biofarm (certified organic fertilizer) and farm fertilizer as
48 organic fertilizer (uncertified) to soil has led to significant increase in dry weight, mineral
49 content and efficiency of corn compared to control. Yazici and Kaynak [4] reported that
50 seaweed increases yield and quality in organic farming, regulate the growth of plants,
51 increase resistance to pests and diseases, improves the structure of the soil. Seker and
52 Ersoy [5] investigated the effects of different doses of compost, cattle manure, chicken
53 manure and leonardit on the soil properties and the development of corn (*Zea mays*
54 L.). They found as a result of the research that type and dose of used organic fertilizer
55 affects soil properties and the corn's growth.

56 Shafiq et al. [6] stated that chicken manure affected positively some parameters such as
57 plant height, seed number, 1000-kernel weight and grain yield values. Selcuk and Tufenkci
58 [7] reported that increasing humic acid application provided significant increase in number of
59 kernel per ear, ear length, plant height and 1000-kernel weight. Cengiz et al. [8] emphasized
60 that effect of organic fertilizers on yield and quality of the corn plant were similar with
61 chemical fertilizers.

62 In this study, it was aimed determining the effect of some organic nutritional sources on
63 some quality properties of popcorn.
64

65

66 2. MATERIAL AND METHODS

67 2.1. Material

68 Ant-Cin-98 popcorn variety was used in the experiment. Organic nutrient sources were used
69 in the study (Table 1). Amount of total pure nitrogen both conventional and organic growing
70 were 17 kg da⁻¹ based on the regulation, principles and applications of organic agriculture in
71 Turkey [9]. According to nitrogen content of organic material, maximum pure nitrogen
72 amount (17 kg da⁻¹) was calculated for organic applications (Table 1). For conventional
73 applications total of 17 kg da⁻¹ nitrogen, 8 kg da⁻¹ phosphor and potassium (15-15-15 NPK
74 as bottom fertilizer and urea as top fertilizer) were given as pure per decare. Nitrogen
75 content of nutritional sources used in the study and the amount of fertilizer thrown per
76 decare were given in Table 1.

77 **Table 1. The nitrogen content of organic nutrient sources and applied amount**

	Nutritional Sources	N content (%)	The amount of applied (kg da ⁻¹)
1	Conventional manure (urea)	46	36.96 kg da ⁻¹
2	Peat	1.2	1416 kg da ⁻¹
3	Compost	2.5	680 kg da ⁻¹
4	Cattle manure	3.5	486 kg da ⁻¹
5	Chicken manure	3.0	567 kg da ⁻¹
6	Horse manure	2.0	850 kg da ⁻¹
7	Sheep manure	2.0	850 kg da ⁻¹
8	Pigeon manure	6.0	283 kg da ⁻¹
9	Seaweed + Cattle manure	2.0 + 3.5	51.5 kg da ⁻¹ + 457 kg da ⁻¹
10	Vermicompost	1.5	1133 kg da ⁻¹ + 140 g da ⁻¹
11	Compost + Humic acid	2.5	680 kg da ⁻¹ + 140 g da ⁻¹
12	Cattle manure + Humic acid	3.5	486 kg da ⁻¹ + 140 g da ⁻¹
13	Chicken manure + Humic acid	3.0	567 kg da ⁻¹ + 140 g da ⁻¹
14	Sheep manure + Humic acid	2.0	850 kg da ⁻¹ + 140 g da ⁻¹
15	Peat + Humic acid	1.2	1416 kg da ⁻¹ + 140 g da ⁻¹
16	Horse manure + Humic acid	2.0	850 kg da ⁻¹ + 140 g da ⁻¹

78

79 2.1.1. Climatic characteristics of the research area

80 Experiment was conducted in Diyarbakır province Cermik district under second crop
81 conditions. Climate values of 2010 and 2011 in which the research was conducted with long
82 years were given in Table 2. The average highest temperature (32.7°C) was observed in
83 July, the lowest temperature (12.0°C) in November between June-November months in
84 2010, and in 2011 the average highest temperature (31.5°C) in July, the lowest temperature
85 (6.6°C) in November in accordance with the data received from Diyarbakır Regional
86 Directorate of Meteorology. The highest value (61.8%) in terms of relative humidity occurred
87 in October 2010, and the lowest value (22.3%) in August 2011. The water need of plants
88 was met through irrigation during the growing period.

89

90 **2.1.2. Soil characteristics of the research area**

91 Total salt content was found to be 0.03%, nitrogen content 1.19%, lime rate 9.8%,
92 phosphorus amount 2.75 kg da⁻¹, potassium amount 82.05 kg da⁻¹ and soil pH 7.4 in the soil
93 sample taken from 30 cm soil depth in the place where trial was established in Diyarbakır
94 province Cermik district.

95 **Table 2. Rainfall, temperature and relative humidity values for 2010, 2011 and long**
96 **years in Diyarbakır province**

Months	Min. Temp. (°C)	Max. Temp. (°C)	Average Temp. (°C)	Rainfall (mm)	Relative humidity (%)	
June	2010	14.9	40.8	27.2	8.0	47.6
	2011	13.2	37.9	26.3	14.6	33.9
	Long years	16.9	33.7	26.3	7.2	36.0
July	2010	18.0	44.0	32.7	0.0	34.3
	2011	18.4	45.0	31.5	0.2	22.6
	Long years	21.7	38.5	31.2	0.7	27.0
August	2010	18.0	43.6	32.4	0.0	32.2
	2011	16.0	43.5	31.2	0.0	22.3
	Long years	21.0	38.1	30.3	0.3	27.0
September	2010	13.6	41.2	26.8	3.0	44.7
	2011	12.8	38.1	25.6	1.9	28.5
	Long years	16.0	33.1	24.8	2.6	31.0
October	2010	7.3	30.0	17.6	49.2	61.8
	2011	3.0	32.8	17.4	57.4	52.5
	Long years	10.1	25.3	17.2	30.8	48.0
November	2010	1.0	26.1	12.0	0.0	57.4
	2011	-4.7	19.9	6.6	104.0	61.1
	Long years	3.6	15.9	9.3	54.6	68.0

97 Resource: Anonymous [10].

98 **2.2. Method**

99 Before the starting of experiment, the trial area was planted with wheat in 2008 and 2009 for
100 making the area suitable for organic farming in which the trial would be established, and
101 wheat was cultivated and harvested without application of any chemical fertilizer and
102 agricultural pesticide. Physical and chemical properties of the trial area were determined by
103 taking soil sample from a depth of 0-30 cm on the trial area before planting.

104 The soil was made ready for planting by processing with goble-disc and then with disc-
105 harrow prior to planting. The trial was established with three replicates according to
106 randomized complete block experimental design. Each parcel comprised 4 lines. The
107 planting was made by hand into a depth of 5-6 cm along the lines 5 meters long with a row
108 spacing of 70 cm and intra row of 20 cm in 15-30 June. Most of the nutritional organic
109 sources were applied with planting. Also some part of seaweed manure was applied before
110 planting and the rest as foliar fertilizer in three times. An equal amount of water was given to
111 the parcels with sprinkler irrigation after planting for germination, and furrow irrigation
112 throughout the growing period due to lack of moisture in sufficient levels for output. A space
113 of 2 meters was left between parcels to hinder water passage between parcels and the
114 parcels were surrounded with berm.

115 Cultural measures (tractor and hand hoeing) were implemented to combat the weed.
 116 Chemical pesticides were not used in the search, *Trichogramma sp.* predator that was
 117 obtained from Adana Agricultural Research Institute Biological Control Unit was used within
 118 scope of biological control against Corn Cob Worm which leads to great productivity losses
 119 for corn plants. Values were taken from two rows in the middle after discarding 0.5 m from
 120 both beginnings and one rows situated at either sides of the parcel as edge effect during the
 121 harvest between 20-25 September.

122 An analysis of variance (ANOVA) was performed by using Totemstat-C software package
 123 for the tested characteristics to evaluate statistical differences between used organic
 124 nutrients. Means were compared by the Duncan's multiple range test ($P \leq 0.05$) [11].

125 3. RESULTS AND DISCUSSION

126 3.1. Cob Ratio (%)

128 Considering 2010-2011 year averages according to Table 3, cob ratio varied between
 129 percent 17.26-19.98 in different nutrient applications. The highest cob ratio value was
 130 determined to be 19.98% in peat + humic acid. The lowest cob ratio was obtained as
 131 17.26% from vermicompost application along combined averages in the trial.

132 **Table 3. Cob ratio (%) values found in popcorn grown using different nutritional**
 133 **sources and the groups formed according to Duncan Test**

Nutritional Sources	2010^{ns}	2011^{ns}	Average[‡]
Conventional fertilizer	15.74	20.24	17.99 AB
Peat	14.44	22.40	18.42 AB
Compost	14.29	20.63	17.46 AB
Cattle manure	15.25	22.06	18.65 AB
Chicken manure	15.41	22.41	18.91 AB
Horse manure	15.37	20.01	17.69 AB
Sheep manure	15.46	19.39	17.43 AB
Pigeon manure	15.93	21.99	18.96 AB
Seaweed + cattle manure	15.73	24.16	19.95 A*
Vermicompost	15.30	19.21	17.26 B
Compost + humic acid	14.67	20.87	17.77 AB
Cattle manure + humic acid	15.50	22.92	19.21 A
Chicken manure+ humic acid	15.36	23.43	19.40 A
Sheep manure. + humic acid	15.71	22.14	18.92 AB
Peat + humic acid	15.51	24.45	19.98 A
Horse manure + humic acid	15.64	21.40	18.52 AB
Average	15.33 B	21.73 A	
LSD	Year: 2.014		
	2010-2011 Average nutritional sources: 2.539		

134 *There is no significant difference at 0.05 level according to Duncan Test among the
 135 averages falling within same letter group.
 136 †: $P \leq 0.01$, ‡: $P \leq 0.05$ ns: No significant

137

138

139

140 **3.2. 1000-Kernel Weight (g)**

141 Considering 2010-2011 year averages according to Table 4, 1000-kernel weights varied
 142 between 122.48-138.65 g in different nutrient applications. While the highest 1000-kernel
 143 weight value was determined to be 138.65 g in seaweed manure + cattle manure, and
 144 afterwards this was followed by horse manure + humic acid (137.41) with sheep manure +
 145 humic acid (137.11), respectively. In the meantime, conventional fertilizer application ranked
 146 sixth among the applications with a 1000-kernel weight value of 132.41 g. The lowest 1000-
 147 kernel weight was obtained as 122.48 g from chicken manure application along combined
 148 averages in the trial. In terms of 1000-kernel weight, it can be said that the abundance of all
 149 applications in second year compared to the first year resulted from both climate and
 150 environmental conditions and positive effect of nutritional sources.

151 The effect of plant nutritional sources on 1000-kernel weight in respect of corn plant was
 152 given by obtaining different results in different studies. Prasanna et al. [12] reported that they
 153 received the highest 1000-kernel weight from vermicompost in respect of corn plant, Shafiq
 154 et al. [6] stated that chemical fertilizer yielded 1000-kernel weight higher than organic
 155 fertilizers (chicken manure, farmyard manure, bio-manure).

156 **Table 4. 1000-kernel weight (g) values found in popcorn grown using different**
 157 **nutritional sources and the groups formed according to Duncan Test**

Nutritional Sources	2010^T	2011^T	Average^T
Conventional fertilizer	123.38 a*	141.43 de	132.41 A-E
Peat	113.97 a-d	131.68 e	122.82 DE
Compost	105.78 cd	141.89 cde	123.84 CDE
Cattle manure	115.6 abc	144.02 bcd	129.81 A-E
Chicken manure	105.22 cd	139.73 de	122.48 E
Horse manure	117.10 ab	148.31 a-d	132.71 A-E
Sheep manure	118.27 ab	150.04 a-d	134.16 ABC
Pigeon manure	120.25 a	148.00 a-d	134.13 A-D
Seaweed + cattle manure	121.47 a	155.84 a	138.65 A
Vermicompost	104.40 d	148.73 a-d	126.56 B-E
Compost + humic acid	107.73 bcd	146.87 a-d	127.30 B-E
Cattle manure + humic acid	117.72 ab	148.64 a-d	133.18 A-E
Chicken manure+ humic acid	108.85 bcd	144.84 a-d	126.85 B-E
Sheep manure. + humic acid	120.30 a	153.92 ab	137.11 AB
Peat + humic acid	116.38 abc	143.64 cde	130.01 A-E
Horse manure + humic acid	121.00 a	153.82 abc	137.41 AB
Average	114.84 B	146.34 A	
	Year: 4.826		
LSD	2010 Nutritional sources: 10.021		
	2011 Nutritional sources: 10.175		
	2010-2011 Average nutritional sources: 9.841		

158 **There is no significant difference at 0.05 level according to Duncan Test among the*
 159 *averages falling within same letter group.*
 160 *†: P ≤ .01, ‡: P ≤ .05 ns: No significant*

161 Various results were obtained in different studies carried out related to 1000-kernel weight in
 162 popcorn. Idikut et al. [13] 114.9-122.9 g; Ertas et al. [14] 54.8-64.6 g; Gokmen et al. [15]
 163 129.0-213.0 g; Ozkan [16] 127.0-133.0 g; Tekkanat and Soyly [17] 114.68-175.93 g; Oktem
 164 et al. [18] 291.0-342.0; Ozkaynak and Samancı [19] reported 1000-kernel weight varying
 165 between 86.0-140.0 g in lines, 83.0-115.0 g in hybrids.

166 **3.3. Test weight (kg hectoliter⁻¹)**

167 Average values of the proportion of test weight determined in different nutritional sources in
 168 popcorn grown organically between 2010 and 2011 and the groups formed according to
 169 Duncan multi comparison test were given in Table 5.

170 In the trial, the difference among fertilizer applications was found to be statistically
 171 significant. Considering 2010-2011 year averages, test weight ranged from 75.62 kg
 172 hectoliter⁻¹ (hl⁻¹) to 81.29 kg hl⁻¹ in different nutrient applications. When examined the Table
 173 5, the highest test weight value was 81.29 kg hl⁻¹ in horse manure + humic acid application,
 174 and afterwards respectively, peat + humic acid (80.58 kg hl⁻¹) and sheep manure + humic
 175 acid (80.56 kg hl⁻¹) applications. Meanwhile, the lowest test weight was obtained as 75.62 kg
 176 hl⁻¹ from vermicompost. The difference of nutrient elements in the structure of organic and
 177 conventional nutritional sources at the end of the study, was seen affecting these fertilizer
 178 sources at different levels.

179 As a result, the highest test weight values were obtained from using organic fertilizers with
 180 humic acid which is a growth regulator. The studies have shown that humic acid effects dry
 181 weight in plant. Some researchers reported that fresh and dry weights increased significantly
 182 ($P \leq 0.05$) with treated humic acid at different levels compared to control [6, 20]. Asli and
 183 Neuman [21] reported that the humic acids reduce the dry weight of corn.

184 **Table 5. Test weight (kg hl⁻¹) values found in popcorn grown using different nutritional**
 185 **sources and the groups formed according to Duncan Test**

Nutritional Sources	2010 [‡]	2011 [†]	Average [†]
Conventional fertilizer	80.57 ab	76.55 e	78.56 AB
Peat	78.70 ab	80.62 abc	79.66 A
Compost	75.68 bc	79.47 bcd	77.58 AB
Cattle manure	76.43 abc	80.60 abc	78.52 AB
Chicken manure	75.98 abc	79.62 bcd	77.80 AB
Horse manure	78.13 abc	81.32 abc	79.73 A
Sheep manure	76.10 abc	79.82 bcd	77.96 AB
Pigeon manure	79.28 ab	80.63 bc	79.96 A
Seaweed + cattle manure	79.38 ab	79.25 cd	79.32 A
Vermicompost	73.47 c	77.77 de	75.62 B
Compost + humic acid	75.95 bc	82.78 a	79.37 A
Cattle manure + humic acid	76.87 abc	81.42 ab	79.14 AB
Chicken manure+ humic acid	80.95 a	79.80 bcd	80.38 A
Sheep manure. + humic acid	80.23 ab	80.88 abc	80.56 A
Peat + humic acid	79.75 ab	81.40 abc	80.58 A
Horse manure + humic acid	80.88 ab	81.70 ab	81.29 A
Average	78.02 B	80.23 A	
	Year: 0.463		
LSD	2010 Nutritional sources: 4.445		
	2011 Nutritional sources: 1.955		
	2010-2011 Average nutritional sources: 3.346		

186 **There is no significant difference at 0.05 level according to Duncan Test among the*
 187 *averages falling within same letter group.*
 188 *†: P ≤ .01, ‡: P ≤ .05 ns: No significant*

189

190 3.4. Popping Volume (cm³ g⁻¹)

191 Considering 2010-2011 year averages according to Table 6, popping volume varied between
 192 17.17 cm³ g⁻¹ and 19.71 cm³ g⁻¹ in different nutrient applications. When examined the Table
 193 6, the highest popping volume value was 19.71 cm³ g⁻¹ in peat application, and afterwards
 194 respectively, vermicompost (19.41 cm³ g⁻¹) and pigeon manure (18.98 cm³ g⁻¹) applications.
 195 Meanwhile, the lowest popping volume was obtained as 17.17 cm³ g⁻¹ from sheep manure +
 196 humic acid. The difference of nutrient elements in the structure of organic and conventional
 197 nutritional sources was seen clearly in test weight values.

198 Besides, even though no study has been carried out about popping volume in organic
 199 popcorn, different study results obtained related to popping volume as 19.79-22.92 cm³ g⁻¹
 200 [14] ; 19.67-25.33 cm³ g⁻¹ [19]; 18.50-35.25 cm³ g⁻¹ [17]; 21.0-27.5 cm³ g⁻¹ [22]; 28.1-28.7 cm³
 201 g⁻¹ [16] have a nature supporting our research results.

202 **Table 6. Popping volume (cm³/g) values found in popcorn grown using different**
 203 **nutritional sources and the groups formed according to Duncan Test**

Nutritional Sources	2010^{ns}	2011^{ns}	Average[‡]
Conventional fertilizer	18.95	17.42	18.18 AB
Peat	20.41	19.00	19.71 A*
Compost	19.93	17.75	18.84 AB
Cattle manure	19.66	19.04	19.35 AB
Chicken manure	18.99	17.68	18.33 AB
Horse manure	19.02	18.43	18.72 AB
Sheep manure	19.00	17.64	18.32 AB
Pigeon manure	19.20	18.76	18.98 AB
Seaweed + cattle manure	18.54	16.65	17.60 AB
Vermicompost	20.56	18.26	19.41 A
Compost + humic acid	18.62	17.15	17.89 AB
Cattle manure + humic acid	18.27	18.09	18.18 AB
Chicken manure+ humic acid	18.95	17.65	18.30 AB
Sheep manure. + humic acid	17.40	16.93	17.17 B
Peat + humic acid	19.42	16.12	17.77 AB
Horse manure + humic acid	17.75	17.61	17.68 AB
Average	19.04 A	17.76 B	
LSD	Year: 0.467		
	2010-2011 Average nutritional sources: 2.011		

204 *There is no significant difference at 0.05 level according to Duncan Test among the
 205 averages falling within same letter group.

206 †: P ≤ .01, ‡: P ≤ .05 ns: No significant

207 3.5. Number of Unpopped kernel (%)

208 In the experiment, the difference among fertilizer applications was found to be statistically
 209 significant. Average values of number of unpopped kernel determined in different nutritional
 210 sources in popcorn grown organically between 2010 and 2011 and the groups formed
 211 according to Duncan multi comparison test were given in Table 7. Considering 2010-2011
 212 year averages, number of unpopped kernel ranged from 3.65% to 5.92% in different nutrient
 213 applications. When examined the Table 7, the highest number of unpopped kernel value was
 214 5.92% in peat application, and afterwards respectively, chicken manure (5.63%) and
 215 compost (5.16%) applications. Meanwhile, the lowest number of unpopped kernel was
 216 obtained as 3.65 % from horse manure + humic acid.

* Tel.: +90 414 3183686; fax: +90 414 3183686.

E-mail address: aoktem@harran.edu.tr.

217 Besides, even though no study has been carried out about number of unpopped kernel in
 218 organic popcorn, different study results obtained related to non-popped grain rates as 12.43-
 219 16.91% [14], 3.49-12.19% in lines and 6.33-9.94% in hybrids [19]; 2.42-9.90% [17]; 2.77-
 220 3.48% [16], have a nature supporting our research results. Many researchers [14, 22] have
 221 found significant differences in non-popped grain rate which is among major quality
 222 parameters of popcorn, and they reported that the impact of varieties had a largest share in
 223 this situation.

224 **Table 7. Number of unpopped kernel (%) values determined in popcorn grown using**
 225 **different nutritional sources and the groups formed according to Duncan Test**

Nutritional Sources	2010 [†]	2011 [†]	Average [†]
Conventional fertilizer	6.48 abc	3.51 b-e	5.00 A-D
Peat	7.05 a*	4.79 a	5.92 A
Compost	6.72 ab	3.59 b-e	5.16 ABC
Cattle manure	4.86 def	3.84 bcd	4.35 CD
Chicken manure	6.97 a	4.28 ab	5.63 AB
Horse manure	5.27 b-f	3.88 abc	4.58 BD
Sheep manure	4.74 ef	3.18 cde	3.96 CD
Pigeon manure	5.47 a-f	3.45 b-e	4.46 BCD
Seaweed + cattle manure	4.88 c-f	2.67 e	3.78 D
Vermicompost	5.91 a-e	3.63 bcd	4.77 A-D
Compost + humic acid	4.41 f	3.46 b-e	3.93 CD
Cattle manure + humic acid	6.33 a-d	3.23 cde	4.78 A-D
Chicken manure+ humic acid	4.63 ef	3.51 b-e	4.07 CD
Sheep manure. + humic acid	4.67 ef	2.81 de	3.74 D
Peat + humic acid	4.65 ef	3.10 cde	3.88 CD
Horse manure + humic acid	4.50 ef	2.79 de	3.65 D
Average	5.47 A	3.48 B	
	Year: 0.467		
	2010 Nutritional sources: 1.393		
	2011 Nutritional sources: 0.889		
	2010-2011 Av. Nutritional sources: 2.011		

LSD

226 **There is no significant difference at 0.05 level according to Duncan Test among the*
 227 *averages falling within same letter group.*

228 *†: P ≤ .01, ‡: P ≤ .05 ns: No significant*

229

230

231 **4. CONCLUSION**

232 It has been determined with this study that organic popcorn production can be made also by
 233 using different nutritional sources under Diyarbakır ecological conditions. Higher values
 234 obtained from organic nutrient sources than conversional application in all tested quality
 235 parameters. Furthermore, it was observed that the implementation of organic fertilizers
 236 together with humic acid in popcorn produced better results in comparison to alone
 237 implementation of organic fertilizers. It has been proved that corn production can be made
 238 without the use of chemical pesticides in the trial. *Trichogramma sp.* beneficial insects can
 239 be introduced to local farmers and its use may be encouraged on corn planted areas.
 240

242 **REFERENCES**

- 243 1. Kun E. Warm Climate Cereals, Ankara University Printing House, Ankara, 1997.
- 244 2. Oktem A, Ulger A.C, Kirtok Y. Effects of different nitrogen doses and distances of row
245 spacing on grain yield and some agronomic characteristics of popcorn. Cukurova
246 University Journal of Agricultural Faculty, 2001;16(2):83-92.
- 247 3. Anac D, Okur B. Improving Soil Fertility in Natural Ways. Ecological (Organic, Biological)
248 Agricultural Organisation Society (ETO), p.37-73, 1996, Izmir.
- 249 4. Yazici K, Kaynak L. Possibilities of using seaweed in organic farming. Turkey 2nd
250 Ecological Symposium, 14-16 November 2001, Antalya, p. 344-352.
- 251 5. Seker C, Ersoy, I. Soil characteristics of different organic fertilizers and leonardit and their
252 effects on development of corn (*Zea mays* L.). Selcuk University Journal of
253 Agricultural Faculty, 2005;19(35):46-50.
- 254 6. Shafiq R, Bukhsh M.A, Ishaque M. Comparative performance and profitability of two corn
255 hybrids with organic and inorganic fertilizers. Pakistan Journal of Agricultural Sciences
256 2008;45(3):8-12.
- 257 7. Selcuk R, Tufenkci S. Effects of ascending doses of zinc and humic acid applications on
258 yield and nutrient content of corn. Yuzuncu Yil University, Institute of Science,
259 Master's Thesis, Van 2009.
- 260 8. Cengiz R, Yanikoglu S, Sezer MC. Effects of synthetic and organic fertilizers on yield and
261 quality of corn (*Zea mays* L.). Ministry of Agriculture and Rural Affairs, Ankara,
262 2010;213-220.
- 263 9. Anonymous. Official gazette, <http://mevzuat.basbakanlik.gov.tr/>, Official Gazette Number:
264 27676, Access date: 18.08.2010. 2010.
- 265 10. Anonymous, Monthly climatic report. Meteorological station. Sanliurfa. 2011.
- 266 11. Acikgoz N, Ilker E, Gokcol A. Assessment of Biological Research on the Computer.
267 TOTEM, 2004;2, Izmir.
- 268 12. Prasanna K, Halepyati A. S, Desai B.K, Pujari B.T. Effect of integrated nutrient
269 management on the productivity and nutrient uptake by maize (*Zea mays* L.).
270 Karnataka Journal of Agricultural Sciences, 2007;20(4):833-834.
- 271 13. Idikut L, Yilmaz A, Yururdurmaz C, Colkesen M. Determination of morphological and
272 agronomical properties of local popcorn genotypes. BİBAD Research Journal of
273 Biological Sciences, 2012;5(2):63-69.
- 274 14. Ertas N, Soylu S, Bilgici N. A research on determination of physical properties of corn
275 relations of quality parameters. Turkey 10. Food Congress, Erzurum, 21-23 May,
276 2008.
- 277 15. Gokmen S, Sakin M.A, Yildirim A., Yield, yield components and quality specifications of
278 some popcorn varieties grown in Tokat-Kazova conditions. Turkey VII. Field Crops
279 Congress, 2007;1:330-333, 25-27 June 2007, Erzurum.
- 280 16. Ozkan A. Effects of different nitrogen levels on grain yields, agricultural properties and
281 some quality parameters of two popcorn (*Zea mays everta* Sturt.) cultivars under
282 Cukurova conditions. Department of Field Crops Institute of Naturel and Applied
283 Sciences University of Cukurova, Phd Thesis, Adana,125p. 2007.
- 284 17. Tekkanat A, Soylu S. Determination of important quality characters and grain yield in
285 popcorn cultivars. Selcuk University Journal of the Faculty of Agriculture,
286 2005;19(37):41-50.
- 287 18. Oktem A., Ulger A.C, Coskun Y. Effects of some weed control methods on grain yield
288 and yield components of corn (*Zea mays* L.) under Harran plain conditions. Journal of
289 University Agricultural Faculty, 2004;8(1):51-57.

- 290 19. Ozkaynak E, Samanci B. Comparison of popcorn (*Zea mays everta* Sturt.) lines and their
291 testcrosses for yield and yield-related traits. *Akdeniz University Journal of the Faculty*
292 *of Agriculture*, 2003;16(1):35-42.
- 293 20. Dogru A, Darcin E.S, Tutar A, Dizman M, Koc Y. Effects of potassium humate to corn
294 (*Zea mays* L.) plant growth. *Sakarya University The Journal of Arts and Science*,
295 2012;1:25-35.
- 296 21. Asli S, Neumann P.M. Rhizosphere humic acid interacts with root cell walls to reduce
297 hydraulic conductivity and plant development. *Plant and Soil*, 2010;336:313-322.
- 298 22. Koc N, Ekiz H, Soysal M, Pamukcu M, Erdal S, Toros A. Popcorn population breeding.
299 Turkey 5. Field Crops Congress, Antalya 5-9 September 2005.
- 300 22. Gokmen S, Sakin M.A. Determination of yield, yield components and some quality
301 properties of popcorn. Turkey 3. Field Crops Congress, s.253-257, 17-21 September
302 2001, Tekirdag.