

# Diversity of ant species (Hymenoptera: Formicidae) Nesting beneath different trees in Periyanaickenpalyam, Coimbatore District, TamilNadu

## ABSTRACT

Ants (Formicidae: Hymenoptera) are one of the most successful organisms of the tropics and have survived many geological changes in the past and play crucial roles in ecosystem functioning. The present study examined ants nesting beneath different trees in Periyanaickenpalyam village, from November 2017 to February 2018. The study observed that 384 ants nests associated with 27 floral species at different locations in Periyanaickenpalyam were recorded. Based on floral ants nest association maximum of 18% ants nest in the *Prosopis juliflora* species followed by 15% of *Azadirachta indica*, *Ficus religiosa* contain 12%, *Mangifera indica* cover 9%, *Shorea robusta* (6%) and *Toona ciliata* equally contain (6%) and rest of floral species *Cocos nucifera*, *Ficus benghalensis*, *Santalum album*, *Gmelina arborea*, *Bambuseae*, *Carica papaya*, *Thespepsia populnea*, *Delonix*, *Tamarinds indica*, *Cassia fistula*, *Musa acuminata* each contain (3%) were observed. According to nest wise, fifty four nest below the *Prosopis juliflora* followed by fifty two nests in *Azadirachta indica*, (49) nests engaged in *ficus religiosa*, forty nests in *Mangifera indica* and twenty three nests beneath the *Shorea robusta*, *Ficus benghalensis* each and in *Toona ciliata*, *Bambuseae* both contain twenty two nests, rest of species contain less than 15 ants in *Gmelina arborea*, *Carica papaya*, *Delonix*, *Cocos nucifera*, *Santalum album*, *Musa acuminata*, *Tamarinds indica*, *Cassia fistula* and *Thespepsia populnea* trees were observed from the study. It can be concluded that this study provides clue information on ants nesting association with floral species.

*Key words:* [Ants diversity, Nests, Mutualism, *Azadirachta indica*].

## 1. INTRODUCTION

Ants are ubiquitous in distribution and occupy almost all terrestrial ecosystems. Ants are one of the ideal model organisms for measuring and monitoring biodiversity for many reasons. Ant belongs to a single large family Formicidae, largest of order Hymenoptera. It is represented by 26 extant subfamilies with 14,711 valid species and 428 valid genera [1] out of these, 152 species are listed by IUCN and 10 subfamilies were reported in India, which are represented by 100 genera with 828 species. Abundance and diversity of ants species is also greater neighbouring areas, for example, In India, Himalaya and the Western Ghats harbor a large number of ant species, 656 species from 88 genera were recorded in Himalaya, and 455 species from 75 genera were recorded in the Western Ghats, especially in Tamil Nadu, 184 species from 51 genera were recorded [2 and 3]. Ants in India, occupy a variety of habitats such as leaf litter, trees, soil and dead logs, while tramp species prefer human-modified habitats. Some species even form symbiotic association with particular group of plants, which produce suitable preformed nest sites to attract the ants to take up residence [4]. The design and architecture of nest is distinctly purposeful and constructed with patience. They construct nests in various types of habitats, some nest in plant cavities, but majority of ants make nests in the ground. Same colony of ants may adopt very different methods of nest building at different periods during the growth and development [5]. The ability of ants to build its nests ranges from subterranean (i.e. terrestrial and/or intertidal habitats) [6], to lignicolous, lithophilic, and arboreal [7 and 8], with nests constructed of various combinations [9]. Ants play an important role in terrestrial ecosystems such as pollinators, seed dispersal, predators of harmful insects, good soil turners and as a food source for other animals. The plants and produce make suitable preformed nest sites to attract the ants to take up residence.

Despite the importance of ants in different ecosystems, there is little information on abundance and diversity of ants nesting beneath different trees in Periyanaickenpalayam, TamilNadu. The present study therefore investigated the abundance and diversity of ant species nesting beneath different trees in Periyanaickenpalayam.

## 2. MATERIALS AND METHODS

### 2.1 Study area

The field work was conducted in the Periyanaickenpalayam village, Coimbatore district, TamilNadu. Coimbatore, which is located between Latitude 11°16'N and Longitude 76°58'21"E, in south India at 411 m.a.s.l on the banks of the Noyyal River, in southwestern Tamil Nadu. The average annual rainfall is around 700 mm (27.6 in) with the northeast and the southwest monsoons contributing to 47% and 28%, respectively, to the total rainfall. Periyanaickenpalayam is a neighborhood in Coimbatore in the Indian state of Tamil Nadu. It is located along National Highway NH 67, Mettupalayam road, an arterial road in Coimbatore.

#### 2.1.1 All out search method

The village was divided into 8 equal size zones. Ants were collected twice a day, each of two hours duration. (6-8 and 16-18 h Indian Standard Time (IST)). One day was devoted to each zone from November 2017 to February 2018. Ants were collected using a brush and forceps during day time in between 11am to 16 h IST twice in every month. Collected ants were transferred into 70% ethanol in plastic vials at the Department of Zoology, PSG College of arts and science and same time maintained for all zones while ants collection. The stored ant specimens were then counted and identified up to genus level (some to species level) using microscope. Species identification was carried out under the help of the keys of "Ants identification guide" [10] collected ants were identified up to the genus level by using based on literature [11, 4, 12, 13 and 14]. Identified specimens will be kept in the air tight insect wooden box. Ant species were listed and each species was counted to calculate and compared composition, richness, species diversity, trees association, habitat type and identification of ants.

## 3. RESULTS

**Table 1 showing the ant nests with floral association on Periyanaickenpalayam village.**

Common name	Scientific Name	Number of nests	Subfamily	Genera	Species
Neem tree	<i>Azadirachta indica</i>	10	Formicinae	Camponotus	radiatus
sacred fig	<i>Ficus religiosa</i>	20	Formicinae	Camponotus	compressus
Sal	<i>Shorea robusta</i>	16	Formicinae	Camponotus	irritans
Coconut	<i>Cocos nucifera</i>	11	Formicinae	Camponotus	parius
Banyan	<i>Ficus benghalensis</i>	23	Formicinae	Camponotus	sericeus
Sacred fig	<i>Ficus religiosa</i>	19	Formicinae	Camponotus	fabricus
Sandal wood	<i>Santalum album</i>	11	Formicinae	Camponotus	maculatus
Mango	<i>Mangifera indica</i>	23	Formicinae	Oecophylla	smaragdina
Toona	<i>Toona ciliata</i>	12	Formicinae	Anoplolepis	gracillipes
White teak	<i>Gmelina arborea</i>	19	Formicinae	Paratrechina	Longicorns
Neem tree	<i>Azadirachta indica</i>	18	Myrmicinae	Monomorium	Minimum

Bamboo	<i>Bambuseae</i>	22	Myrmicinae	Monomorium	Destructor
Papaya	<i>carica papaya</i>	15	Myrmicinae	Monomorium	Pharaonis
Karuva	<i>Prosopis juliflora</i>	11	Myrmicinae	Crematogaster	Species
Porita tree	<i>Thespepsia populnea</i>	6	Myrmicinae	Crematogaster	Subnuda
Gulmohar	<i>Delonix</i>	13	Myrmicinae	Solenopsis	Invicta
Mango	<i>Mangifera indica</i>	10	Myrmicinae	Solenopsis	Germinata
Neem tree	<i>Azadirachta indica</i>	11	Myrmicinae	Solenopsis	Diplorhoptom
Sacred fig	<i>Ficus religiosa</i>	4	Myrmicinae	Phediole	Spp
Tamarind	<i>Tamarinds indica</i>	7	Myrmicinae	Phediole	Megacephala
Indian laburnum	<i>Cassia fistula</i>	7	Ponerinae	Lepitogenys	Processionalis
Banana tree	<i>Musa acuminate</i>	10	Dolichoderinae	Tapinoma	Indicum
Karuva	<i>Cinnamoumverum</i>	5	Dolichoderinae	Tapinoma	Sessile
Karuva	<i>Cinnamoumverum</i>	13	Pseudomyrmicinae	Tetraponera	Species
Neem tree	<i>Azadirachta indica</i>	7	Pseudomyrmicinae	Tetraponera	Nigra
Toona	<i>Toona ciliata</i>	10	Pseudomyrmicinae	Tetraponera	Rufonigra
Karuva	<i>Cinnamoumverum</i>	9	Pseudomyrmicinae	Tetraponera	Allaborans
Sal	<i>Shorea robusta</i>	7	Formicinae	Camponotus	flying ants
Karuva	<i>Cinnamoumverum</i>	9	Myrmicinae	Solenopsis	Germinata
Neem tree	<i>Azadirachta indica</i>	6	Pseudomyrmicinae	Tetraponera	nigra(flying)
Mango	<i>Mangifera indica</i>	7	Formicinae	Camponotus	Species
Sacred fig	<i>Ficus religiosa</i>	6	Myrmicinae	Tetramorium	Species
Karuva	<i>Prosopis juliflora</i>	7	Myrmicinae	Crematogaster	Species

The study observed that 384 ants nests associated with 27 floral species at different locations in Periyanaickenpalyam Village, TamilNadu state were recorded (Table 1). The study recorded the approximate height (H) of tree, diameter (D) of tree and ants nest entrance size (NES). Based on floral ants nest association, eighteen percentage of ants species associated with *Prosopis juliflora* tree with the height of (14-16ft), diameter (14-21cm) and nest entrance size ranges in between (0.8-1.2 cm) followed by fifteen percentage of ants species allied in *Azadirachta indica* with (9-16ft) in height, diameter of (60-70cm) and nest size of (1-1.5 cm), *Ficus religiosa* species contain twelve percentage of ants species with the distinctive features of trees height (40-66 ft) with diameter (40-50cm) and nest size (0.9-1.5 cm), *Mangifera indica* species contain nine percentage of ants species with the height of (19-22 ft) and the diameter of (23-28cm) with the nest size (0.9-1.5 cm), *Shorea robusta* contain six percentage of ants species with (H: 19-23 ft, D:40- 55 cm and NES: 1-12 cm) and six percentage of species associated with *Toona trees* with the height of ( 9-12 ft), including the diameter of (20-30cm) and nest size of (1-1.5 cm) were recorded. The rest of floral species of *Cocos nucifera species* with the height of (20ft) and the diameter of (30cm) and nest size of (0-7-1.2 cm), *Ficus benghalensis* with height (9-16 ft) and diameter of (60cm) with nest size of (1-1.7 cm), *Santalum album* with the height of twenty five feet with the diameter of sixteen centimeter and the nest size of (1-2 cm), *Gmelina arborea* with height and diameter of (H: 13 ft, D:12cm, NS: 0.5-1 cm), *Bambuseae* with height and diameter of (5 ft, 5 cm), and nest size of opening of (1-1.5 cm), *Carica papaya* approximate height of five feet with the diameter of (10 cm) and Nest size of (1.3-1.5 cm), *Thespepsia populnea* with height of sixteen feet and diameter of (44cm) with the nest size of (1-1.2 cm), *Delonix* with height of eight feet and diameter of (41 cm) with the nest size of (0.8-1.4 cm), *Tamarinds indica* height of thirty two feet and the (26cm) diameter with the nest size of (1-1.6 cm), *Cassia fistula* with height of sixteen feet and diameter of (54 cm) with nest size of (1-1.5 cm), *Musa*

*acuminate* floral species with height of fifteen feet with the diameter of ( 10 cm) and the nest size(0.4-.7 cm), *Prosopis juliflora* tree species with height of sixteen feet and diameter of (20-33cm) along with the nest size( 0.7-1.1 cm) were observed from the study.

According to nest wise, out of 384 ant's nests, fifty four nests (14%) in *Prosopis juliflora* floral species under three subfamilies of ants were observed. Twenty seven nests (50%) in Myrmicinae, twenty two nests (41%) in Pseudomyrmicinae and five nests (9%) in Dolichodeni species were recorded. In *Azadirachta indica* (13.5%) species which contain fifty two nests. Out of 52 nests, subfamily Myrmicinae have twenty nine nests (56%), thirteen nest in Pseudomyrmicinae (25%) and ten Formicinae nests consist of (19%) were observed.

Forty nine ants' nests species present in *Ficus religiosa* trees which restrain (12.7%) and further undergoes into thirty nine nests (80%) in subfamily Formicinae and ten nests (20%) in Myrmicinae subfamilies were recorded. Forty nests occupied in *Mangifera indica* contain (10.5%) nests which was further undergoes into thirty (75%) nest in subfamily Formicinae and ten nest (25%) in Myrmicinae nest were recorded.

twenty three nests present in *Shorea robusta* species which comprise of 6% ants nests further undergoes twenty three ants nests (100%) in subfamily *Formicinae* nests. Twenty two nests present in *Toona ciliata* species enclose (5.8%) nests under subfamily Formicinae contain twelve nests (55%) and ten nest (45%) in *Pseudomyrmicinae* species were observed. The other faunal varieties like twenty three nests in *Ficus benghalensis* species contain 5.8%, *twenty two Bambuseae* nest comprise of 5.7%, *nineteen nest in Gmelina arborea* contain 4.9%, *fifteen nests in Carica papaya* enclose 3.9%, *thirteen nests in Delonix* contain 3.4%, *eleven nests in Cocos nucifera* species contain 2.8% nests, eleven nests in *Santalum album* includes 2.8%, *ten nests in Musa acuminate* include 2.6%, seven nests in *Tamarinds indica* contain 1.9% ,seven nests in *Cassia fistula* species enclose 9% and six nests in *Thespepsia populnea* comprise of 1.6% of nests were observed from the study.

#### 4. DISCUSSION

Domatia are internal plant structures that appear to be specifically adapted for habitation by ants [15]. These cavities are found primarily in the stems, leaves, and spines of plants. Many different genera of plants offer domatia. The study observed that 384 ants nests associated with 27 floral species at different locations in Periyannayakenpalayam were recorded based on our photos and searching method. Based on floral -ants nest association maximum 18% of *Prosopis juliflora* followed by 15% of *azadirachta indica*. Plants of the *Acacia* genus have some of the most widely recognized forms of domatia and offer some of the best examples of ant-plant obligate mutualism [15]. Different species of *Acacia* provide a variety of resources needed for their codependent counterparts. One of these resources is the need for shelter. *Acacia* have enlarged thorns on their stems that are excavated by ants for use as housing structures.

*Ficus religiosa* contain 12%, *Mangifera indica* contain 9%, *Shorea robusta* (6%) and *Toona ciliata* both contain (6%). The *Shorea robusta*, *Mangifera indica* are formed building up leaf domatia. The rolling pattern of leaf domatia in *Pterospermum sp* and the complete leaf domatia on a hemi parasitic angiosperm (*Lorenthus longifolia*) in Manikara plant species. In the interesting, hanging like leaf domatia found in *Swietenia mahogany*. Some plants produce food bodies for use by other organisms [16].

Floral species *Cocos nucifera*, *Ficus benghalensis*, *Santalum album*, *Gmelina arborea*, *Bambuseae*, *carica papaya*, *Thespepsia populnea*, *Delonix*, *Tamarinds indica*, *Cassia fistula*, *Musa acuminate* each contain(3%) were observed. Mutualism between plants and ants is widespread. Two of these mutualisms involving ants in protecting plants from herbivores, and in seed dispersal (myrmecochory), are well known and intensively studied [17, 18, 19 and 20].

Colony structure can be highly variable, with some species establishing nests through either single or multiple queens [21 and 22] or obligatory parasitic relationships with other species of ants [23]. Individual nests can contain numerous dimorphic queens, each of which has a full set of thoracic clerites and seemingly functional wings. Both queen morphs appear capable of reproduction, possessing apparently functional ovaries, and together present a typical bimodal but continuous size frequency distribution [24].

According to nest wise showed that, out of 384 ants nests associated with 27 floral species at different locations in periyannayakenpalayam were recorded. Ants nests maximum in (n=54) nest with the *Prosopis juliflora* species which undergoes three subfamily; 27(50%) nest of myrmicinae, 22(41%) nests in pseudomyrmicinae and 5(9%) dolichodeni nests followed by (n=52) nests in *azadirachta indica* which under goes 29 (56%) in Myrmicinae, Pseudomyrmicinae 13(25%) and Formicinae

consist of 10(19%) nests, (n=49) nests in *Ficus religiosa* contain 39(80%) Formicinae nests and 10(20%) Myrmicinae nest, (n=40) nests in *Mangifera indica* contain 30(75%) Formicinae nests and 10(25%) myrmicinae nest and (n=23) *Shorea robusta* contain nests under 23(100%) Formicinae nests and (n=22) *Toona ciliata* contain nests under formicinae 12 (55%) nests and 10 (45%) Pseudomyrmicinae. Worker size is strongly bimodal [25]: Smaller (minor) workers perform tasks within the nest while larger (major) workers carry out a range of tasks both within and outside the nest [26] reported that ant species richness generally increased with increase in vegetation. Tree hollow, tree holes and dead limbs are the most common nesting site for this species [27 and 28]. Many myrmecophytes are defended from both herbivores and other competing plants by their ant counterparts (Acacia cornigera, for example, is thoroughly guarded by its obligate ant partner, Pseudomyrmex ferruginea. A single colony of P. ferruginea may contain more than 30,000 ants, and can tend multiple Acacia trees. The soldier ants are extremely aggressive, patrolling the trees twenty-four hours a day. Any disturbance to the tree alerts ants, who then recruit more workers from inside the horn domatia. These ants defend the Acacia by biting, violently stinging, and pruning any trespassers. The ants keep the plant free from other insects and vertebrate herbivores, but also from invading fungi and other plants [19].

Faunal varieties like *ficus benghalensis* (n=23) contain 5.8%, *bambuseae* (n=22) contain 5.7%, *gmelina arborea* (n=19) contain 4.9%, *carica papaya* (n=15) contain 3.9%, *delonix* (n=13) contain 3.4%, *cocos nucifera* (n=11) contain 2.8% nests, *santalum album* (n=11) contain 2.8%, *musa acuminata* (n=10) contain 2.6%, *tamarinds indica* (n=7) contain 1.9%, *cassia fistula* (n=7) contain 1.9% and *thespepsia populnea* (n=6) contain 1.6%, nests were observed from the study. Nesting location within Polyrhachis species for example can vary from intertidal and subterranean to arboreal, the presence of silk nests and/or larval cocoons is highly variable and disjunct, and even the source of silk within nests can vary from their own larvae to spiders silk.

Since the tree contains their nest, these aggressive ants react strongly to any disturbance of the tree, providing the myrmecophyte with defense from grazing herbivores and encroaching vines. The ants continuously patrol the surface of their host plant and protected it from depend on this protection and grow poorly in the absence of their ant partner.


Plants of the Acacia genus have some of the most widely recognized forms of domatia and offer some of the best examples of ant-plant obligate mutualism [15]. Different species of Acacia provide a variety of resources needed for their codependent counterparts. One of these resources is the need for shelter. Acacia have enlarged thorns on their stems that are excavated by ants for use as housing structures. Since the tree contains their nest, these aggressive ants react strongly to any disturbance of the tree, providing the myrmecophyte with defense from grazing herbivores and encroaching vines.

Recent work involving principally taxonomic and ecological studies indicates that nesting habits and ecology in the taxonomically extensive ant genus Polyrhachis are almost as diverse as that of all ants in general [29], offering the unique potential to explore the evolution of nest weaving within a single genus.

## CONCLUSION

A total of 33 floral species associated with ants nesting habitat have been recorded from Periyankenpalayam village, Coimbatore district. During this study, out of thirty three floral species, Karuva tree followed by Neem tree, Sacred fig, Mango tree, Sal and toona species accounted for contain 63% of ants nests was occupied. The present study will yield valuable information on ant species availability in this region. Finally, to sum up, this study provides a little information about ants nesting association with floral species. In future study we developed to research soil developing and decomposing of soil during the nesting behavior

### PLATE 1



1 *Camponotus* spp, 2 *Camponotus compressus*, 3 *Camponotus sericeus*, 4 *Camponotus fabricius*, 5 *Monomorium destructor*, 6 *Crematogaster subnuda*, 7 *Camponotus* spp, 8 *Camponotus compressus*, 9 *Camponotus* spp, 10 *Monomorium pharaonis*, 11 *Oecophylla smaragdina*, 12 *Paratrechina longicornis*

## REFERENCE

1. Bolton B. Bolton's Catalogue and Synopsis, in <http://gap.entclub.org/> Version: 1, 2011.
2. Bharti H. List of Indian Ants (Hymenoptera: Formicidae). Halteres, 2011; 3:79-87.

3. Bharti H, Guénard B, Bharti M, Economo EP. An updated checklist of the ants of India with their specific distributions in Indian states (Hymenoptera, Formicidae). *ZooKeys*, 2016; 551:1-83.
4. Bolton B. Identification guide to the ant genera of the world. Cambridge, Mass: Harvard University Press, 1994, 222
5. Wheeler, W.M. Ants, their structure, Development and Behaviour. Colombia University Biological Series 9, 1913.
6. Nielsen M.G. Nesting biology of the mangrove mud-nesting ant *Polyrhachis sokolova* Forel (Hymenoptera, Formicidae) in northern Australia. *Insect. Soc.* 1997; 44: 15–21.
7. Liefke C., Dorow W.H.O., Hölldobler B. and Maschwitz U. Nesting and food resources of syntopic species of the ant genus *Polyrhachis* (Hymenoptera, Formicidae) in West-Malaysia. *Insect. Soc.* 1998; 45: 411–425.
8. Robson S.K. Comparative nesting biology of two species of Australian lithocolous ants: *Polyrhachis* (Hedomyrma) turneri Forel and *P.* (Hagiomyrma) thusnelda Forel (Hymenoptera : Formicidae : Formicinae). *Aust. J. Entomol.* 2004; 43:5–9.
9. Robson S.K.A. and Kohout R.J. Evolution of nest-weaving behaviour in arboreal nesting ants of the genus *Polyrhachis* Fr. Smith (Hymenoptera: Formicidae). *Aust.J.Entomol.* 2005; 44:164–169.
10. Bayer. Ant identification guide. *Bayer environmental science*, 2010.
11. Mathew R, and Tiwari, R. N. Insecta: Hymenoptera: Formicidae. *Zool. Surv. India. Fauna of Meghalaya. State Fauna Series. Part 7*, 2000; 4.
12. Bingham, C. T. *The Fauna of British India, including Ceylon and Burma. Hymenoptera, Ants and Cuckoo-wasps* London: Taylor and Francis (02) .1903; 1-506.
13. Hölldobler, B, Wilson, E. O. The ants. Cambridge, MA: Harvard University Press.732; 1990.
14. Krebs, C.J, Addison- S. Prasad, P Puyravaud, B.R. Ramesh, K.A. Ecological methodology. *Educational Publishers*. California. 1999; 581.
15. Janzen, D.H. Coevolution of mutualism between ants and acacias in Central America. *Evolution*.1966; 20: 249– 275.
16. Shilpa Dinda & Amal Kumar Mondal. Biodiversity of Myrmecophytes in Eastern India. 2015; 6 (4): 625-631
17. Herrera CM, Pellmyr O Plant-animal interactions: an evolutionary approach. Blackwell Publishing, Malden, 2002.
18. Bronstein JL, Ruben AR, Geber M (2006) Tansley review: the evolution of plant–insect mutualisms. *New Phytol* 172:412–428.
19. Rico-Gray V, Oliveira PS the ecology and evolution of antplant interactions. University of Chicago Press, Chicago. 2007.
20. Schaefer HM, Ruxton GD. Plant-animal communications. Oxford University Press, Oxford. 2011.
21. Sasaki K, Jibiki E, Satoh T, and Obara Y. Queen phenotype and behaviour during cooperative colony founding in *Polyrhachis moesta*. *Insect. Soc.* 2005; 52: 19–25.
22. Yamauchi K., It Y., Kinomura K. and Takamine H. Polycalic colonies of the weaver ant *Polyrhachis dives*. *Kontyu* 1987; 55: 410–420.
23. Maschwitz U., Go C., Dorow W.H.O., Buschinger A. and Kohout R.J. *Polyrhachis loweryi* (Formicinae): A guest ant parasitizing *Rhytidoponera* sp. (Ponerinae) in Queensland, Australia. *Insect. Soc.* 2003; 50: 69–76.
24. Heinze J. and Hölldobler B. Queen polymorphism in an Australian weaver ant, *Polyrhachis* cf. *doddi* Psyche.1993; 100: 83–92.
25. COLE, A.C. Jr. & JONES, J.W. A study of the weaver ant, *Oecophylla smaragdina* (Fab.). – *American Midland Naturalist* 1948; 39: 641-651.
26. Sunil kumar M KT, Nair P, Varghese T, Gadagkar R. Ant species richness at selected localities of Bangalore *Insect Environment* 1997; 3:3-5.
27. Robinson W. *Urban Entomology* Chapman Hall, London. 1996; 262-284.
28. Hölldobler B, Wilson EO. *The Super-Organism: The Beauty elegance, and Strangeness of Insect Societies*, 2009.
29. Robson SKA, Kohout RJ. A review of the nesting habits and socioecology of the ant genus *Polyrhachis* Fr. Smith. *Asian Myrmecol.* 2007; 1:81–99.

