

Effects of Different Types of Organic Fertilizers on Growth Performance of *Amaranthus caudatus* (Samaru Local Variety) and *Amaranthus cruentus* (NH84/452)

ABSTRACT

Aims: To evaluate the effect of different types organic fertilizers on growth performance of *Amaranthus caudatus* (Samaru local variety) and *Amaranthus cruentus* (NH84/452).

Study design: A randomized complete block design (RCBD) was used for the experiment.

Place and Duration of Study: The field experiment was carried out in the nursery of a homestead garden at No 20, Isaiah Balat Street, Sabo GRA, Kaduna State, Nigeria.

Methodology: The study consists of seven treatments which includes control (no fertilizer), 5 t ha⁻¹ and 10 t ha⁻¹ poultry manure, 5 t ha⁻¹ and 10 t ha⁻¹ sewage sludge, 35 kg ha⁻¹ and 70 kg ha⁻¹ NPK compound fertilizer and also with *Amaranthus caudatus* (Samaru local variety) and *Amaranthus cruentus* (NH84/452) in factorial arrangement fitted into a randomized complete block design (RCBD) and replicated three times. Growth performance data were collected on plant height, number of leaves, leaf area and leaf area index from 2 weeks after transplanting (WAT) to 6 weeks after transplanting (WAT).

Results: The plant height and number of leaves of the two varieties were found in the range of 18.30 - 135.67cm and 13.33 - 78.33cm respectively. Leaf area and leaf area index of the two varieties had values in the range of 41.71 - 258.29cm² and 1.76 - 41.72 respectively. At 6WAT, 10 t ha⁻¹ poultry manure recorded the highest value for all the growth parameters for both varieties except for leaf length, leaf width and leaf area of *Amaranthus caudatus* (Samaru local variety), where 10 t ha⁻¹ sewage sludge and 70 kg ha⁻¹ NPK compound fertilizer were highest.

Conclusion: The experimental results of this study have shown that poultry manure had higher growth performance on the two varieties of Amaranth when compared with sewage sludge and NPK compound fertilizer. The application of poultry manures at 10 t ha⁻¹ is therefore recommended for farmers to use to obtain higher yields of Amaranth.

Keywords: Growth, Organic and Inorganic fertilizers, Amaranth, Soil and Insecticides

1. INTRODUCTION

Increasing population of the world has doubled the food demands and inundated the available land resources. Alongside other food alternatives, vegetables are considered cheap source of energy (1). Vegetables are rich sources of essential biochemicals and nutrients such as carbohydrates, carotene, protein, vitamins, calcium, iron, ascorbic acid and palpable concentration of trace minerals (2).

Amaranth has been one of the most important vegetables of Amaranthaceae family. Amaranth has been naturalized in central parts of Asia and possibly Iran (3) and has cultivation history of more than 2000 years (4). Cultivation of the various *Amaranthus* species is acquiring increasing importance in Nigeria and other parts of African continent where the available species are grown for their leaves (5)

Organic and inorganic fertilizers are essential for plant growth as it supplies plants with the nutrients needed for optimum performance. Organic fertilizer has been used for many centuries whereas chemically synthesized inorganic fertilizers were only widely developed during the industrial revolution.

47 Inorganic fertilizers have significantly supported global population growth, as it has been estimated that
48 almost half the people on the earth are currently fed as a result of artificial nitrogen fertilizer use (6).
49 Commercial and subsistence farming has been and is still relying on the use of inorganic fertilizers for
50 growing crops (7). This is because they are easy to use, quickly absorbed and utilized by crops. The
51 continued dependence of developing countries on inorganic fertilizers has made prices of many
52 agricultural commodities to skyrocket (7).

53 Moreover, most vegetable farmers in tropical Africa are small holders who cannot afford the cost of
54 inorganic fertilizers, although soil fertility limits yield of vegetables especially in urban centres (8). In
55 Nigeria, fertilizers, being costly and sometimes scarce can make farmers not apply enough for good
56 growth (5). Fertilizer application rates in intensive agricultural systems have increased drastically during
57 recent years in Nigeria. Farmers depend largely on locally sourced organic fertilizers (8). In Nigeria, huge
58 amount of organic wastes such as poultry waste, animal excreta, sewage sludge, refuse soil and palm oil
59 mill effluent are generated and heaped on dump sites, posing potential environmental hazard.
60 Incorporating these waste materials into the soil for crop production is expected to be beneficial to the
61 buildup of organic matter layer that is needed for a steady supply of nutrients by tropical soils (9).

62 Oyediji *et al.* (10) reported that NPK and poultry manure improved the growth and yield of three different
63 species of amaranth (*Amaranthus hybridus*, *Amaranthus deflexus* and *Amaranthus cruentus*) but
64 influenced proximate composition differently. Emede *et al.* (11) reported that poultry manure influenced
65 the plant growth and yield of *Amaranthus cruentus* L. positively. Therefore, the objective of this study was
66 to determine the effect of different types of organic fertilizers on the growth performance of *Amaranthus*
67 *caudatus* (Samaru local variety) and *Amaranthus cruentus* (NH84/452)

68

69 2. MATERIALS AND METHODS

70 2.1 Seeds

71 The seeds of *Amaranthus caudatus* (Samaru local variety) were obtained from local farmers in Samaru,
72 Zaria, Nigeria while the seeds of *Amaranthus cruentus* (NH84/452) were obtained from National
73 Horticultural Research Institute (NIHORT), Ibadan, Nigeria. *Amaranthus caudatus* samples collected were
74 authenticated at the herbarium unit of Biological Sciences Department, Ahmadu Bello University Zaria,
75 Nigeria and a voucher specimen was deposited.

76 2.2 Study Area

77 The field experiment was carried out in the nursery of a homestead garden at No 20, Isaiah Balat Street,
78 Sabo GRA, Sabo Tasha, Kaduna State, Nigeria. Kaduna metropolis has a tropical savanna climate with
79 dry winters characterized by maritime air and rainfall is between April and October with annual raining
80 days ranging from 81 to 103 mm. During the reference period, the annual mean rainfall values range from
81 145.37mm to 318.67mm. From the figures above, ample rains are available for the production of many
82 agricultural crops. During harmattan, dry desert wind blows between December and mid February while
83 night temperature is very low. The geographical location of Kaduna metropolis is Latitude 9⁰03' N and
84 11⁰32' N north of the equator and Longitudes 6⁰05' E and 8⁰38' E East of the Greenwich meridian.
85 Kaduna metropolis has a sub-humid semi arid tropical climate with maximum annual mean temperature
86 ranging from 25.30⁰C to 36.20⁰C while the minimum annual mean temperature range of 28.45⁰C to
87 34.38⁰C (12).

88 2.3 Soil sampling

89 Surface soil sample was taken from the experimental site at a depth of 0 – 15cm at land preparation (after
90 ploughing and harrowing) using the zigzag method. The sample was collected from twenty points and
91 bulked to form a composite sample. The composite sample was air-dried, crushed and sieved through a
92 2mm mesh sieve and stored for chemical analysis (13).

93 2.4 Fertilizers

- 94 i. Poultry manure: The poultry manure was collected at Ishaya's poultry farm in Sabo GRA,
95 Sabo Tasha Kaduna State, Nigeria.
- 96 ii. Sewage sludge: The dried packed sewage sludge was collected at the sewage site of
97 Ahmadu Bello University Zaria, Nigeria.
- 98 iii. NPK compound fertilizer: NPK compound fertilizer (15:15:15) was bought at Kawo market
99 Kaduna State, Nigeria.

100 2.5 Soil analysis

101 The sampled soil was analyzed at the Soil Science Department of the Institute of Agricultural Research,
102 Ahmadu Bello University Zaria, Nigeria. The following parameters were analyzed in the sampled soil;
103 particle size, pH (in water), organic carbon, available phosphorus, total nitrogen, cation exchange
104 capacity (CEC) and exchangeable bases (14).

105 2.6 Experimental design and fertilizer treatment

106 The experiment included seven fertilizer treatments for each of the two varieties of Amaranth which are in
107 factorial arrangement fitted into a randomized complete block design (RCBD) and replicated three times.
108 Hence, the experiment had a total of 42 experimental plots. The treatments were: Control (no fertilizer), 5
109 t ha⁻¹ poultry manure, 10 t ha⁻¹ poultry manure, 5 t ha⁻¹ sewage sludge, 10 t ha⁻¹ sewage sludge, 35 kg ha⁻¹
110 NPK compound fertilizer, 70 kg ha⁻¹ NPK compound fertilizer (13).

112 2.7 Planting and nursery management

113 Prior to planting, the amaranth seeds were soaked in water for about 24 hours in order to enhance
114 germination. The soaked seeds were first sown in the nursery of about 1.9 cm deep and were watered
115 twice daily. Appropriate nursery management practices were carried out as at when needed to obtain
116 healthy and uniform seedlings. The experimental site was ploughed, harrowed and prepared into slightly
117 raised beds (plots) of 25cm width × 80cm length dimension preparatory to transplanting the crop
118 seedlings. Poultry manure and sewage sludge were incorporated according to treatment level to specific
119 plots during land preparation, thoroughly mixed with the soil and then left for two weeks to allow for
120 mineralization. Half of the NPK Compound fertilizer was applied at day of transplanting while the balance
121 was applied one week later. After two weeks in the nursery, randomly picked seedlings were transplanted
122 to the well prepare beds (plots). The seedlings were watered twice daily using watering can and the
123 surrounding areas were weeded regularly. The experimental area and the surroundings were kept clean
124 to prevent harbouring of pests. Insects were controlled by using "Dime Force Insecticide" with
125 concentration of 1.5 L/ha (15).

127 2.8 Data collection for growth performance

128 Data were first collected two weeks after transplanting (WAT) and subsequently at one week interval for
129 up to six weeks after transplanting. Two randomly selected plants were tagged and used in each plot for
130 data collection. Data collected included plant height, number of leaves, leaf length and leaf width, while
131 the leaf area and leaf area index were computed (13).

132 **2.8.1 Determination of plant height**

133 Plant height is the length of the plant from the base of the stem (surface of the soil) to the apex of the
134 leaves. Plant height was measured using a measuring tape for the two tagged plants per plot and the
135 average computed (13).

136 **2.8.2 Determination of number of leaves**

137 The number of leaves was counted from the two tagged plants and the average computed (13).

138 **2.8.3 Determination of leaf area**

139 The Leaf Area (LA) was computed by multiplying the Leaf Length (LL) by the Leaf Width (LW) and the
140 product multiplied by the correction factor (13).

141 Calculation;

142 Leaf Area = (Leaf Length × Leaf Width) 0.578.

143 **2.8.4 Determination of leaf area index**

144 The leaf area index (LAI) was computed using this formula (16)

$$145 LAI = Y \times N \times LA \times (AP)^{-1}$$

146 Where: Y = Population of plants per plot (5 plants), N = Average number of leaves, LA = Leaf area, AP =
147 Area of plot (25cm width * 80cm length = 2000cm²)

148

149 **2.9 Statistical analysis**

150 Data was analyzed using the Statistical Package for Social Sciences (SPSS) version 21.0 computer
151 package. Descriptive statistics was used to determine the measures of central tendency. Means were
152 separated using Duncan Multiple Range test. Values with different superscripts down the column are
153 significantly different at $p < 0.05$.

154

155 **3. RESULTS AND DISCUSSION**

156 **3.1 Soil analysis results**

157 Results of analyses of the soil used for this experiment are shown in Table 1. The texture class of the soil
158 is sandy clay loam in which sand was highest with value of $66 \pm 2.0\%$, followed by clay with $24 \pm 3.0\%$
159 and silt was the lowest with value of $10 \pm 1.0\%$. The soil organic carbon, total nitrogen and available
160 phosphorus were $0.46 \pm 0.02\%$, $0.32 \pm 0.01\%$ and $7.4 \pm 0.30\text{ppm}$ respectively. The exchangeable bases
161 of Sodium, magnesium, calcium, potassium and cation exchange capacity (CEC) contents were $0.34 \pm$
162 0.02Cmol/Kg , $0.84 \pm 0.02\text{Cmol/Kg}$, $3.26 \pm 0.05\text{Cmol/Kg}$, $0.65 \pm 0.03\text{Cmol/Kg}$ and $5.7 \pm 0.20\text{Cmol/Kg}$
163 respectively. Soil pH value was 7.7 ± 0.2 .

164

165 **Table 1: Physical and Chemical Properties of Soil used in this Experiment.**

Particulars	Value	Methods
Particle size		
• Clay	$24 \pm 3.0 \%$	USDA
• Silt	$10 \pm 1.0 \%$	USDA
• Sand	$66 \pm 2.0 \%$	USDA
Texture Class	Sandy Clay Loam	USDA
pH (in Water)	7.70 ± 0.20	
Organic Carbon	$0.46 \pm 0.02 \%$	Walkley-Black method
Available Phosphorus	$7.40 \pm 0.30 \text{ ppm}$	Bray and Kurts method
Total Nitrogen	$0.32 \pm 0.01 \%$	Kjeldahl method
Exchangeable bases		
• Calcium (Ca)	$3.26 \pm 0.05 \text{ Cmol/Kg}$	AAS
• Magnesium (Mg)	$0.84 \pm 0.02 \text{ Cmol/Kg}$	AAS
• Potassium (K)	$0.65 \pm 0.03 \text{ Cmol/Kg}$	AAS
• Sodium (Na)	$0.34 \pm 0.02 \text{ Cmol/Kg}$	AAS
• Cation Exchange Capacity (CEC)	$5.70 \pm 0.20 \text{ Cmol/Kg}$	Ammonium saturation

166 Values are mean \pm standard deviation of triplicate analysis.

167 **3.2 Organic fertilizer analysis results**

168 Results of analyses of the organic fertilizers used for this experiment are shown in Table 2. Poultry
169 manure showed a pH of 7.62 ± 0.04 , while the concentrations for total nitrogen, available phosphorus and
170 potassium were found to be $3.53 \pm 0.02\%$, $0.71 \pm 0.05\%$ and $1.61 \pm 0.03\%$ respectively. Sewage sludge

171 pH was found to be 8.25 ± 0.09 ; the concentrations of total nitrogen, available nitrogen and potassium
172 were gotten as $2.44 \pm 0.03\%$, $0.97 \pm 0.02\%$ and $1.33 \pm 0.05\%$ respectively.

173

174 **Table 2: Chemical Properties of Organic Fertilizers used in this Experiment**

Chemical Properties of the Organic Fertilizers used in the Experiment				
	pH (in H ₂ O)	Total Nitrogen (%)	Available Phosphorus (%)	Potassium (%)
Poultry manure	7.62 ± 0.04	3.53 ± 0.02	0.71 ± 0.05	1.61 ± 0.03
Sewage sludge	8.25 ± 0.09	2.44 ± 0.03	0.97 ± 0.02	1.33 ± 0.05

175 Values are mean \pm standard deviation of triplicate analysis.


176

177 **3.3 Plant Height**

178 Plant height was significantly ($P = .05$) higher in plants derived from poultry manure treated plots applied
179 at 10 t ha^{-1} treatment and lowest in plants derived from no fertilizer treatment plots for both *Amaranthus*
180 *caudatus* (Samaru local variety) and *Amaranthus cruentus* (NH84/452) as shown in Fig 1 and Fig 2. At 2
181 Weeks After Transplanting (2WAT), the plant height was $26.28 \pm 1.07 \text{ cm}$ and $45.97 \pm 0.88 \text{ cm}$ from poultry
182 manure applied at 10 t ha^{-1} treatment for *Amaranthus caudatus* (Samaru local variety) and *Amaranthus*
183 *cruentus* (NH84/452) respectively, which was also consistently highest till maturity (6WAT) as shown in
184 Fig1 and Fig 2. The highest plant height for the two varieties of amaranth at 6WAT were both observed in
185 plants treated with 10 t ha^{-1} poultry manure which was significantly ($p < 0.05$) different from the other
186 treatments. This position was earlier reported by Egharevba and Ogbe (17) and Okokoh and Bisong (18).
187 The highest plant height exhibited by plants treated with 10 t ha^{-1} poultry manure might have been due to
188 the presence of the primary nutrients plus other minerals found in inorganic manure, and also it may be
189 probably due to favourable nutrient mineralization of poultry manure as a result of the influence of the
190 mineral component on the organic content of the manure (19). The control plants had the lowest height as
191 they had to depend mainly on the intrinsic soil fertility as exhibited by the soil chemical analysis to be low.
192 A similar effect for control was reported for *Amaranthus caudatus* by Abayomi and Adebayo (19) and on
193 radish stems amaranth-indian spinach by Islam *et al.* (20). The height of the plant is an important growth
194 character directly linked with the productive potential of plants. An optimum plant height is claimed to be
195 positively correlated with productivity of plants (21).

196

197



198

199 **Fig 1: Effect of Organic Fertilizers on Plant Height of *Amaranthus caudatus* (Samaru local variety)**

200 Mean values \pm standard deviation of triplicate analysis

201 **WAT=Week after transplanting Control= No fertilizer PM1=Poultry manure at 5 t ha⁻¹ PM2= Poultry**


202 **manure at 10 t ha⁻¹ SS1= Sewage sludge at 5 t ha⁻¹ SS2= Sewage sludge at 10 t ha⁻¹**

203 **NPK1= NPK compound fertilizer at 35 kg ha⁻¹ NPK2= NPK compound fertilizer at 70 kg ha⁻¹**

204

205

206



207

208 **Fig 2: Effect of Organic Fertilizers on Plant Height of *Amaranthus cruentus* (NH84/452)**

209 Mean values \pm standard deviation of triplicate analysis

210 **WAT=Week after transplanting Control= No fertilizer PM1=Poultry manure at 5 t ha⁻¹ PM2= Poultry**

211 **manure at 10 t ha⁻¹ SS1= Sewage sludge at 5 t ha⁻¹ SS2= Sewage sludge at 10 t ha⁻¹**

212 **NPK1= NPK compound fertilizer at 35 kg ha⁻¹ NPK2= NPK compound fertilizer at 70 kg ha⁻¹**

213

214


215

216

217 **3.4 Number of leaves**

218 The number leaves were highest for plants treated with 10 t ha⁻¹ poultry manure for both varieties of
 219 amaranth and were not significantly (P= .05) different among the treatments except for the plants in the
 220 control group as shown in Fig 3 and Fig 4. At 6WAT, poultry manure applied at 10tons/ha gave the
 221 highest number of leaves with values of 78.67 ± 5.03 and 64.50 ± 3.50 for *Amaranthus caudatus* (Samaru
 222 local variety) and *Amaranthus cruentus* (NH84/452) respectively. At the start of the experiment, the
 223 average number of leaves was highest for poultry manure and NPK compound fertilizers for both varieties
 224 of Amaranth. However, between 2WAT and 4WAT, the highest development of new leaves was observed
 225 in 70 kg ha⁻¹ NPK compound fertilizer but not significantly different from 35 kg ha⁻¹ NPK and poultry
 226 manure. Relatively high content of nitrogen in the NPK compound fertilizer increase the growth and
 227 development of new leaves. Normally inorganic fertilizer nutrients are soluble, so the nitrogen was quickly
 228 released into the soil leading to fast leaf growth and development. Although, during maturity leaf
 229 development declined because the nutrients were probably exhausted in the soil; however, the reason for
 230 the high number of leaves for plants treated with poultry manure compared to the sewage sludge at the
 231 early stages was attributed to the high amount of nitrogen in the poultry manure than sewage sludge from
 232 chemical analysis, and also due to faster mineralization and release of nutrients from the poultry manure
 233 than sewage sludge. At maturity, the 10 t ha⁻¹ poultry manure showed the highest average number of
 234 leaves for both varieties of Amaranth, which was also reported by Law-Ogbomo and Ajayi (22) for
 235 *Amaranthus cruentus*. This also agrees with reports by previous workers such as Sanwal *et al.*(23) in
 236 turmeric (*Curcuma longa*); Premesekhar and Rajashree (24) in Okra (*Abelmoschus esculentus*) who
 237 separately attributed higher leaf yield to released nutrients from organic manure application which
 238 improved chemical, physical and biological properties of soil. This high leaves development in the poultry
 239 manure compared to the sludge is due to the higher amount of nitrogen in poultry manure and continuous
 240 release of the nutrients. However, the reason behind the higher number of leaves for plants treated with
 241 organic fertilizers than the NPK compound fertilizer may be due to availability of nutrients as affected by
 242 the water holding capacity of the soil (25). Most probably because as the manure quantities increased the
 243 water holding capacity of the soil and subsequent nutrient release increases, while the NPK compound
 244 fertilizer nutrients have been exhausted as the early stages due to the solubility of the nutrients.

245



246 **Fig 3: Effect of Organic Fertilizers on Number of Leaves of *Amaranthus caudatus* (Samaru local**
 247 **variety)**

248 Mean values ± standard deviation of triplicate analysis

249 **WAT=Week after transplanting Control= No fertilizer PM1=Poultry manure at 5 t ha⁻¹ PM2= Poultry**
 250 **manure at 10 t ha⁻¹ SS1= Sewage sludge at 5 t ha⁻¹ SS2= Sewage sludge at 10 t ha⁻¹**
 251 **NPK1= NPK compound fertilizer at 35 kg ha⁻¹ NPK2= NPK compound fertilizer at 70 kg ha⁻¹**

252
 253
 254




Fig 4: Effect of Organic Fertilizers on Number of Leaves of *Amaranthus cruentus* (NH84/452)

Mean values \pm standard deviation of triplicate analysis


WAT=Week after transplanting **Control**= No fertilizer **PM1**=Poultry manure at 5 t ha⁻¹ **PM2**= Poultry

manure at 10 t ha⁻¹ **SS1**= Sewage sludge at 5 t ha⁻¹ **SS2**= Sewage sludge at 10 t ha⁻¹

NPK1= NPK compound fertilizer at 35 kg ha⁻¹ **NPK2**= NPK compound fertilizer at 70 kg ha⁻¹

3.5 Leaf Area

At maturity, leaf area which is a measure from the leaf length and leaf width was significantly ($P= .05$) higher in plants derived from plots treated with 10 t ha⁻¹ sewage sludge with area of $127.36 \pm 3.40\text{cm}^2$ and 10 t ha⁻¹ poultry manure with area of $258.29 \pm 23.96\text{cm}^2$ for *Amaranthus caudatus* (Samaru local variety) and *Amaranthus cruentus* (NH84/452) respectively and was lowest in plants derived from plots with no fertilizer (control treatment) as shown in Fig 5 and Fig 6. Leaf area were found to be highest in plants from the 70 kg ha⁻¹ NPK compound fertilizer for green type and the 10 t ha⁻¹ poultry manure recorded the highest for *Amaranthus cruentus* (NH84/452) but there no significant ($P= .05$) difference among the treatments except for the control treatment. Similar work also reported by Mshelia and Degri (26) on effect of different levels of poultry manure on performance of *Amaranthus caudatus* L. Okokoh and Bisong (18) reported similarly in a research in Calabar that application of poultry manure significantly influenced performance of amaranth. The increase in leaf area had been claimed to be directly influence by nitrogen supply in fertilizer applied (27). The insignificant difference among the treatments suggests that the nutrients in both the organic and inorganic fertilizers increases leaf width but higher amount of nutrients in individual fertilizers may not necessary influence noticeable difference in the width of the plants.




281
282 **Fig 5: Effect of Organic Fertilizers on Leaf Area of *Amaranthus caudatus* (Samaru local variety)**

283 Mean values \pm standard deviation of triplicate analysis

284 **WAT=Week after transplanting Control= No fertilizer PM1=Poultry manure at 5 t ha⁻¹ PM2= Poultry**
 285 **manure at 10 t ha⁻¹ SS1= Sewage sludge at 5 t ha⁻¹ SS2= Sewage sludge at 10 t ha⁻¹**
 286 **NPK1= NPK compound fertilizer at 35 kg ha⁻¹ NPK2= NPK compound fertilizer at 70 kg ha⁻¹**

287
288



289
290 **Fig 6: Effect of Organic Fertilizers on Leaf Area of *Amaranthus cruentus* (NH84/452)**

291 Mean values \pm standard deviation of triplicate analysis


292 **WAT=Week after transplanting Control= No fertilizer PM1=Poultry manure at 5 t ha⁻¹ PM2= Poultry**
 293 **manure at 10 t ha⁻¹ SS1= Sewage sludge at 5 t ha⁻¹ SS2= Sewage sludge at 10 t ha⁻¹**
 294 **NPK1= NPK compound fertilizer at 35 kg ha⁻¹ NPK2= NPK compound fertilizer at 70 kg ha⁻¹**

295
296
297

298 **3.6 Leaf Area Index**

299 Leaf area index which indicates the photosynthetic ability of the plants was significantly ($P= .05$) higher in
 300 plants derived from plots treated with 10 t ha^{-1} poultry manure for both *Amaranthus caudatus* (Samaru
 301 local variety) and *Amaranthus cruentus* (NH84/452) with values of 23.74 ± 1.96 and 41.72 ± 5.48
 302 respectively and was lowest for plants derived from plots with no fertilizer added for both varieties as
 303 shown in Fig7 and Fig 8. The 10 t ha^{-1} poultry manure treatment resulted in the highest leaf area index for
 304 both varieties of amaranth which is consistent with report on red lettuce (7). The positive effect of poultry
 305 manure increasing leaf area index of amaranth was earlier reported by Egharevha and Ogbe, (17). Law-
 306 Ogbomo and Ajayi (22) also reported similar results on *Amaranthus cruentus*. Leaf area index and
 307 number of leaves follow the same pattern as both are directly related. The higher leaf area index in
 308 poultry manure was caused by the relatively higher nutrient availability which increased the leaf length,
 309 number of leaves and leaf width per unit area of the plot. Normally, inorganic chemical fertilizer nutrients
 310 are soluble, so the nitrogen was quickly released into the soil thus leading to fast leaf growth and
 311 development. However, during maturity, leaf development declined because the nutrients were probably
 312 exhausted in the soil. This resulted in the leveling of the leaf growth and development between NPK
 313 compound fertilizer, sewage sludge and poultry manure at maturity as sewage sludge and poultry manure
 314 was continuously releasing nitrogen. Organic manures like cattle manure and poultry manure have been
 315 reported to release both micro and macro nutrients slowly resulting in subsequent promotion of vegetable
 316 growth (28; 29; 30).

317
318




319 **Fig 7: Effect of Organic Fertilizers on Leaf Area Index of *Amaranthus caudatus* (Samaru local**
 320 **variety)**

321 Mean values \pm standard deviation of triplicate analysis

322 **WAT=Week after transplanting Control= No fertilizer PM1=Poultry manure at 5 t ha^{-1} PM2= Poultry**
 323 **manure at 10 t ha^{-1} SS1= Sewage sludge at 5 t ha^{-1} SS2= Sewage sludge at 10 t ha^{-1}**
 324 **NPK1= NPK compound fertilizer at 35 kg ha^{-1} NPK2= NPK compound fertilizer at 70 kg ha^{-1}**

325
326
327
328



329 **Fig 8: Effect of Organic Fertilizers on Leaf Area Index of *Amaranthus cruentus* (NH84/452)**

330 Mean values \pm standard deviation of triplicate analysis

331 **WAT=Week after transplanting Control= No fertilizer PM1=Poultry manure at 5 t ha⁻¹ PM2= Poultry**
 332 **manure at 10 t ha⁻¹ SS1= Sewage sludge at 5 t ha⁻¹ SS2= Sewage sludge at 10 t ha⁻¹**
 333 **NPK1= NPK compound fertilizer at 35 kg ha⁻¹ NPK2= NPK compound fertilizer at 70 kg ha⁻¹**

336 4. CONCLUSION

337 The rapidly rising cost of chemical fertilizers has forced small scale vegetative farmers to look for
 338 alternatives such as organic fertilizers; the result of this present study justifies the use of some organic
 339 fertilizers over chemical fertilizer due to high vegetative growth and plant development. The present study
 340 revealed that application of poultry manure at 10 t ha⁻¹ on average effected the highest growth and
 341 development of *Amaranthus Amaranthus caudatus* (Samaru local variety) and *Amaranthus cruentus*
 342 (NH84/452) with highest values for plant height, number of leaves, leaf length, leaf width, leaf area and
 343 leaf area index. This effect can be as a result of the slow decomposition and release of nutrients from the
 344 organic fertilizers.

345 COMPETING INTERESTS

346 No competing interests

347 REFERENCES

- 348 1. Hussain J, Khan AL, Rehman N, Hamayun M, Shah T, Nisar M, Lee I. Proximate and nutrient
 349 analysis of selected vegetable species: A case study of Karak region, Pakistan. African Journal of
 350 Biotechnology. 2009; 8(12).
- 351 2. Salunkhe DK, Kadam SS. Handbook of Vegetable Science and Technology: Production,
 352 Composition, Storage and Processing. Marcel Dekker, INC. 1998; 721.
- 353 3. Kawazu Y, Okimura M, Ishii T, Yui S. Varietal and seasonal differences in oxalate content of
 354 spinach. Scientia Horticulturae 2003; 97(3):203-210.
- 355 4. Daneshvar H. Vegetable Farming. Chamran University Press Martyr. 2000; 190:461.
- 356 5. Alonge SO, Alonge FO, Bako SP, Olarewaju JD, Adeniji OB. Effects of rates and split application
 357 of compound NPK fertilizer on the growth and yield of three *Amaranthus* species in Nigeria
 358 guinea savanna. Asian J. Plant Sci. 2007; 6:906-912.
- 359 6. Erisman JW, Sutton MA, Galloway J, Klimont Z, Winiwarter W. How a century of ammonia
 360 synthesis changed the world. Nat. Geosci. 2008; 1:636-639.

- 361 7. Masarirambi MT, Hlawe MM, Oseni OT, Sibiya TE. Effects of organic fertilizers on growth, yield,
362 quality and sensory evaluation of red lettuce (*Lactuca sativa* L.) 'Veneza Roxa'. *Agric. Biol. J.*
363 North America. 2010; 1(6):1319-1324.
- 364 8. Makinde EA, Ayeni LS, Ojeniyi SO, Odedina JN. Effect of organic, organomineral and NPK
365 fertilizer on nutritional quality of Amaranthus in Lagos, Nigeria. *Researcher*. 2008; 2:91-96.
- 366 9. Agboola AA, Omueti JA. Soil fertility problem and its management in the tropical Africa. Paper
367 presented at the International Institute of Tropical Agriculture, Ibadan, Nigeria. 1982; 25.
- 368 10. Oyedeji S, Animasaun DA, Bello AA, Agboola OO. Effect of NPK and Poultry Manure on Growth,
369 Yield, and Proximate Composition of Three Amaranths. *Hindawi Publishing Corporation Journal*
370 *of Botany*. 2014; 10:11–55.
- 371 11. Emede TO, Law-Ogbomo KE, Osaigbovo AU. Effects of poultry manure on the growth and
372 herbage yield of amaranth (*amaranthus cruentus* L.). *Nigerian Journal of Agriculture, Food and*
373 *Environment*. 2012; 8(4):26-31.
- 374 12. Abdullahi J, Shaibu-Imodagbe EM, Mohammed F, Said A, Idris UD. Rural – Urban migration of
375 the Nigerian work populace and climate change effects on food supply. A case study of Kaduna
376 city in northern Nigeria. 2009.
- 377 13. Masarirambi MT, Mbokazi MB, Wahame PK, Oseni TO. Effect of kraal manure, chicken manure
378 and inorganic fertilizer on growth and yield of lettuce (*Lactuca sativa* L. var Commander) in a
379 semi-arid environment. *Asian Journal of Agricultural Sciences*. 2012; 4(1):56-64.
- 380 14. Burt R. Soil survey field and laboratory methods manual. *Soil survey investigations report*. 2014;
381 2(51):181-387.
- 382 15. Musa A, Ogbadoyi EO, Oladiran JA, Ezenwa MIS, Akanya HO. Effect of fruiting on
383 micronutrients, antinutrients and toxic substances in *Tilfairia occidentalis* grown in Minna, Niger
384 State, Nigeria. *African Journal of Environmental Science and Technology*. 2011; 5(9):710-716
- 385 16. Msibi BM, Mukabwe WO, Manyatsi AM, Mhazo N, Masarirambi MT. Effect of liquid manure on
386 growth and yield of spinach (*Beta vulgaris* var *Cicla*) in a sub-tropical environment in Swaziland.
387 *Asian Journal of Agricultural Sciences*. 2014; 6(2):40-47.
- 388 17. Egharevba RKA, Ogbe FM. The effects of different levels of organic and mineral fertilizers on the
389 yield performance of two *Amaranthus* (*A. cruentus*) cultivars. *The Plant Scientists*. 2002; 3:62-72.
- 390 18. Okokoh SJ, Bisong BW. Effect of Poultry Manure and Urea- N on Flowering Occurrence and
391 Leaf Productivity of *Amaranthus cruentus* in Calabar. *Journ.of Apl. Sic. Environmental Management*.
392 2011; 15(1):13-15.
- 393 19. Abayomi OA, Adebayo OJ. Effect of fertilizer types on the growth and yield of *Amaranthus*
394 *caudatus* in Ilorin, Sothern Guinea, Savanna Zone of Nigeria. *Advances in Agriculture*. 2014; 14:5
- 395 20. Islam MM, Karim AJM, Jahiruddin M, Majid NM, Miah MG, Ahmed MM, Hakim MA. Effects of
396 organic manure and chemical fertilizers on crops in the radish-stem amaranth-indian spinach
397 cropping pattern in homestead area. *Australian journal of crop science*. 2011; 5(11):1370-1378.
- 398 21. Saeed IN, Abbasi K, Kazim M. Response of maize (*Zea mays*) to nitrogen and phosphorus
399 fertilization under agro-climatic condition of Rawalokot Azad Jammu and Kashmir. *Pak. J. Biol.*
400 *Sci*. 2001; 4:53-55.
- 401 22. Law-Ogbomo KA, Ajayi SO. Growth and yield performance of *Amaranthus cruentus* influenced by
402 planting density and poultry manure application. *Natulae Botanicae Horti Agrobotanici Cluj-*
403 *Napoca*. 2009; 37(1): 195-199.
- 404 23. Sanwal SK, Lakminaragana K, Yadav RK, Yadav DS, Mousumi B. Effect of organic manures on
405 soil fertility, growth, physiology, yield and quality of turmeric. *Indian Journal of Horticulture*. 2007;
406 64 (4): 444-449.
- 407 24. Premsekhar M, Rajashree V. Influence of organic manures on growth, yield and quality of okra.
408 *American-Eurasian Journal of sustainable Agriculture*. 2009; 3(1): 6-8.
- 409 25. Jacobs RD, Loan D, Jacob J. Cage Layer Manure: An Important Resource for Land use. 2003.
410 Retrieved from: <http://edis.ifas.ufl.edu/PS005>, (Accessed on: May 19, 2009).
- 411 26. Mshelia JS, Degri MM. Effect of different levels of poultry manure on the performance of
412 *Amaranthus caudatus* L. in Bama, Nigeria. *International Journal of Science and Nature*. 2014;
413 5(1):121-125
- 414 27. Ehigiator JO. Farm yard manure: Need for its adoption as an alternative to chemical fertilizer uses
415 in Nigeria. *Nigerian Journal of Horticultural Science*. 1990; 3: 1– 9.
- 416 28. Van-Averbeke W, Yoganathan S. Using Kraal Manure as Fertilizer. Department of Agriculture
417 Cape Town, University of Fort Hare, Republic of South Africa. 2003.
- 418 29. Pimentel D, Hepperly P, Hanson J, Douds D, Sidel R. Environmental, energetic and economic
419 comparisons of organic and conventional farming systems. *BioSci*. 2005; 55(7): 573-582.

420
421
422
423
424

30. Kuntashula E, Shileshi G, Mafongoya PL, Banda J. Farmer participatory evaluation of the potential for organic vegetable production in the wetlands of Zambia. Outlook Agr. 2006; 35(4): 299-305.

UNDER PEER REVIEW