

Effect of Planting Date on the Performance of High Yield Potential Varieties of Rice in Bangladesh

ABSTRACT

The experiment was conducted at the Agronomy Field Bangladesh Agricultural University, Bangladesh to investigate the effect of planting dates on the performance of high yield potential varieties of rice in Boro season. The experiment consisted of five dates of transplanting viz. 10 and 25 December 2015, 10 and 25 January and 10 February 2016: four high yield potential varieties viz. hybrid rice Sonar Bangla-1, Jagoran, BRRI Dhan-29 and BINA Dhan-6. Results indicated that there was a significant effect of date of transplanting on yield of potential varieties of rice in Boro season. It was observed that plant height, panicle length, grain yield, straw yield, number of tillers per hill⁻¹ gradually increased up to 10 January. After 10 January transplanting yield reduced. The highest grain yield 6.41 t ha⁻¹ was observed in hybrid rice Sonar Bangla-1. The result revealed that Sonar Bangla-1 emerged as the best variety in Boro season regarding grain and straw yields among the varieties studied and it should preferably be transplanted between 25 December to 10 January to obtain appreciable better yield.

12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

Keywords: Hybrid rice, planting date, growth, yield

1. INTRODUCTION

Agriculture is the heart rhythm of Bangladesh which provides the ultimate entity of most of her population. The pressing issue of alarming population growth, rapid industrialization and urbanization has emerged as the biggest challenge for Bangladesh to ensure food security and socio-economic development. The economy of Bangladesh depends predominantly on agriculture sector accounting for about 31.08% of its gross domestic product [1]. Rice is grown in 11.79 million hectare of land with total production of 33.80 million tons [2]. Rice is extensively grown in Bangladesh in three season's viz. Aus, Aman and Boro and it covers 74.85 % of the total cultivable area of Bangladesh [2].

FAO (Food and Agriculture Organization) considers hybrid rice technology as an important avenue for increasing global rice production. China's success with hybrid rice encourages the prospect of this technology for the tropics and subtropics. Realizing hybrid rice technology as an important option to increase rice yield, IRRI rejuvenated its research on its development and Bangladesh is no exception to it as it is the prime need to increase rice production in Bangladesh.

Planting date is an important factor for obtaining higher yields and there is an optimum planting date to obtain higher yield of a crop [3]. Generally Boro rice is transplanted from early December to mid-March [4, 5]. Early transplantation of Boro rice prolongs field duration due to low temperature and involves high cost of production, particularly for management

36 practices including irrigation while delayed planting reduces the yield in some cases [6, 7]. A
37 compromise is therefore, needed between sacrificing grain yield by adjusting planting date or
38 incurring extra expenditure by irrigating the crop for a longer period in case of early planting.
39 Yield is the cumulative effect of the inherent characteristics of a variety as well as
40 management practices under which it is grown. Variety is one of the important factors for
41 increasing yield. In general, it is believed that there are differences in morpho-physiological
42 aspects among the traditional and modern varieties. Generally speaking modern **inbred** rice
43 varieties in Bangladesh have a longer growth duration of (150-160 days in Boro season) with
44 a low daily yield (lower than 30kg/ha/day) while the hybrid one because of its hybrid vigor
45 needs only 120-130 days to mature. If hybrid rice is cultivated, 20-40 days in crop duration
46 can be reduced. This may **facilitate** the accommodation of succeeding crop in the cropping
47 systems.

48 Boro rice has been gaining much importance by the farmers for its higher yield per hectare
49 than other rice crops. Recently several private seed traders are introducing rice seeds from
50 India and China. But as introduced plant materials, they need thorough evaluation under the
51 prevailing climatic conditions of Bangladesh for morphological and physiological
52 characteristic before they are going to the end users i.e., for large scale cultivation by the
53 farmers. As per available information regarding the yield and yield contributing characters,
54 both morphological and physiological characteristics of hybrid rice varieties are meager in
55 Bangladesh. That is why, it is a prime need to conduct more research work to find out and
56 develop sustainable technology of hybrid rice cultivation under the prevailing local edaphic
57 conditions. Therefore present study was undertaken to investigate the adaptability of high
58 yield potential rice varieties in boro season in relation to the effect of different planting dates
59 on higher growth and yield.

60

61 **2. MATERIAL AND METHODS**

62

63 **2.1 Experimental site**

64

65 The experiment was conducted at the Agronomy field, Bangladesh Agricultural University,
66 Bangladesh from November 2015 to June 2016 to study the effect of different planting date
67 on the performance of high yield potential varieties of rice in Boro season. The experimental
68 site is under the Old Brahmaputra Flood Plain of Agro-ecological Zone, AEZ 9 [8]. The land
69 was medium high with sandy loam in texture with pH value of 6.9. The experimental site was
70 under the subtropical climatic condition.

71

72 **2.2 Planting material**

73

74 Two inbred rice varieties (BRRI Dhan29 and BINA Dhan 6) and two hybrid rice (Sonar
75 bangla-1 and Jagoran) used as the test crops. Seeds were collected from BRRI, BINA for
76 the inbred varieties. Meanwhile Sonar Bangla-1 was imported from China and approved by
77 the National Seed Board Bangladesh. Jagoran was developed by India and marketed in
78 Bangladesh by BARC, Seed Ltd. Bangladesh.

79

80 **2.3 Experimental Design and Treatments**

81

82 The experiment was laid out in a Randomized Complete Block Design (RCBD) with three
83 replications. The treatments were comprised of two factors- Factor A: Variety viz. i) BRRI
84 Dhan29, ii) BINA Dhan6, iii) Sonar Bangla-1, iv) Jagoran and Factor B: Planting Date viz., i)
85 10 December 2015, ii) 25 December 2015, iii) 10 January 2016, iv) 25 January 2016 and v)
86 10 February 2016. There were 60 unit plots in the experiment. The size of the each plot was
87 5 m² where treatments were allotted at random.

88

2.4 Growth condition and measurement of parameters

Seeds of each cultivar were soaked in water in separate buckets for 24 hours and then placed under gunny bag for sprouting. After sprouting seeds were sown after 72 hours in the well prepared nursery bed. The 30 day old seedlings were then transplanted in the main field as per planting dates. Fertilizers were applied at the rate of 270, 130, 120, 70, and 10 kg/ha in the form of Urea, Triple Super Phosphate, Murate of Potash and Zinc Sulphate respectively following standard application procedures. Intercultural operations were done as per requirement. Maturity of crops was determined when some 90% of seeds became golden yellow color. The harvest crops were threshed manually and the fresh weights of grain and straw were recorded plot-wise. Finally, grain and straw yield per plot were converted to t/ha.

2.5 Data collection and Statistical Analysis

Data on individual plant parameters were recorded from the sample hill¹ and those on grain yield, straw yield, biological yield and harvest index were recorded from the whole plot at harvest. The collected data were analyzed statistically by MSTAT C software. The mean differences among the treatments were adjudged with Duncan's Multiple Range Test (DMRT) at 5% level of probability [9].

3. RESULTS AND DISCUSSION

3.1 Plant height

Plant height differed significantly among the varieties (Table 1). Results showed that BINA Dhan6 produced the tallest plants of 99.26 cm while Sonar Bangla-1 produced the shortest plant stature of 84.39 cm which was statistically identical to that produced by BRR1 Dhan29. Significant variation among the varieties was also reported by [10,11] and our results are in consistent with their result. This variation in plant height is probably due to the genetical makeup of the varieties.

Date of planting also exerted significant effect on plant height (Table 1). It was observed that the longest plant height (95.10 cm) when transplanted on 10 January. Plant height gradually decreased with delay in planting after 10 January. The shortest plant (85.47 cm) was observed when transplanted on 10 December.

Plant height was significantly affected by the combined effect of variety and planting time (Table 2). The tallest plant height was recorded in BINA Dhan6, transplanted on 10 January. The shortest plant height was observed in BRR1 Dhan29, transplanted on 10 December which was statistically similar to that of Sonar Bangla-1, transplanted on 10 December.

3.2 Number of effective tillers hill⁻¹

Number of effective tillers per hill varied significantly due to variety (Table 1). The results showed that the highest number (11.63) of effective tillers per hill was produced by BRR1 Dhan29. Number of effective tillers per hill varied significantly for different date of planting (Table1). It was observed that the highest number of effective tillers per hill (11.83) was produced on 10 January transplanting. On the other hand, lowest number of effective tillers per hill (8.46) was produced on 10 February planting. Effect of interaction of variety and planting date on the number of effective tillers per hill (Table 2). Maximum number of effective tillers per hill (15.73) was produced by BRR1 Dhan29 planted on 10 January. Meanwhile the lowest number of effective tillers per hill (6.39) was produced from 10 February planting in hybrid rice Jagoran.

142 **3.3 Number of non-effective tillers hill⁻¹**

143

144 The results showed that there were significant differences in the number of non-bearing
 145 tillers hill⁻¹ among the varieties studied (Table 1). The results revealed that the number of
 146 non-effective tillers hill⁻¹ ranged from 1.04 to 1.30. It was found that the highest number of
 147 non-effective tillers hill⁻¹ was produced by hybrid rice Jagoran whereas the lowest number of
 148 non-effective tillers hill⁻¹ was produced by BRRI Dhan29. This might be due to the genetic
 149 makeup of the varieties. This was reported by [12, 13]. Production of non-effective tillers per
 150 hill was found to be significantly affected by the date of planting (Table 1). A maximum
 151 number of non-effective tillers per hill (2.03) were produced when planting was done on 10
 152 February whereas production of non-effective tillers was minimum when transplanted on 10
 153 January. Number of non-effective tillers per hill was significantly affected by the interaction
 154 between variety and planting date (Table 2). Maximum number of non-effective tillers per hill
 155 (2.04) was produced by Sonar Bangla-1 transplanted on 10 February.

156

157 **3.4 Length of Panicle**

158

159 Length of panicle was not significantly affected by the variety and the planting date (Table 1).
 160 Effect of interaction of variety and planting date was found to be insignificant on panicle
 161 length (Table 2).

162

163 **Table1: Effect of variety and date of planting on the crop characters of rice in Boro**
 164 **season**

165

Treatment	Plant height (cm)	Effective tillers hill ⁻¹ (no)	Non-effective tillers hill ⁻¹ (no)	Length of panicle (cm)
Variety				
BRRI Dhan-29	86.03b	11.63a	1.04b	22.19
BINADhan-6	99.26a	9.19c	1.16ab	22.06
Sonar Bangla-1	84.39b	9.96b	1.10ab	20.84
Jagoran	97.19a	8.07d	1.30a	22.33
Significance level	0.01	0.01	0.05	NS
Transplanting date				
10 December	85.47b	9.49b	1.23b	22.00
25 December	92.48a	9.73b	0.83c	22.19
10 January	95.01a	11.83a	0.63c	22.53
25 January	92.63a	9.03bc	1.04b	21.62
10 February	93.33a	8.46c	2.03a	20.95
Significance level	0.01	0.01	0.05	NS
CV (%)	5.27	10.20	15.13	9.76

166

167

168

169

170

171 **Table 2: Combined effect of variety and date of planting on crop characters of rice in**
 172 **Boro season**

173

Treatment	Plant height (cm)	No. of total tillers hill ⁻¹	No. of Effective tillers hill ⁻¹	No. of Non-Effective tillers hill ⁻¹	Length of panicle(cm)
V ₁ D ₁	77.01e	11.66cd	11.00bcd	1.20efg	22.63

V ₁ D ₂	88.45cd	13.27b	11.20bc	0.66hij	22.98
V ₁ D ₃	90.56c	16.93a	15.73a	0.66hij	23.41
V ₁ D ₄	85.49cd	10.86cde	10.27bcde	0.66hij	21.36
V ₁ D ₅	89.99c	10.26defg	9.93bcdef	2.06ab	20.60
V ₂ D ₁	96.68b	10.40def	9.13defgh	1.40def	22.12
V ₂ D ₂	98.55b	10.93cde	9.20defgh	0.60hij	22.24
V ₂ D ₃	105.60a	11.20cde	9.80bcdef	0.40j	22.24
V ₂ D ₄	97.99b	9.60efgh	9.00efgh	1.60cde	21.96
V ₂ D ₅	97.48b	9.60efgh	8.80efgh	1.80bcd	21.77
V ₃ D ₁	77.87e	11.06cde	10.26bcde	0.93fghi	20.97
V ₃ D ₂	85.03cd	12.00bcd	10.26bcde	0.80fghi	20.98
V ₃ D ₃	84.26d	12.53bc	11.60b	0.53ij	21.26
V ₃ D ₄	88.26cd	10.80cde	9.46cdefg	0.86ghij	20.91
V ₃ D ₅	86.56cd	9.06fgh	8.20fghi	2.40a	20.10
V ₄ D ₁	90.32c	9.47efgh	7.60ghi	1.40cdef	22.29
V ₄ D ₂	97.92b	9.66efgh	8.26fghi	1.26fgh	22.58
V ₄ D ₃	99.65b	11.13cde	10.20bcde	0.93fghi	23.20
V ₄ D ₄	98.80 b	8.66gh	7.00hi	1.06fgh	22.24
V ₄ D ₅	99.30b	8.00h	6.93i	1.86fgh	21.33
Significance level	0.05	0.01	0.05	0.01	NS
CV (%)	5.27	8.31	10.20	15.13	9.76

174 Means in a same column followed by different letter (s) are significantly different at $P < 0.05$; (V₁- Sonar
175 Bangla-1, V₂- Jagoran; V₃- BRRJ Dhan-29; V₄- BINA Dhan-6 and D₁ -10 December, D₂ -25 December,
176 D₃ -10 January, D₄ -25 January; D₅ -10 February)

177

178 3.5 Grain weight hill⁻¹

179

180 Variety exerted significant influence on grain weight hill⁻¹ (Table 3). The highest grain weight
181 hill⁻¹ (26.31 g) was recorded in hybrid rice Jagoran whereas BINA Dhan6 gave the lowest
182 grain weight hill⁻¹ (21.86 g). The grain weight hill⁻¹ also varied due to different dates of
183 planting (Table 3). The highest weight (25.58 g) grains hill⁻¹ was produced from 10 January
184 whereas the lowest was recorded at 10 February planting. The variation due to interaction
185 between variety and planting date was significant for the parameter grain weight hill⁻¹ (Table
186 4). Maximum grain weight hill⁻¹ (27.34 g) was observed in Jagoran on 10 January planting
187 whereas minimum was observed from BRRJ Dhan29 on 10 February planting.

188

189 3.6 Weight of 1000-grains

190


191 Varieties differed significantly among themselves regarding weight of 1000-grains (Table 3).
192 The results revealed that highest 1000-grain weight (30.18 g) was obtained from Sonar
193 Bangla-1. The lowest 1000-grain weight (22.68 g) was found in BRRJ Dhan-29. This result is
194 in corroborate with the results of reported by [14, 15] who stated that 1000-grain weight
195 differed among the varieties. 1000-grain weight varied significantly due to different dates of
196 planting (Table 3). The highest grain weight was produced from 10 January planting that was
197 statistically at par with 25 December planting. The lowest 1000-grain weight was observed
198 from 10 February planting. However 1000-grain weight did not vary significantly due to the
199 interaction between variety and date of planting (Table 4).

200

201 3.7 Grain yield (t ha⁻¹)

202


203 Grain yield varied significantly among the varieties (Fig 1). The results elicited that hybrid
 204 rice variety Sonar Bangla-1 produced the maximum grain yield (6.41 t ha^{-1}) which was
 205 statistically alike to that of hybrid variety Jagoran (6.11 t ha^{-1}). On the other hand BINA
 206 Dhan6 produced the minimum grain yield (4.9 t ha^{-1}). The highest grain yield of hybrid rice
 207 Sonar Bangla-1 was the consequence of the maximum 1000-grains weight and the second
 208 highest number of productive tillers hill^{-1} . Though Jagoran produced the highest grain weight
 209 and second highest 1000-grains weight but failed to produce the highest yield because of
 210 mainly due to the minimum number of productive tillers hill^{-1} . This was also supported by [16,
 211 17, 18, 19].
 212
 213



214 **Figure 1: Effect of variety on the grain yield and straw yield of rice in Boro season**

215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225

Grain yield varied significantly due to different dates of planting (Fig 2). The highest grain
 yield (6.61 t ha^{-1}) was obtained at 10 January planting. The lowest (5.02 t ha^{-1}) yield was
 recorded on 10 February planting. The highest grain yield on 10 January might be due to the
 prevailing favorable temperature. The grain yield was significantly influenced by the
 interaction of variety and date of planting (Table 4). The highest grain yield (6.99 t ha^{-1}) was
 produced from the combination of Sonar Bangla-1 with 10 January Planted while the lowest
 (4.21 t ha^{-1}) was in BINA Dhan6 with planted on 10 February.



226

227

Figure 2: Effect of planting dates on the grain yield and straw yield of rice in boro season

228

229

230

3.8 Straw yield

231

232

233

234

235

236

237

238

239

240

241

242

3.9 Biological yield

243

244

245

246

247

248

249

250

251

252

253

254

255

The varietal effect on biological yield was highly significant (Table 3). The highest biological yield (13.76 t ha⁻¹) was recorded from Sonar Bangla-1 which has the highest grain yield producer. The biological yield of Jagoran and BRR1 Dhan29 was statistically similar to the hybrid rice Sonar Bangla-1. The lowest biological yield (11.10 t ha⁻¹) was produced by BINA Dhan6. Biological yield was also significantly affected by the date of transplating (Table 3). It was observed that the highest biological yield (14.12 t ha⁻¹) was obtained in 10 January planting. Meanwhile the lowest biological yield (11.34 t ha⁻¹) was obtained in 10 February planting. The result is in agreement with the findings of [20].

Effect of interaction of variety and date of planting was significant (Table 4). The highest biological yield (14.57 t ha⁻¹) was observed in interaction at Jagoran and on 10 January planting. The lowest biological yield was observed at BRR1 Dhan29 on 10 February planting.

256

257

258

259
260

Table 3: Effect of variety and planting date on crop characters of rice in boro season

Treatment	Grain weight hill ⁻¹ (g)	1000-grain weight (g)	Biological yield (t ha ⁻¹)	Harvest Index (%)
Variety				
BRR1 Dhan-29	22.68c	21.49b	13.32a	44.60ab
BINADhan-6	21.79c	26.02c	11.09b	43.60b
Sonar Bangla-1	24.39b	30.18a	13.75a	46.20a
Jagoran	26.31a	28.35b	13.37a	45.20ab
Level of Significance	0.01	0.01	0.01	NS
Transplanting date				
10 December	23.32b	26.28b	13.16bc	44.75abc
25 December	25.18a	27.01ab	13.36ab	44.75ab
10 January	25.58a	27.71a	14.10a	46.50a
25 January	23.87b	26.39b	12.44c	44.00bc
10 February	21.51c	25.17c	11.34d	43.50c
Significance level	0.01	0.01	0.01	0.05
CV (%)	5.81	4.21	7.11	5.09

261 *Means in a same column followed by different letter (s) are significantly different at P<0.05*

262
263
264
265

Table 4: Combined effect of variety and plating dates on crop characteristics of rice in Boro season

Treatment	Grain weight hill ⁻¹	1000-grain weight (g)	Grain Yield (t ha ⁻¹)	Straw Yield (t ha ⁻¹)	Biological Yield (t ha ⁻¹)	Harvest Index (%)
V ₁ D ₁	22.56 e-h	21.46	6.27 bcd	7.75 ab	14.02 ab	44.00
V ₁ D ₂	23.11 efg	22.03	6.55 abc	7.54 abc	14.09 ab	46.00
V ₁ D ₃	23.61 def	22.62	6.74 ab	7.52 abc	14.26 ab	47.00
V ₁ D ₄	22.08 e-h	21.34	6.18 b-e	7.90 a	14.08 ab	43.00
V ₁ D ₅	22.05 e-h	20.04	4.45 h	5.70 f	10.15 e	43.00
V ₂ D ₁	20.49 hi	26.00	4.81 gh	6.10 ef	10.91 de	44.00
V ₂ D ₂	24.57 b-e	26.71	5.09 fg	6.26 def	11.35 cde	44.00
V ₂ D ₃	24.63 b-e	26.76	5.96 cde	7.17 abc	13.13 ab	45.00
V ₂ D ₄	20.92gh	25.62	4.47 h	5.78 f	10.25 e	43.00
V ₂ D ₅	18.34 i	25.01	4.21 h	5.64 f	9.85 e	42.00
V ₃ D ₁	23.81 def	29.41	6.63 ab	7.60 abc	14.23 ab	46.00
V ₃ D ₂	26.06 a-d	30.77	6.69 ab	7.32 abc	14.31 ab	47.00
V ₃ D ₃	26.75 ab	32.04	6.99 a	7.52 abc	14.51 a	48.00
V ₃ D ₄	23.55 def	30.22	5.88 cde	7.01 abcd	12.89 abc	45.00
V ₃ D ₅	21.79 fgh	28.46	5.88 cde	6.96 bcd	12.84 abc	45.00
V ₄ D ₁	26.41 abc	28.23	6.16 bcde	7.33 abc	13.49 ab	45.00
V ₄ D ₂	26.98 ab	28.54	6.33 ab	7.38 abc	13.71 ab	46.00
V ₄ D ₃	27.34 a	29.43	6.76 ab	7.81 ab	14.57 a	46.00
V ₄ D ₄	26.96 ab	28.40	5.77 de	6.78 cde	12.55 bcd	45.00
V ₄ D ₅	23.89 c-f	27.71	5.56 ef	6.97 bcd	12.53 bcd	44.00
Significance level	0.05	NS	0.05	0.05	0.05	NS
CV (%)	5.81	4.21	6.01	6.82	7.11	5.09

266 Means in a same column followed by different letter (s) are significantly different at $P < 0.05$; (V_1 - Sonar
267 Bangla-1, V_2 - Jagoran; V_3 - BRRI Dhan-29; V_4 - BINA Dhan-6 and D_1 -10 December, D_2 -25 December,
268 D_3 -10 January, D_4 -25 January; D_5 -10 February)

269 3.10 Harvest index

270 Varieties exerted a significant effect on harvest index (Table 3). It is evident that the highest
271 harvest index (46.20%) was recorded from Sonar Bangla-1 which was statistically similar
272 with Jagoran (44.60%) and BRRI Dhan29 (45.20 %). The lowest harvest index (43.60%) was
273 obtained from BINA Dhan6 which led to the lowest grain yield (5.6 t ha^{-1}).

274 Harvest index also varied significantly due to date of transplanting (Table 3). The highest
275 harvest index (46.50 %) was observed in 10 January planting which statistically similar with
276 25 December and 10 December planting. The lowest harvest index (43.50 %) was obtained
277 in 10 February planting. However, interaction of variety and date of planting on harvest index
278 was non-significant.

279

280

281 4. CONCLUSION

282

283 The result revealed that there was significant effect of date of transplanting on yield of
284 potential varieties of rice in Boro season. The highest grain yield (6.41 t ha^{-1}) was obtained
285 from Sonar Bangla-1. Therefore, it may be concluded that Sonar Bangla-1 has emerged as the
286 best variety in Boro season regarding grain and straw yields among the varieties studied
287 under the present investigation and it should be preferably be transplanted between 25
288 December to 10 January to have higher yield.

289

290

291

292 REFERENCES

293

294

295 1. Biswas PK, Salokhe VM. Effects of planting date, intensity of tiller separation and plant
296 density on the yield of transplanted rice. J Agric Sci Cambridge, 2001; 137(3): 279–287

297 2. BBS (Bangladesh Bureau of Statistics). Statistics Division., Ministry of Planning, Govt.
298 Peoples Republic of Bangladesh, Dhaka. P. 56. 2017.

299 3. BRRI, Adunik Dhaner Chash (Modern Rice Cultivation), vol. 7, Bangladesh Rice Research
300 Institute, Joydebpur, Bangladesh, 6th edition, 2008, Booklet No. 5.

301 4. Julfquar AW, "BRRI: research and development of hybrid rice," Te Guardian, 2009, 19 (3):
302 33, Regd no. DA 816.

303 5. Pandey N, Verma AK, Tripathi RS. Effect of planting time and nitrogen on tillering pattern,
304 dry matter accumulation and grain yield of hybrid rice. Indian J Agric Sci. 2001; 71(5): 337–
305 338.

306 6. Akram HM, Ali A, Nadeem MA, Iqbal MS. Yield and yield components of rice varieties as
307 affected by transplanting dates. J Agric Res. 2007; 45(2): 105–111.

308 7. Zhende Y, 1988. Agronomic management of rice hybrids compared with conventional
309 varieties. International Rice research Institute, Manila, Philippines.

- 310 8. Khalifa AABA. Physiological evaluation of some hybrid rice varieties under different
311 sowing dates. Australian J. Crop. Sci.2009; 3: 178-183.
- 312 9. Gomez KA, Gomez AA. Statistical Procedures for Agricultural Research. 2nd ed. New
313 York: John Wiley and Sons.1984 p. 1–340
- 314 10. Kanfany G, El-Namaky R, Ndiaye K, Traore K and Ortiz R. Assessment of rice inbred
315 lines and hybrids under low fertilizer levels in Senegal. Sustainability.2014; 6: 1153-1162
- 316 11. Kushwaha UKS, Khatiwada SP and Upreti HK. Delayed transplanting of aged rice
317 seedlings causes the yield reduction in farmer's field. Genom. Appl. Biol.2016; 7: 1-9
- 318 12. Murthy KNK, Shankaranarayana V, Murali K and Jayakumar BV. Effect of different dates
319 of planting on spikelet sterility in rice genotypes (*Oryza sativa* L.). Res. Crops.2004; 5: 143-
320 147
- 321 13. Shah LM and Bhurer KP. Response of wet seeded rice varieties to sowing dates. Nepal
322 Agric. Res. J.2005; 6: 35-38
- 323 14. Swain P, Annie P and Rao KS. Evaluation of rice (*Oryza sativa*) hybrids in terms of
324 growth and physiological parameters and their relationship with yield under transplanted
325 condition. Indian J. Agric. Sci.2005; 76: 496-499
- 326 15. Abou-Khalif BAA. Evaluation of some hybrid rice varieties under different sowing times.
327 Afr J Plant Sci. 2009 ; 3(4): 53–58
- 328 16. Khakwani AA, Zubair M, Mansoor M, Naveed K, Shah IH, Wahab A, Ilyas M, Aahmed I.
329 Agronomic and morphological parameters of rice crop as affected by date of transplanting. J
330 Agron. 2006; 5(2): 248–250.
- 331 17. Choudhury DA, Hamid H, Miah GU and Haque MM, “Phenology, growth and yield ability
332 of modern and old rice cultivars of different maturity groups,” Bangladesh Agronomy
333 Journal,1998, 8: 47–52.
- 334 18. Peng S, Cassman KG, Virmani SS, Sheehy J and Khush GS, “Yield potential trends of
335 tropical rice since the release of IR8 and the challenge of increasing rice yield potential,”
336 Crop Science.1999; 39(6):1552–1559
- 337 19. Chowdhury PKD and Guha, B. Performance of rice varieties under different time of
338 planting in boro season. Annals of Biology Lundhiana.2000;16 (1): 41-44.
- 339 20. Iqbal S, Ahmad A, Hussain A, Ali MA, Khaliq T and Wajid SA. Influence of transplanting
340 date and nitrogen management on productivity of paddy cultivars under variable
341 environments. Int. J. Agric. Biol.2008; 10: 288-292.