

**HEPATOTHERAPEUTIC TENDENCY OF *CITRULLUS*
LANATUS RIND METHANOLIC EXTRACT ON LIVER
MARKERS IN MALE WISTAR RATS**

ABSTRACT

It is a common practice to discard the peel or rinds of fruits. Interestingly, some parts of fruit humans find inedible actually possess bioactive nutrients that may be used for medicinal purposes. The effect of methanolic extract of *Citrullus lanatus* rind on liver function in normal male wistar rats was studied. 24 wistar rats with body weight of 150-250g were used for this study. The animals were simple randomly divided into four groups, 6 rats in each. Group 1 contained the control given normal saline and feed; group 2, a low dose, 50mg/kg of methanolic extract of *Citrullus lanatus* rind was administered, group 3 and 4 were administered medium and high dose of *Citrullus lanatus* rind extract 100mg/kg and 200mg/kg respectively. The *Citrullus lanatus* rind extract was administered via orogastric route and the experiment lasted for a period of 56 days. Blood samples were collected by left ventricular cardiac puncture for liver function test at the last day of the experiment. The outcome of this research revealed that medium and high dose administration of *citrullus lanatus* rind significantly ($p \leq 0.05$) reduced the serum level of liver enzymes alanine transaminase, aspartate transaminase and alkaline phosphatase and also total protein. There was a non-significant ($p \leq 0.05$) change in serum total bilirubin and albumin when all doses were compared to the control. Prolonged and moderate ingestion of *Citrullus lanatus* rind may be of benefit in regulating blood level of liver enzymes; hence, this part of the fruit has therapeutic value.

KEY WORDS: *Citrullus lanatus*, Liver function, Methanolic extract, Transaminase

INTRODUCTION

The use of plants as source of medicines can never be underestimated^[1]. Plant application in medicine can be dated back to the days of ancient Egypt and beyond.^[2] The discovery of the bioactive phytoconstituents derived from various plants has provided indispensable knowledge concerning the therapeutic effect of various types of plants and their varying species^[3]. *Citrullus lanatus* is commonly called watermelon^[4]. It is a popular vegetable that is consumed globally for diverse reasons due to its nutritional equivalent. The rind of watermelon is usually green in color but may vary depending on the specie^[5]. The rinds make up the outer part of the fruit that cover the inner fleshy and commonly edible part usually composed of seeds^[6]. In countries like Nigeria and some other parts of western Africa, the only part regarded as edible is the reddish inner fleshy part of the fruit^[7]. Despite reports from other studies carried out generally, that revealed that the phytonutrient composition of the outer part or peel and seeds of most fruits are more abundant than the edible fleshy part^[8], most individuals still dispose the rind of watermelon because they believe it has no nutritional importance or better still it may be poisonous if ingested. There is always a belief that the rinds have no value nutritionally, economically and therapeutically.

Some tribes in Asia have already adopted it as a practice to prepare the rind as a special delicacy^[9]. The rind can be fried, boiled, roasted or consumed in raw form by some natives in Asia and Europe^[10]. The liver is a vital abdominal organ. Survival without the liver is very much impossible^[11]. Liver function tests are used to help detect, monitor or evaluate liver diseases or damages and recovery from such changes in response to various therapeutic methods or agents^[12]. Liver function tests include test for liver enzymes like alanine transaminase (ALT), alkaline phosphatase (ALP), aspartate transaminase (AST)^[10] and test for albumin and bilirubin^[13]. This study is aimed at evaluating the hepatotherapeutic potential of *Citrullus lanatus* rind.

MATERIALS AND METHODS

Plant and extract preparation

Large, fresh and healthy watermelons were purchased from Creek road market in Port Harcourt. The fruits were properly washed and the rinds were collected and extracted by maceration process for 48 hours using methanol. Qualitative phytochemistry and phytochemical analysis was carried out using standard laboratory techniques^[14] to determine the phytoconstituents or phytoactive agents present in the rind. The preparation of the plant extract was carried out in the Department of Phytochemistry and Pharmacognosy, Faculty of Pharmaceutical Sciences, Madonna University.

Extract preparation

The watermelon rind was collected. It was ensured that it was well peeled out, separating it from the fleshy red part of the fruit using a knife. The rinds were dried for 2 days in an incubator at 70°C. The rind appeared smaller in size after drying due to heat induced shrinkage. The dried rind was then grinded using a manual grinder. The powdered rind was then measured with a weighing balance. 50gram of the extract was introduced into 250ml of methanol using a measuring cylinder. The mixture containing 250ml methanol and 50gram of powdered extract was allowed to stay for about 2 days. After 2 days, the mixture was filtered using a filter paper and surgical gloves. The extract (filtrate) derived after the filtration process was dried for about 4 days in an incubator. The shaft (residue) derived from the filtration process was disposed. After about 4 days, the extract reduced in quantity due to the evaporation of methanol when exposed to heat in an incubator (Heat induced evaporation). The extract had a dark brown coloration. Concentration of the extract was 0.2mg/ml.

Phytochemical analysis

Test for alkaloids

0.2ml dilution of the extract was measured into a 250ml beaker and 50ml of 10% acetic acid in ethanol was added and allowed to stand for some minutes. This was filtered and the extract was concentrated on a water bath for one quarter of the original volume. Concentrated ammonium hydroxide was added drop wise to the extract until the precipitation was complete. The whole solution was allowed to settle and the precipitate was collected and washed with dilute ammonium hydroxide and then filtered. The residue is the alkaloid which was dried and weighed.

Test for saponins

0.2ml of the extract was added to warm water. The tube containing the extract and warm water was mixed properly and the presence of soapy lather signified the presence of saponins.

86 **Test for flavonoids**

87 The total flavonoid content was determined using a slightly modified method reported by
88 Minotti and Aust, the extract was measured into three test tubes in the range of 50, 100 and
89 200 ml and each was mixed with 500 ml of methanol. Water was added to mark up to 200 ml.
90 50ml 10% AlCl₃ followed by 50ml of 1M potassium acetate and 1400ml water was added
91 and allowed to incubate at room temperature for 30 minutes. The absorbance of the reaction
92 mixture was subsequently measured at 415nm; the total flavonoid content was subsequently
93 calculated. The non-flavonoid polyphenols were taken as the difference between the total
94 phenol and total flavonoid content.

95 **Test for tannins**

96 About 1ml of the methanol extract was added in 2ml of water in a test tube. 2 to 3 drops of
97 diluted ferric chloride solution was added and observed for green to blue-green (Cathechic
98 tannins) or a blue-black (Gallic tannins) coloration.

99 **Test for phenols**

100 The total phenol content was determined according to the method of singleton^[20]. Briefly,
101 appropriate dilution of the extracts were oxidized with 2.5ml of 10% Foli-Ciocalteau's
102 reagent (v/v) and neutralized by 2.0ml of 7.5% sodium carbonate to pH 7.4. The reaction
103 mixture was incubated for 40 minutes at 45°C and the absorbance was read at 765nm in the
104 spectrophotometer. The total phenol content was subsequently calculated as Gallic acid
105 equivalent.

106 **Test for steroids**

107 Sterols and Steroids were sought by the reaction of Liebermann^[24]. Ten (10ml) ml of
108 methanolic extract was evaporated. The residue was dissolved in 0.5ml of hot acetic
109 anhydride; we added 0.5ml of the filtrate chloroform Treated with the reagent of Libermann
110 Burchardt. The appearance, at the interphase, a ring of blue-green, showed a positive
111 reaction.

112 **Test for terpenoids**

113 With CHCl₃ (3ml), dissolve the extract (same 3ml), include H₂SO₄ (conc. 2ml) after drying.
114 For 2 minutes, allow to heat. Terpenoids indicted by solution that is gray.

115 **Test for cardiac glycosides**

116 Added about 2ml of HCL (dilute), to the extract (same 2ml) then pyridine (containing
117 sodium-nitroprusside) and NaOH were included in the initial solution. Glycosides indicated
118 by the appearance of scarlet red or pinkish color solution^[21].

119 **Experimental animals and protocols**

120 Twenty four (24) adult male wistar rats weighing 150 to 250 grams were obtained from the
121 experimental animal unit, Department of Human Physiology, Madonna University. All
122 animals were physically healthy. Using simple random technique of sampling, the animals
123 were divided into four (4) groups containing six (10) rats per group. The animals were
124 allowed to acclimatize for 2 weeks before the start of the experiment which lasted for 42
125 days. All animals had access to food and water *ad libitum*. The cages were properly cleaned
126 twice daily to avoid coprophagy.

127 Group 1: Control

128 Group 2: Low dose of *citrullus lanatus* rind

129 Group 3: Medium dose of *citrullus lanatus* rind

130 Group 4: High dose of *citrullus lanatus* rind

131 **Extract treatment**

132 The LD₅₀ of methanolic extract was 1500mg/kg. Methanolic extract of watermelon rind was
133 administered in 3 doses;

134 Low dose : 50mg/kg

135 Medium dose : 100mg/kg

136 High dose : 200mg/kg

137 The route of administration was the orogastric route.

138 **Sacrifice and collection of blood samples**

139 Several hours after treatment on the last day (day 42) of the experimental period, the animals
140 were anesthetized using diethyl ether from sigma chemicals® and then they were placed in a
141 supine position after which 5ml of blood was collected from the left ventricular chamber
142 using a syringe. The blood samples were collected into well labeled heparinized bottles. All
143 samples were taken to the laboratory for hematology, Madonna University Teaching
144 Hospital, for analysis of liver function enzymes and substrates.

145 **LIVER FUNCTION TESTS**

146 Experiment to determine the liver function biomarkers were carried out using the standard
147 laboratory procedures^[14]. The biomarkers tested for include; alanine transaminase (ALT),
148 aspartate transaminase (AST), alkaline phosphatase (ALP), total proteins (TP), bilirubin and
149 albumin^[19]. This test was carried out on blood samples collected on day 42 of the
150 experimental period.

151 **Test for alanine transaminase (ALT) and aspartate transaminase (AST)**

152 Activities of serum Aspartate transaminase (AST) and Alanine transaminase (ALT) were
153 assayed by the reitman and frankel calorimetric method^[15] in which 0.2 ml of serum reacted
154 with 1ml of substrate (Aspartate and α -ketoglutarate for AST, while alanine and α -
155 ketoglutarate for ALT, in phosphate buffer pH 7.4) and was incubated for an hour in the case
156 of AST and 30 minutes for ALT. Then 1ml of DNPH (Dinitrophenyl-hydrazine) solution
157 was added to arrest the reaction and kept for 20 minutes in room temperature. After
158 incubation, 1 ml of 0.4 N NaOH was added and absorbance was read at wavelength of
159 540nm.

160 **Test for alkaline phosphatase (ALP)**

161 Alkaline phosphatase in serum is determined by measuring the rate of hydrolysis of various
162 phosphate esters under specified condition. The principle in the test includes;

163 ALP

164 ρ - Nitro phenyl Phosphate + H₂O -----> ρ -Nitro phenol + H₃PO₄

165 P-Nitro phenyl Phosphate is hydrolyzed to ρ -Nitro phenol and inorganic phosphate. The rate
166 at which the ρ -Nitro phenol Phosphate is hydrolyzed, measured at 405nm, is directly
167 proportional to the alkaline phosphatase activity.

168 **Test for total protein (TP)**

169 The assay is based on a polypeptide chelation of cupric ion (colored chelate) in strong alkali.
170 In general, biuret assays are useful for samples containing -1 to 10 mg protein/ml, which is
171 diluted -5-fold by the added reagent to give a concentration of 0.2 to 2 mg/ml final assay
172 volume (F.A.V.). Most proteins produce a deep purple color, with a maximum absorbance
173 (λ_{max}) at about 550nm.

174 **Test for bilirubin**

175 Method of estimation of bilirubin in serum was based on an indirect reaction method of Van
176 den Berg: the bilirubin in serum reacted with a freshly prepared solution of Van den Berg's
177 diazotized sulphonilic acid (0.5 ml). Afterwards, purple colored azobilirubin compound was
178 formed which was measured at a wavelength of 540nm. This color was observed after the
179 addition of methanol and serum was diluted with distilled water, (0.2 ml + 1.8 ml distilled
180 water) (Klot, 2005).

181 **Test for albumin**

182 A bromocresol green (BCG) dye binding procedure was first proposed in 1964^[16]. This
183 procedure exhibited greater sensitivity and much lower susceptibility to interfering
184 substances^[15]. Albumin is bound by the BCG dye to produce an increase in the blue-green
185 color measured at 630nm. The color increase is proportional to the concentration of albumin
186 present.

187 **Ethical Approval**

188 This study was approved by Madonna University Research Ethics Committee. All
189 experimental procedures were done strictly following the guidelines provided by the research
190 ethics committee. The animals were sacrificed after exposure to diethyl ether according to EC
191 directives 86/609/EEC. In addition, the laid down standards according to the 1964 declaration
192 of Helsinki were strictly adhered to.

193 **Statistical Analysis**

194 SPSS 20.0 was used for all calculations and statistical analysis such as One-way analysis of
195 variance (ANOVA). Values are significant at $p \leq 0.05$ or at confidence interval of 95%.

196 **RESULTS**

197 **Table 1: Phytochemical constituents of methanolic extract of *Citrullus lanatus* rind.**

Phytochemicals	Indication
Saponin	—
Tannin	—
Flavonoids	++
Steroids	—

Alkaloids	++
Terpenoids	+
Phenol	+
Cardiac glycosides	+
Oils	+

+ = present - = Absent

Experimental data are presented in Mean±SEM. Percentage change was also calculated to make the data well translated.

Table 2:Effect of methanolic extract of *citrullus lanatus* rind on liver enzymes.

Treatments	ALT(U/L)	ALP(U/L)	AST (U/L)
Normal saline	1.1086±0.02	2.0874±0.17	39.3420±4.1
% Change	-0.02	-0.03	-2
50mg/kg of Extract	1.1083±0.01	2.0802±0.16	38.6000±33.5
% Change	-60	-50	-50
100mg/kg of Extract	0.476±0.02*	1.0498±0.41*	21.3350±148.3*
% Change	-70	-50	-50
200mg/kg of Extract	0.300±0.01*	1.0124±0.16*	21.0600±9.68*

Values are expressed in Mean±SEM, n=6, * $p \leq 0.05$ compared to control




Fig. 1; segmented pie chart illustration of the effect of treatments on alanine transaminase (ALT)




Fig. 2; segmented pie chart illustration of the effect of treatments on alkaline phosphatase (ALP)




Fig. 3; segmented pie chart illustration of the effect of treatments on aspartate transaminase (AST)

From **table 1**, the Phytoconstituents of methanolic extract of *Citrullus lanatus* (watermelon) rind include flavonoids, alkaloids, terpenoids, phenols and cardiac glycosides.

From **table 2**, the data showed that there was a significant decrease ($p \leq 0.05$) in the alanine transaminase (ALT) 100mg/kg medium dose ($0.476 \pm 0.02^*$) with a % change of -60 and 200mg/kg high dose ($0.300 \pm 0.01^*$) groups with a % change of -70 when compared with the control (1.1086 ± 0.02). There was no significant difference in the 50mg/kg low dose (1.1083 ± 0.01) with a % change of -2, compared to the control.

The result for alkaline phosphatase (ALP) showed that there was a significant decrease ($p \leq 0.05$) in the group 3 ($1.0498 \pm 0.41^*$) with a % change of -50 and group 4 ($1.0124 \pm 0.16^*$)

with a % change of -50 when compared with the control group (2.0874 ± 0.17). There was no significant difference in the group 2 (2.0802 ± 0.16) with a % change of **-0.03**, compared to the control..

The result for aspartate transaminase (AST) showed that there was a significant decrease ($p \leq 0.05$) in medium dose ($21.3350 \pm 148.3^*$) with a % change of -50 and in high dose ($21.0600 \pm 9.68^*$) with % change of -50 when compared to the control group (39.3420 ± 4.1). Low dose group (38.6000 ± 33.5) with a % change of -0.2 had no significant difference compared to the control.

Table 3: Effect of methanolic extract of *Citrullus lanatus* rind on liver substrates.

Treatments	TP($\mu\text{mol/L}$)	ALB($\mu\text{mol/L}$)	TB ($\mu\text{mol/L}$)
Normal saline	238.0468 ± 25.0	59.2660 ± 4.1	1.1370 ± 0.06
% Change	30	-0.1	20
50mg/kg of Extract	$298.4138 \pm 4.7^*$	58.4980 ± 3.8	1.3800 ± 0.13
% Change	50	-10	40
100mg/kg of Extract	$361.2048 \pm 23.3^*$	54.1712 ± 2.6	1.6177 ± 0.02
% Change	50	-10	-0.03
200mg/kg of Extract	$362.7020 \pm 5.2^*$	55.5514 ± 1.8	1.1340 ± 0.03

Values are expressed in Mean \pm SEM, n=6, * $p \leq 0.05$ compared to control




Fig. 4; segmented pie chart illustrating the effect of treatments on total protein (TP)




Fig. 5; segmented pie chart illustrating the effect of treatments on total bilirubin (TB)




Fig. 6; segmented pie chart illustrating the effect of treatments on Albumin (ALB)

249

250

251 From **table 3**, the result for total protein (TP) showed that there was a significant increase
252 ($p \leq 0.05$) in low dose ($298.4138 \pm 4.7^*$) with % change 30 , medium dose ($361.2048 \pm 23.3^*$)
253 with % change 50 and high dose ($362.7020 \pm 5.2^*$) with % change 50, when compared to the
254 control group ($238.0468 \pm 25.0^*$).

255 The result for albumin (ALB) showed that there was no significant change ($p \leq 0.05$) in all
256 doses; low dose (58.4980 ± 3.8), medium dose (54.1712 ± 2.6) and high dose
257 (55.5514 ± 1.8) when compared to the control group (59.2660 ± 4.1), at a % change of -0.1, -10
258 and -10 respectively.

259 The result for total bilirubin showed that there was no significant change ($p \leq 0.05$) in all
260 doses; low dose (1.3800 ± 0.13), medium dose (1.6177 ± 0.02) and high dose (1.1340 ± 0.03)
261 when compared to the control group (1.1370 ± 0.06), at a % change of 20, 40 and -0.03
262 respectively.

263 DISCUSSIONS

264 The phytoconstituents of methanolic extract of *Citrullus lanatus* rind is in correspondence
265 with earlier studies ^{[4][5][7]}. The phytoconstituents present in the methanolic extract of *Citrullus*
266 *lanatus* rind includes flavonoids, alkaloids, terpenoids, phenols and cardiac glycosides.
267 Methanolic form of extraction yielded better concentration of the phytochemicals probably
268 due to the non-polar biochemical nature of the various agents extracted for ^[4]. From previous
269 studies ^[4] ^{[5][7]}, Flavonoid is one of the major phytoconstituent in the methanolic extract of
270 *Citrullus lanatus* rind. Flavonoid has free radical scavenging properties ^[4]. *Citrullus lanatus*
271 rind is also an abundant source of lycopene, also a known antioxidant ^[2]. The antioxidant
272 agents in *Citrullus lanatus* may be the reason it has a dose-dependent hepatotherapeutic
273 function, hence, increasing the dose of *Citrullus lanatus* rind treatment may probably increase
274 its therapeutic manifestations in relation to liver function. The ALT, ALP, AST ^[15], Total
275 protein, Albumin, Total Bilirubin are the most sensitive biochemical markers employed in
276 the diagnosis of hepatic dysfunction ^[16]. Treatment with moderate and high doses of *citrullus*
277 *lanatus* rind methanolic extract resulted in significant reduction in serum levels of ALT, ALP
278 and AST. Low dose treatment of the extract caused no significant change in serum ALT,
279 ALP and AST in comparison with the control. The findings of this study is in agreement with
280 earlier reports ^{[3][7]}. Watermelon rind contains significant quantity of antioxidant
281 phytochemicals ^[4]. Alkaloids, flavonoids and phenols possess antioxidant properties ^[18].
282 Several studies have revealed the positive correlation between oxidative stress and cellular
283 damage ^[4] ^[18]. The *Citrullus lanatus* rind methanolic extract also significantly increased
284 serum level of total proteins (TP). This increase further reflects the ability for the extract to
285 enhance the synthetic function of the liver as well as its hepatoprotective function. It is
286 believed that flavonoid content in the rind extract may promote the synthesis of
287 immunocompetent proteins by the hepatocytes ^[3]. This may be the reason behind the
288 significant increase in total proteins after treatment with all doses of the extract. The total
289 bilirubin and albumin showed no significant change in all treatment doses. The *Citrullus*
290 *lanatus* rind methanolic extract may, at the level of this study, be of low potency in affecting
291 the rate of synthesis of albumin and secretion of bilirubin. Also the absence of saponins, a
292 reported hemolytic phytoagent, may probably reduce or maintain the blood level of
293 bilirubin ^[4].

Conclusion

Citrullus lanatus rind should be ingested as part of the fruit due to its therapeutic phytoconstituents. It has the tendency of being effective in management of defects in liver function.

Recommendation

Ethical approval

An ethical approval was obtained from Madonna University Research Ethics Committee.

This research work should be replicated on human subjects.

CONSENT

Not applicable in this study.

REFERENCES

- [1] Shaogui Guo, Jianguo Zhang, Honghe Sun, Jerome Salse, the draft genome of watermelon (*Citrullus lanatus*) and resequencing of 20 diverse accessions. *Nature Genetics.*; **2013** 45:51–58.
- [2] Naresh Singh Gill. Evaluation of Antioxidant activity of *citrullus lanatus* Seed extract in Rats. *Latin American journal of pharmacy* (formerly *Acta Farmaceutica Bonaerense*), *Lat. Am. J. Pharm.*; **2011**;30(3):429-34.
- [3] Madhavi P, Maruthi Rao, Kamala Vakati, Habibur Rahman, M. Chinna Eswaraiah, Evaluation of Anti-Inflammatory Activity of *Citrullus lanatus* Seed Oil by *In-vivo* and *In-vitro* Models. *Int. Res J Pharm. App Sci.*, **2012**; 2(4):104-108.
- [4] Alok Bhardwaj, Rajeev Kumar, Vivek Dabas, Niyaz Alam, Evaluation of Anti-Ulcer Activity of *Citrullus Lanatus* Seed Extract in Wistar Albino Rats. *International Journal of Pharmacy and Pharmaceutical Sciences*, **2012**; 4(5): 135-139.
- [5] Saxena, M., Saxena, J., Nema, R., Singh, D. and Gupta, A. Phytochemistry of medicinal plants. *Journal of Pharmacognosy and Phytochemistry*, 1(6): **2013**; 168-182.
- [6] Aruna Poduri, Debra L. Rateri, Shubin K. Saha, Sibin Saha, Alan Daugherty. *Citrullus lanatus* ‘sentinel’ (watermelon) extract reduces atherosclerosis in LDL receptor-deficient mice. *Journal of Nutritional Biochemistry*, **2012**;
- [7] Sevcan Altas, Göksel Kızıl, Murat Kızıl, Aydın Ketani, Parvez I. Haris, Protective effect of Diyarbakır watermelon juice on carbon tetrachloride-induced toxicity in rats. *Food and Chemical Toxicology*, **2011**; 49: 2433–2438

- 329 [8] Hoque MS, Uddin MF, Islam MA. A market model for watermelon with supply under
330 rational expectations: An empirical study on Bangladesh. *European Scientific*
331 *Journal*. **2015**; 11(9):236.
- 332 [9] Khanam M, Hafsa U. Market model analysis and forecasting behavior of watermelon
333 production in Bangladesh. *Bangladesh J. Sci. Res.* **2013**; 26(1&2):47-56.
- 334 [10] Imai, K.; Takai, K.; Watanabe, S.; Hanai, T.; Suetsugu, A.; Shiraki, M.; Shimizu, M.
335 Sarcopenia Impairs Prognosis of Patients with Hepatocellular Carcinoma: The Role of
336 Liver Functional Reserve and Tumor-Related Factors in Loss of Skeletal Muscle.
337 *Nutrients* **2017**, 9, 1054.
- 338 [11] Yoh, K.; Nishikawa, H.; Enomoto, H.; Ishii, A.; Iwata, Y.; Miyamoto, Y.; Ishii, N.; Yuri,
339 Y.; Hasegawa, K.; Nakano, C.; et al. Predictors Associated with Increase in Skeletal
340 Muscle Mass after Sustained Virological Response in Chronic Hepatitis C Treated
341 with Direct Acting Antivirals. *Nutrients* **2017**, 9, 1135.
- 342 [12] Yang, C.H.; Perumpail, B.J.; Yoo, E.R.; Ahmed, A.; Kerner, J.A., Jr. Nutritional Needs
343 and Support for Children with Chronic Liver Disease. *Nutrients* **2017**, 9, 1127.
- 344 [13] Marciano, F.; Savoia, M.; Vajro, P. Celiac disease-related hepatic injury: Insights into
345 associated conditions and underlying pathomechanisms. *Dig. Liver Dis.* **2016**, 48,
346 112–119.
- 347 [14] Orso, G.; Mandato, C.; Veropalumbo, C.; Cecchi, N.; Garzi, A.; Vajro, P. Pediatric
348 parenteral nutrition-associated liver disease and cholestasis: Novel advances in
349 pathomechanisms-based prevention and treatment. *Dig. Liver Dis.* **2016**, 48, 215–222.
- 350 [15] Tang, X.; Wei, R.; Deng, A.; Lei, T. Protective Effects of Ethanolic Extracts from
351 Artichoke, an Edible Herbal Medicine, against Acute Alcohol-Induced Liver Injury in
352 Mice. *Nutrients* **2017**, 9, 1000.
- 353 [16] Rangboo, V.; Noroozi, M.; Zavoshy, R.; Rezadoost, S.A.; Mohammadpoorasl, A. The
354 Effect of Artichoke Leaf Extract on Alanine Aminotransferase and Aspartate
355 Aminotransferase in the Patients with Nonalcoholic Steatohepatitis. *Int. J. Hepatol.*
356 **2016**.
- 357 [17] Mehmetçik, G.; Ozdemirler, G.; Koçak-Toker, N.; Cevikba,s, U.; Uysal, M. Effect of
358 pretreatment with artichoke extract on carbon tetrachloride-induced liver injury and
359 oxidative stress. *Exp. Toxicol. Pathol.* **2008**, 60, 475–480.
- 360 [18] Ilochi, O.N., Chuemere, A.N., Olorunfemi, O.J.; Evaluation of Antihyperglycaemic
361 Potential of *Allium cepa*, Coffee and Oxidative stress. *International Journal of*
362 *Biochemistry and Physiology* **2018**; Medwin Publishers, 3-1(pg1-9).
- 363 [19] Perumpail, B.J.; Li, A.A.; Cholaneril, G.; Kumari, R.; Ahmed, A. Optimizing the
364 Nutritional Support of Adult Patients in the Setting of Cirrhosis. *Nutrients* **2017**, 9,
365 1114.
- 366 [20] Huisman, E.J.; Trip, E.J.; Siersema, P.D.; van Hoek, B.; van Erpecum, K.J. Protein
367 energy malnutrition predicts complications in liver functions. *Eur. J. Gastroenterol.*
368 *Hepatol.* **2011**, 23, 982–989.

- 369 [21] Yadav, R. and Agarwala, M. Phytochemical analysis of some medicinal plants. *Journal*
370 *of Phytology*, **2011**; 3(12): 10-14.
- 371 [22] Tiwari, P., Kumar, B., Kaur, M., Kaur, G. and Kaur, H. Phytochemical screening and
372 extraction: A review. *Internationale Pharmaceutica Scientia*, **2011**; 1(1): 98-106.
- 373 [23] Rahman, H., Priyanka, P., Lavanya, T., Srilakshmi, N. and Kumar, P. R. (). A review on
374 the ethnobotany, phytochemistry and pharmacology of *Citrullus lanatus*.
375 *International Research Journal of Pharmaceutical and Applied Sciences*,
376 **2013**;3(2): 77-81.
- 377 [24] Tyagi, S., Singh, G., Sharma, A. and Aggarwal, G. Phytochemicals as candidate
378 therapeutics: An overview. *International Journal of Pharmaceutical Sciences*
379 *Review and Research*, **2012**; 3(1):53-55.